


META 2021 Warsaw - Poland

The 11th International Conference on Metamaterials, Photonic Crystals and Plasmonics



Program

July 20 – 23, 2021
Warsaw - Poland

metaconferences.org

META 2021 Warsaw - Poland

The 11th International Conference on Metamaterials, Photonic Crystals and Plasmonics

Program

**July 20 – 23, 2021
Warsaw - Poland**

metaconferences.org



META 2021 Warsaw - Poland

The 11th International Conference on Metamaterials, Photonic Crystals and Plasmonics

Please share your comments, photos & videos !

facebook

www.facebook.com/metaconference

twitter

[#meta2021 @metaconference](https://twitter.com/metaconference)

Edited by

Said Zouhdi | Paris-Saclay University, France
Dorota Pawlak | ENSEMBLE3 Centre of Excellence, Poland
Andrzej Kudelski | University of Warsaw, Poland

CONTENTS

| | |
|--------------------------------------|----|
| META 2021 ORGANIZATION | 5 |
| SPONSORS AND SUPPORTERS | 7 |
| PLENARY SPEAKERS | 8 |
| KEYNOTE SPEAKERS | 12 |
| CONFERENCE TUTORIALS | 15 |
| TECHNICAL PROGRAM | 19 |

META 2021 ORGANIZATION



Said Zouhdi
General Chair
Paris–Saclay University
France



Dorota Pawlak
General Co-Chair
ENSEMBLE3 Centre of Excellence
Poland



Andrzej Kudelski
General Co-Chair
University of Warsaw
Poland

LOCAL ORGANIZING COMMITTEE

Piotr Piotrowski, University of Warsaw, Poland
Emil Tymicki, ITME, Poland

Kasia Kołodziejak, ITME, Poland

INTERNATIONAL ADVISORY COMMITTEE

Federico Capasso, USA
Nader Engheta, USA
Teruya Ishihara, Japan
Yuri Kivshar, Australia
Graeme W. Milton, USA

Manuel Nieto-Vesperinas, Spain
Susumu Noda, Japan
Franco Nori, Japan & USA
Masaya Notomi, Japan
Vladimir Shalaev, USA

David R. Smith, USA
Martin Wegener, Germany
Xiang Zhang, Hong Kong
Nikolay Zheludev, UK

TECHNICAL PROGRAM COMMITTEE

Pierre-Michel Adam, France
Ishwar Aggarwal, USA
Benfeng Bai, China
Mohamed Bakr, Canada
Xavier Begaud, France
Henri Benisty, France
Svetlana V. Boriskina, USA
Che Ting Chan, Hong Kong
Debashis Chanda, USA
Pai-Yen Chen, USA
Johan Christensen, Spain
Alfredo De Rossi, France
Alexander Dmitriev, Sweden
Mohamed Farhat, Saudi Arabia

Monika Fleischer, Germany
Pedro D. G. Fernandez, Spain
Alexander Govorov, USA
Stephen Hanham, UK
Ortwin Hess, UK
Eugene Kamenetskii, Israel
Daniel Lanzillotti Kimura, France
Howard (Ho Wai) Lee, USA
Qiang Li, China
Yang Li, China
Anatole Lupu, France
Nicolò Maccaferri, Luxembourg
Martin McCall, UK
Miguel Navarro-Cia, UK

Taiichi Otsuji, Japan
Namkyoo Park, Korea
Virginie Ponsinet, France
Min Qiu, China
Ilya Razdolski, Poland
Jun Suk Rho, Korea
Almas Sadreev, Russia
Takuo Tanaka, Japan
Andrea Toma, Italy
Tetsuya Ueda, Japan
Paolo Vavassori, Spain
Hong Wei, China

SPECIAL SESSIONS ORGANIZERS

Antonio Ambrosio, Italy
Benfeng Bai, China
Mohamed Bakr, Canada
Henri Benisty, France
Jamal Berakdar, Germany
Mohamed B. Larosi, Spain
Shah Nawaz Burokur, France

Vincenzo Caligiuri, Italy
Debashis Chanda, USA
Xianzhong Chen, UK
Xianfeng Chen, China
Dibakar Roy Chowdhury, India
Jayasri Dontabhaktuni, India
Ramy El-Ganainy, USA

Didier Felbacq, France
Denis Garoli, Italy
Patrice Genevet, France
Jonathan Gratus, UK
Junpeng Guo, USA
Kotaro Kajikawa, Japan
Harshavardhan Kalathur, India

Paul Kinsler, UK
 Manas Kulkarni, India
 Stéphane Lanteri, France
 Yang Li, China
 Erik Lotfi, Bahrain
 Anatole Lupu, France
 Nicolò Maccaferri, Luxembourg
 Konstantinos Makris, Greece
 Martin McCall, UK
 Bumki Min, Korea
 Igor V. Minin, Russia

Oleg V. Minin, Russia
 Mario Miscuglio, USA
 Sang Soon Oh, UK
 Sahin Ozdemir, USA
 Willie Padilla, USA
 Dorota Pawlak, Poland
 Lucia Petti, Italy
 Virginie Ponsinet, France
 Juan Pou Saracho, Spain
 Ciyuan Qiu, China
 Badreddine Ratni, France

Emmanuel Rousseau, France
 Matthias Saba, Switzerland
 Sergey Saveliev, UK
 Jonghwa Shin, Korea
 Artur Sowa, Canada
 Junichi Takahara, Japan
 Hong Wei, China
 Jeong Weon Wu, Korea
 Fajun Xiao, China
 Alexandre Zagoskin, UK
 Weiren Zhu, China

SPECIAL SYMPOSIA ORGANIZERS



Jerome Plain
France



Alexander Govorov
USA



Davy Gérard
France



Pedro H. Martinez
Singapore



Junsuk Rho
Korea



Hakjoo Lee
Korea



Namkyoo Park
Korea



Howard Lee
Korea



Yu-Jung (Yuri) Lu
Taiwan



Zi Jing Wong
USA



Eugene Kamenetskii
Israel



Jensen Li
Hong Kong



Guoliang Huang
USA

SPONSORS AND SUPPORTERS

META 2021 gratefully acknowledges the support of these institutions and companies for their contribution to the success of this conference.

SUPPORTERS



PLENARY SPEAKERS



Robert W. Boyd

University of Ottawa, Canada and University of Rochester, USA

How Light Behaves when the Refractive Index Vanishes

Robert Boyd received the B.S. degree in physics from the Massachusetts Institute of Technology and the Ph.D. degree in physics in 1977 from the University of California at Berkeley. His Ph.D. thesis was supervised by Charles Townes and involved the use of nonlinear optical techniques in infrared detection for astronomy. Professor Boyd joined the faculty of the Institute of Optics of the University of Rochester in 1977 and in July 2001 he became the M. Parker Givens Professor of Optics. In 2010, he became Professor of Physics and Canada Excellence Research Chair in Quantum Nonlinear Optics at the University of Ottawa. His research interests include studies of nonlinear optical interactions, studies of the nonlinear optical properties of materials, the development of photonic devices including photonic biosensors, and studies of the quantum statistical properties of nonlinear optical interactions. Professor Boyd has written two books, co-edited two anthologies, published over 200 research papers, and has been awarded five patents. He is a fellow of the Optical Society of America and of the American Physical Society and is the past chair of the Division of Laser Science of the American Physical Society.



Federico Capasso

Harvard University, USA

Structuring Light and Dark with Metaoptics

Federico Capasso is the Robert Wallace Professor of Applied Physics at Harvard University, which he joined in 2003 after 27 years at Bell Labs where he was Member of Technical Staff, Department Head and Vice President for Physical Research. He is visiting professor at NTU with both the School of Physical and Mathematical Sciences and Electrical and Electronic Engineering. His research has focused on nanoscale science and technology encompassing a broad range of topics. He pioneered band-structure engineering of semiconductor nanostructures and devices, invented and first demonstrated the quantum cascade laser and investigated QED forces including the first measurement of a repulsive Casimir force. His most recent contributions are new plasmonic devices and flat optics based on metasurfaces. He is a member of the National Academy of Sciences, the National Academy of Engineering, the American Academy of Arts and Sciences. His awards include the King Faisal Prize, the IEEE Edison Medal, the SPIE Gold Medal, the American Physical Society Arthur Schawlow Prize in Laser Science, the Jan Czochralski Award for lifetime achievements in Materials Science, the IEEE Sarnoff Award in Electronics, the Materials Research Society Medal, the Wetherill Medal of the Franklin Institute, the Rank Prize in Optoelectronics, the Optical Society Wood Prize, the Berthold Leibinger Future Prize, the Julius Springer Prize in Applied Physics, the European Physical Society Quantum Electronics Prize.



Nader Engheta

University of Pennsylvania, USA

4D Structured Waves

Nader Engheta is the H. Nedwill Ramsey Professor at the University of Pennsylvania in Philadelphia, with affiliations in the Departments of Electrical and Systems Engineering, Materials Science and Engineering, Physics and Astronomy, and Bioengineering. He received his B.S. degree from the University of Tehran, and his M.S and Ph.D. degrees from Caltech. His current research activities span a broad range of areas including nanophotonics, metamaterials, nano-scale optics, graphene optics, optical metatronics, imaging and sensing inspired by eyes of animal species, optical nanoengineering, microwave and optical devices, and physics and engineering of fields and waves. He has received several awards for his research including the 2017 William Streifer Scientific Achievement Award from the IEEE Photonics Society, the 2015 Gold Medal from SPIE, the 2015 Fellow of US National Academy of Inventors (NAI), the 2015 National Security Science and Engineering Faculty Fellow (NSSEFF) Award (also known as Vannevar Bush Faculty Fellow Award) from US Department of Defense, the 2015 IEEE Antennas and Propagation Society Distinguished Achievement Award, the 2015 Wheatstone Lecture in King's College London, the 2014 Balthasar van der Pol Gold Medal from the International Union of Radio Science (URSI), the 2013 Inaugural SINA Award in Engineering, the 2012 IEEE Electromagnetics Award, 2006 Scientific American Magazine 50 Leaders in Science and Technology, the Guggenheim Fellowship, and the IEEE Third Millennium Medal. He is a Fellow of seven international scientific and technical societies, i.e., IEEE, URSI, OSA, APS, MRS, SPIE, and American Association for the Advancement of Science (AAAS). He has received the honorary doctoral degrees from the Aalto University in Finland in 2016 and from the University of Stuttgart, Germany in 2016.



Maiken H. Mikkelsen

Duke University, USA

Applications of metasurfaces : From multispectral imaging to optical communications and biosensing

Maiken H. Mikkelsen is the James N. and Elizabeth H. Barton Associate Professor at Duke University in the Department of Electrical and Computer Engineering, and by courtesy, in the Departments of Physics and Mechanical Engineering and Materials Science. She received her B.S. in Physics from the University of Copenhagen in 2004, her Ph.D. in Physics from the University of California, Santa Barbara in 2009 and was a postdoctoral fellow at the University of California, Berkeley before joining Duke University in 2012. Her research explores nanophotonics and new quantum materials to enable transformative breakthroughs for optoelectronics, quantum science, the environment and human health. Her awards include the Maria Goeppert Mayer Award from the American Physical Society, the NSF CAREER award, the Moore Inventor Fellow award from the Gordon and Betty Moore Foundation, Young Investigator Program Awards from the Office of Naval Research, the Army Research Office and the Air Force Office of Scientific Research, the Cottrell Scholar Award from the Research Corporation for Science Advancement, and the Early Career Achievement Award from SPIE – the International Society for Optics and Photonics.



Masaya Notomi

NTT Basic Research Labs., Japan

Integrated Nanophotonics for Optoelectronic Computation

Masaya Notomi received his B.E., M.E. and Ph.D. degrees in applied physics from The University of Tokyo, Japan in 1986, 1988, and 1997, respectively. He joined NTT Optoelectronics Laboratories, Nippon Telegraph and Telephone Corporation in 1988 and moved to NTT Basic Research Laboratories in 1999. Since then, his research interest has been to control the optical properties of materials and devices by using artificial nanostructures, and engaged in research on quantum wires/dots and photonic crystal structures. In 1996-1997, he was a visiting researcher of Linköping University, Sweden. He was a guest associate professor of Applied Electronics in 2003-2009 and is currently a guest professor of Physics in Tokyo Institute of Technology. He was appointed as Senior Distinguished Scientist of NTT since 2010. He is currently a director of NTT Nanophotonics Center. He received IEEE/LEOS Distinguished Lecturer Award in 2006, Japan Society for the Promotion of Science (JSPS) prize in 2009, Japan Academy Medal in 2009, the Commendation for Science and Technology by the Minister of Education, Culture, Sports, Science and Technology (Prize for Science and Technology, Research Category) in 2010, and IEEE Fellow grade in 2013. He served as a member of National University Corporation Evaluation Committee in the Japanese government. He is a research director of JST CREST program from 2015. He is also a member of the Japan Society of Applied Physics, APS, IEEE, and OSA.



Deirdre O'Carroll

Rutgers University, USA

Metasurfaces for Light Management in Semiconductor Thin Films

Deirdre O'Carroll is an Associate Professor in the Departments of Materials Science, Engineering and Chemistry and Chemical Biology at Rutgers University. Her research areas include nanophotonics, plasmonics, organic optoelectronics and energy materials. She obtained her B.E. in Electrical Engineering in 2002, and a PhD in Microelectronics in 2008 at University College Cork and the Tyndall National Institute, Ireland. Prior to joining Rutgers in 2011, she conducted postdoctoral research in plasmonics at California Institute of Technology in the US and at the University of Strasbourg and CNRS in France. She is a recipient of a National Science Foundation CAREER Award (2016), an American Chemical Society Young Investigator Award in Polymer Material Science and Engineering (2017) and a Science Foundation Ireland Future Research Leaders Award (2018). She is an associate editor for the SPIE Journal of Photonics for Energy and a member of the editorial advisory board for APL Photonics.



Vladimir M. Shalaev

Purdue University, USA

Empowering Quantum Photonics with Nanoplasmonics and Machine Learning

Vladimir M. Shalaev, Scientific Director for Nanophotonics at Birck Nanotechnology Center and Distinguished Professor of Electrical and Computer Engineering at Purdue University, specializes in nanophotonics, plasmonics, and optical metamaterials. Vladimir M. Shalaev has received several awards for his research in the field of nanophotonics and metamaterials, including the Max Born Award of the Optical Society of America for his pioneering contributions to the field of optical metamaterials, the Willis E. Lamb Award for Laser Science and Quantum Optics, IEEE Photonics Society William Streifer Scientific Achievement Award, Rolf Landauer medal of the ETOPIM (Electrical, Transport and Optical Properties of Inhomogeneous Media) International Association, the UNESCO Medal for the development of nanosciences and nanotechnologies, OSA and SPIE Goodman Book Writing Award. He is a Fellow of the IEEE, APS, SPIE, MRS and OSA. Prof. Shalaev has authored three books, thirty invited book chapters and over 500 research publications.

KEYNOTE SPEAKERS



Ali Adibi

Georgia Institute of Technology, USA

Analysis and Knowledge Discovery of Metastructures Using Deep Learning and Machine Learning Approaches in Reduced-dimensionality Spaces



Harry Atwater

California Institute of Technology, USA

Tunable and Time-Modulated Flat Optics



Konstantin Bliokh

RIKEN, Japan

A new spin for acoustics



Paul V. Braun

University of Illinois, USA

Volumetric microscale gradient refractive index lenses and waveguides for ultra-dense 3D optics



Hyuck Choo

Samsung Electronics Co., Ltd., Korea

Commercializing Metaphotonics



Miguel A. Correa-Duarte

University of Vigo, Spain

Photophysical Effects behind the Efficiency of Hot Electron Injection in Plasmon-Assisted Catalysis



Javier García de Abajo

ICFO-Institut de Ciències Fotòniques, Spain

Nanophotonics with Two-Dimensional Materials



Hilmi Volkan Demir

Nanyang Technological University (NTU), Singapore

Semiconductor Nanocrystal Optoelectronics : Pushing the Limits



Alexander Govorov

Ohio University, USA

Optical and photochemical properties of chiral plasmonic nanostructures



Deep Jariwala

University of Pennsylvania, US

Tunable Light-Matter Coupling in Low-Dimensional Excitonic Semiconductors



Mona Jarrahi

University of California Los Angeles, USA

Wavelength conversion through plasmonic photoconductive nanostructures



Seokwoo Jeon

Korea Advanced Institute of Science and Technology, Korea

Fast, Low Cost Fabrication of Optimized 3D Nanostructures for Energy Transfer and Transport Properties



Philippe Lalanne

Institut d'Optique Graduate School, France

Rigorous modal analysis of micro and nanoresonators



Howard Lee

UC Irvine, USA

Active Epsilon-near-zero Photonics



Prineha Narang

Harvard University, USA



Hiromi Okamoto

Institute for Molecular Science, Japan

Chiral Near-Field Properties of Plasmonic Nanomaterials : Imaging and Functions



Sir John B. Pendry

Imperial College London, UK

Metamaterials that travel faster than light : putting the squeeze on photons



Junsuk Rho

Pohang University of Science and Technology (POSTECH), Korea

Dielectric metasurfaces for flat optics : wavefront engineering and future applications



Volker J. Sorger

George Washington University, USA

Strainoptronics : A New Degree of Freedom for 2D Material Device Engineering



Martin Wegener

KIT, Germany

3D Laser Nanoprinting of 3D Metamaterials



Rachel Won

Nature Photonics (United Kingdom)

3D Publishing in Nature Journals



Eli Yablonovitch

UC Berkeley, USA

The Challenge of META is (Aperiodic) Inverse Electromagnetic Design

CONFERENCE TUTORIALS

META 2021 features several technical tutorials instructed by world-leading experts on various topics of interest to the META community. Tutorials are intended to provide a high quality learning experience to conference attendees. The tutorials address an audience with a varied range of interests and backgrounds : beginners, students, researchers, lecturers and representatives of companies, governments and funding agencies who wish to learn new concepts and technologies.

The tutorials are part of the conference technical program, and are free of charge for the conference attendees.

Organizer



Prof. Ishwar Aggarwal, UNC Charlotte, USA

Tutorials & Instructors



Prof. Federico Capasso

Harvard University, USA

Tuesday 20th July

18:00 - 19:00 — Tutorial Room

Tutorial I : Metasurface Flat Optics : from components to mass manufacturing to systems

Flat optics based on metasurfaces has emerged in recent years as a promising alternative to refractive and Fresnel optics in many applications, due to the smaller footprint, mass-manufacturing using the same technology of semiconductor chips, easier control of aberrations and multifunctionality. I will cover recent advances in components and show how they have led to breakthroughs in cameras and other systems such as ultra compact spectrometers.



Prof. Harry Atwater

California Institute of Technology, USA

Wednesday 21st July

18:00 - 19:00 — Tutorial Room

Tutorial II : Design of Active and Reconfigurable Metasurfaces

A grand challenge for nanophotonics is the realization of comprehensively tunable metasurface nanoantenna arrays enabling dynamic, active control of the key constitutive properties of light – amplitude, phase, wavevector and polarization. Achieving this will open new photonics applications in phased-array optical beam steering, visible light modulation for communications and thermal radiation management. This tutorial will discuss design approaches for active and reconfigurable metasurfaces including selection of active materials, electromagnetic design and time-modulation. We will also survey status and outlook for electronically tunable and reconfigurable plasmonic and all-dielectric metasurfaces, whose elements are arbitrarily reprogrammable, enabling a wide array of functions, including steering, focusing, and frequency multiplexing of scattered radiation.



Prof. Ali Adibi

Georgia Institute of Technology, USA

Thursday 22nd July

18:00 - 19:00 — Tutorial Room

Tutorial III : Artificial Intelligence in Meta-optics

A survey of new artificial-intelligence-based approaches for analysis, design, optimization, and knowledge discovery in electromagnetic metastructures will be presented. Recent advances in using both deep learning and machine learning techniques, and their application to practical problems will be covered. These techniques will not only enable more efficient designs of the electromagnetic metastructures (e.g., photonic metasurfaces) but also provide valuable insight about the complex physics of light-matter interactions in such structures. Details of the training process for these algorithms as well as the challenges and limitations of these techniques for different classes of metastructures will be discussed. Knowledge discovery using these techniques includes the study of feasibility of a certain optical response from a given class of metastructures and comparing the roles of different design parameters to facilitate the inverse design process.

GUIDELINES FOR PRESENTERS

META 2021 Online - Don't miss a moment !

META 2021 will be held exclusively online and include both oral and poster presentations.

Oral presentations

Oral presenters at META 2021 will be required to submit a pre-recorded presentation (regular : 10-minutes duration, invited : 15-minutes, keynote : 25-minutes and plenary : 30-minutes) which will be made available to view by the participants up to four weeks after the conference.

All speakers have the possibility to give their talk **live** in the allotted time specified in the technical program. At the time of the presentation, the session chairperson will allow the speaker to share his/her computer screen with the audience (the pre-recording will be used as a backup in case of a no-show or technical difficulty). Speakers also have the option to have their **pre-recorded** presentation played during the live session. **However, all authors, no matter what format they select (live or on replay) should submit a pre-recorded presentation by the deadline of June 30th.**

For both live or on-demand presentations, a 5-minute live Question and Answer (Q&A) session will follow each presentation. The session chairperson will read questions submitted by participants. At least one of the paper's authors must be online to answer questions after the talk.

More information on oral presentations can be found below.

Poster presentations

Poster presenters will be required to submit a poster in **digital format**. They will also have the opportunity to provide a **pre-recorded presentation** (5-minute duration). Both poster and video presentations will be made available for viewing by participants up to three weeks after the conference. Questions can be posted at any time via special chat channels. During the conference, there will be poster sessions scheduled over different time zones where poster presenters will discuss their posters with the attendees via video conferencing. Additional information on poster presentations can be found below.

If a paper is not presented live in the allotted time or a pre-recorded presentation is not submitted by the deadline, then that paper will be considered a "no-show" and will not be published in the Conference Proceedings. If you are unable to make this deadline, please inform contact@metaconferences.org as soon as possible with the date by which your presentation will be available.

Online Platform

The online platform Whova will allow you to join the conference from the comfort of your own home or office and at the convenience of your own schedule all streamlined through one seamless platform. In addition to all poster and oral presentations from the **162 technical sessions.**, the single interface will provide around-the-clock access to :

- 7 plenary lectures
- 21 keynote lectures
- 3 conference tutorials
- Breakout rooms for networking or further discussions 1-on-1 or in a group
- Social and networking activities

All live sessions and lectures will be recorded and available for our participants for viewing after the conference so that they don't miss a thing !

ORAL PRESENTATIONS

Please refer to our website for further details :

<https://metaconferences.org/ocs/index.php/META21/index/pages/view/oral-presentations>

POSTER PRESENTATIONS

Please refer to our website for further details :

<https://metaconferences.org/ocs/index.php/META21/index/pages/view/poster-presentations>

TECHNICAL PROGRAM

All sessions in the program are scheduled in Central European Time (CET/GMT+2/UTC+2).

The program has been put together considering the authors respective time zones.

Tuesday 20th July, 2021

08:45 - 08:45 — Victor Veselago Room

Session 1A1

Opening Address

09:00 - 09:35 — Victor Veselago Room

Session 1A2

Plenary Session I

09:00 : **Plenary talk**

Integrated Nanophotonics for Optoelectronic Computation

Masaya Notomi

Tokyo Institute of Technology (Japan)

In this talk, we show that integrated nanophotonics technologies enable energy-effective O/E and D/A conversions in an integrable fashion. In addition, it is essentially important to add integrable energy-efficient nonlinearities at appropriate places in linear-optic circuits. We present our recent results of such nonlinear elements based on OEO transistors and nanomaterial-loaded nanophotonics. Finally, we will discuss possible applications of these technologies for optoelectronics computations.

Coffee Break and Exhibit Inspection

Session 1P1

Poster session I

09:35 - 10:20

P1: Anapole Excitations in Oxygen-Vacancy-Rich TiO_{2-x} Nanoresonators: Tuning the Absorption for Photocatalysis in the Visible Spectrum

Ludwig Huttenhofer¹, Felix Eckmann², Alberto Lauri³, Javier Cambiasso³, Evangelina Pensa³, Yi Li⁴, Emiliano Cortés¹, Ian D. Sharp², Stefan A. Maier¹

¹Ludwig-Maximilians-Universität München (Germany), ²Technische Universität München (Germany), ³Imperial College London (United Kingdom), ⁴Southern University of Science and Technology (China)

We realize engineered nanoantennas composed of photocatalytic dielectrics and demonstrate increased light harvesting capabilities in otherwise weakly absorptive spectral regions. In particular, we employ anapole excitations, which are known for their strong light confinement, in nanodisks of oxygen-vacancy-rich TiO_{2-x}, a prominent photocatalyst that provides a powerful platform for exploring concepts in absorption enhancement in tunable nanostructures. This concept is general and can be extended to other catalytic materials.

P2: Photonic dualism of icosahedral quasicrystals

Artem Sinelnik, Ivan Shishkin, Xiaochang Yu, Kirill Samusev, Pavel Belov, Mikhail Limonov, Pavel Ginzburg, Mikhail Rybin

ITMO University (Russia)

Here we report on the first fabrication of 3D submicron-size dielectric icosahedral quasicrystals and show a photonic dualism of these structures. Relying on far-field measurements, we found pronounced patterns of unconventional Bragg diffraction, which indicates the existence of multiple photonic pseudogaps

P3: Optical properties of a spatiotemporally modulated surface**Daigo Oue, Kun Ding, John Pendry***Imperial College London (United Kingdom)*

In this research, we develop a differential formalism [J. Chandezon et al., JOSA, 72, 839 (1982).] in order to analytically calculate the optical response of spatiotemporal metasurfaces.

P4: Electron energy-loss spectroscopy of hybrid silicon-on-gold nanoresonators**Artyom Assadillayev¹, Tatsuki Hinamoto², Minoru Fujii², Radu Malureanu¹, Tim Booth¹, Mark L. Brongersma³, Soren Raza¹**¹ *Technical University of Denmark (Denmark)*, ² *Kobe University (Japan)*, ³ *Stanford University (USA)*

Using electron energy-loss spectroscopy (EELS), we show that high-refractive-index silicon nanoparticles on top of a thin gold layer give rise to low-loss hybrid resonances and strong plasmon launching. We characterize the surface plasmon (SP) modes which are launched by the spherical silicon nanoparticle in the presence of the thin gold layer and the Mie modes of the nanoparticle.

P5: Generating vivid colours from black disordered plasmonic network**Changxu Liu¹, Peng Mao², Shuang Zhang², Stefan A. Maier¹**¹ *Ludwig Maximilian University of Munich (Germany)*, ² *University of Birmingham (United Kingdom)*

We design an optical platform that reconciles the two seemingly contradictory criteria, generating vivid structural colours from a disordered network with original black colour. By manipulating the coupling between the disordered system and its optical environment, the random plasmonic network switches from a broadband absorber to a system with tunable resonance. Not limited to the structural colour formation, the platform can be utilised for ultra-sensitive stimuli detection such as nanometer scale height fluctuation and minute refractive index variation.

P6: Self-detunable MRI Signal Enhancement Metasurface On-Bench Characterization**Endri Stoja¹, Diego Betancourt¹, Robin Niklas Wilke², Dennis Philipp², Simon Konstandin², Reiner Umathum², Jurgen Jenne², Thomas Bertuch¹, Matthias Gunther³**¹ *Fraunhofer FHR (Germany)*, ² *Fraunhofer MEVIS (Germany)*, ³ *University of Bremen (Germany)*

A simple technique for the detuning during transmission (Tx) of an MRI signal enhancement metasurface is proposed. The metasurface itself is composed of a linear alignment of wire resonators, whereas the detuning is achieved via inductive coupling to a varactor-loaded loop which senses the RF field strength. On-bench measurements validate the detuning efficiency. Detuning during the Tx phase avoids excitation field distortion, targeted flip angle modification and limits patient exposure.

P7: Optical Extinction Cross-Section Enhancement of a Dielectric Cylinder through Meta-surface made of Graphene-Coated Spherical Nanoparticles**Shiva Hayati Raad, Zahra Atlasbaf***Tarbiat Modares University (Iran)*

Graphene-coated spherical nano-particles are arranged around a dielectric cylinder in order to enhance its extinction cross-section. Initially, a 1D array of the particles is considered in different loci with respect to the incident wave. It is observed that regardless of the position of the particles, the extinction cross-section is considerably enhanced with respect to the bare cylinder due to the excitation of localized surface plasmons in shells. By increasing the number of particles, extinction cross-section is further enhanced.

P8: Absorption control in epsilon-near-zero hyperbolic metamaterials based on InAs**Michal Dudek, Alessandro Pianelli, Rafal Kowrdziej, Janusz Parka***Military University of Technology (Poland)*

Here we present the possibility to control the epsilon-near-zero (ENZ) region by properly designing a hyperbolic metamaterial (HMM) based on InAs, which allows for an adjustable absorption in mid-IR. Numerical results show that by increasing thickness of the undoped InAs layers we are able to redshift the absorption by 14 microns. Also, transition from hyperbolic dispersion type I to ENZ and type II is observed. The designed HMM stack may provide a platform for perfect metamaterial absorbers in mid-IR.

P9: Photon Nanosources Using Low Energy Tunnel Electrons**Gérald Dujardin, Elizabeth Boer-Duchemin, Rémi Bretel, Shuiyan Cao, Eric Le Moal, Delphine Pommier***Université Paris-Sud (France)*

We explore new types of electrically-driven photon nanosources based on the use of low energy tunnel electron excitations with the Scanning Tunneling Microscope (STM).

P10: Differential field Enhancement using Non-Reciprocal Material for Plasmon-Enhanced Upconversion at a Particular Wavelength

Jasvith Raj Basani, Nikhil Navaratna, Waseem A. Wani, Kannan Ramaswamy

Birla Institute of Technology and Science Pilani (India)

Upconversion, the process of changing the energy of a photon from a lower energy to a higher energy, is an area with a vast multitude of applications - from efficient hybrid solar cells to background-free bioimaging. Presently, however, upconverting techniques used are inefficient and are economically unfeasible. To maximize the efficiency of upconverting technologies, the system could ideally react to a band of frequencies - particularly, to have different enhancements for different frequencies, by introducing a nonreciprocal spacer.

P11: Ultra-wide Bandwidth Electromagnetic Wave Absorbers with Multilayered Frequency Selective Surfaces

Tian Liu, Sung-Soo Kim

Chungbuk National University (Korea)

A high-capacitive frequency selective surface (FSS) with a new structure of folded spiral conductors has been proposed as the small-array periodicity and low-frequency resonance FSS for the aim of ultra-wide bandwidth absorbers in multilayer structure. Through a combination of the high capacitive spiral FSS with other conventional FSS (square loop, square patch) with a medium- and high-frequency resonance, an ultra-wide absorption bandwidth (4.7-50.0 GHz for -10 dB reflection loss) can be designed with a small total thickness close to the theoretical limit.

P12: Engineering spontaneous valley coherence in two dimensional heterostructures

Anshuman Kumar

Indian Institute of Technology Bombay (India)

In order to harness the valley degree of freedom in TMDCs, it is critical to be able to actively control the coherence between excitons in the two valleys. Here we demonstrate how spontaneous valley coherence (SVC) can be achieved by creating a heterostructure of the TMDC with other 2D materials. Subsequently we will present our recent results to show tunability of SVC using certain commonly available 2D semiconductor heterostructures.

P13: High-efficient phase modulator based on all dielectric metasurface electrically tunable by thin layer liquid crystal for visible light modulation

Mingyu Sun

University of Shanghai for Science and Technology (China)

We propose phase modulator by LC composited TiO₂ metasurface. Under effect of Huygens' condition, obvious binary phase shift with high transmittance about 80% is obtained with small driving voltage of 4~8 V, and a large dipole resonance extension above 150 nm to be expected. The convolutional neural network based inverse design setup helps to anticipate the proper LC layout with respect to the measured optical output. Such LC composited metasurface device could be a solution to the high-efficient phase modulator.

P14: Sorting full angular momentum states with all-dielectric Pancharatnam-Berry spiral transformation metasurfaces

Baiming Wang¹, Yuanhui Wen¹, Jiangbo Zhu², Yujie Chen¹, Siyuan Yu³

¹Sun Yat-sen University (China), ²Northumbria University (United Kingdom), ³University of Bristol (United Kingdom)

Spin and orbital angular momenta (SAM and OAM), constitute a complete and higher state space of light. Here we propose a scheme that combines the spiral coordinate transformation with all-dielectric Pancharatnam-Berry metasurfaces for high-resolution simultaneous sorting SAM and OAM states of light. Simulations and experiments are demonstrated sorting 7 OAM states and 2 SAM states. This compact, high-resolution, and high-efficiency full angular momentum state sorter could have applications in both classical and quantum information systems.

P15: Influence of ordering of gold nanohole arrays on plasmonic resonances

Brindhu Malani S.¹, P. Viswanath²

¹Mangalore University (India), ²Centre for Nano and Soft Matter Sciences (India)

Gold nanoholes of varying hole sizes and ordering were fabricated using colloidal lithography. The degree of ordering was quantified using pair-correlation function and bond-orientational order parameters. Reflectance studies from experiments and simulations of these samples shows that it supports both localized surface plasmon resonance and propagating surface plasmon polaritons. We found that spectral resolution improves for sample with long range order and higher bond-orientational order parameter. Based on plasmonic responses and order quantification optimum sample is chosen for refractive index sensing.

P16: Realization of photonic time crystal in a trapped exciton-polariton condensate

Ting-Wei Chen¹, Shih-Da Jheng², Szu-Cheng Cheng²

¹National Chiayi University (Taiwan), ²Chinese Culture University (Taiwan)

A photonic time crystal in a trapped polariton condensate is theoretically realized. Numerical modeling based on the open-dissipative Gross-Pitaevskii equation suggests the formation of a quantized vortex surrounded by azimuthal superflows within the annular excitation area.

P17: Optomagnetic field in nonmagnetic plasmonic nanostructures

Vage Karakhanyan, Clement Eustache, Yannick Lefier, Thierry Grosjean

University of Bourgogne - Franche-Comté (France)

Using simplified hydrodynamic model, we theoretically investigate resonant inverse Faraday effect within individual plasmonic nanostructures. Upon illumination with circularly polarized light, resonant nanostructures are shown to develop an optomagnetic field that is controllable by the helicity of the light. Given their sub-micron footprint, individual plasmonic nanostructures open new prospects towards ultrafast and polarization-controlled tunable magnetism on the nanoscale, thus potentially impacting large panel of application and techniques including all optical magnetization switching, spin-wave excitation and optomagnetic tweezing of nano-objects.

10:20 - 12:40 — Victor Veselago Room

Session 1A3

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

10:20 : Invited talk

Nanophotonic systems for image processing

Lukas Wesemann, Timothy Davis, Ann Roberts

The University of Melbourne (Australia)

Conventional all-optical image processing requires relatively bulky optics limiting its potential use in mobile electronic systems. Here, the use of nanoscale resonant gratings and other thin film structures for image processing of both amplitude and phase objects is presented.

10:40 : Invited talk

Spatial and dynamical multi-level control over thermal emission

Ziquan Xu¹, Huanzheng Zhu¹, Min Qiu², Qiang Li¹

¹Zhejiang University (China), ²Westlake University (China)

We demonstrate a thermal emitter with spatial and dynamic tunability by using phase-transition material vanadium dioxide. Emissivity varies from 0.19 to 0.91 during the insulator-to-metal transition. Laser-induced phase transition and its application in multi-level spatial control of thermal emission are also demonstrated.

11:00 : Invited talk

High refractive index contrast meta-surfaces for sensing and emitting devices

Yuusuke Takashima, Masanobu Haraguchi, Yoshiki Naoi

Tokushima University (Japan)

High contrast meta-surfaces support several optical eigenmodes, and the interference of the excited modes provides the extraordinary optical properties. We experimentally have demonstrated unique ultraviolet and blue-violet devices with high contrast meta-surface, such as a highly polarized ultraviolet light-emitting diode with very low loss and highly sensitive refractive index sensor with very simple optical system operating blue-violet wavelength.

11:20 : Invited talk

Synchronously wired infrared antennas for resonant single-quantum-well photodetection up to room temperature

Hideki T. Miyazaki¹, Takaaki Mano¹, Takeshi Kasaya¹, Hiroataka Osato¹, Kazuhiro Watanabe¹, Yoshimasa Sugimoto¹, Takuya Kawazu¹, Yukinaga Arai¹, Akitsu Shigetou¹, Tetsuyuki Ochiai¹, Mel F. Hainey Jr.¹, Yoji Jimba², Hiroshi Miyazaki³

¹National Institute for Materials Science (Japan), ²Nihon University (Japan), ³Tohoku University (Japan)

To apply optical patch antennas sandwiching dielectrics between metal layers for optoelectronic devices, wiring to each electrically isolated antenna is indispensable. We show that geometrically engineered metallic wires interconnecting the antennas can function to synchronize the optical phases for promoting coherent resonance, not only as electrical conductors. Antennas connected with optimally folded wires are applied to quantum well infrared photodetectors, and a polarization-independent quantum efficiency as high as 61 % at 78 K, even extending to room temperature, is demonstrated.

11:40 : Invited talk

Bloch surface wave platform in the near- and mid-infrared regions

Chih-Zong Deng¹, Ya-Lun Ho¹, Yang-Chun Lee², Zhiyu Wang¹, Yi-Hsin Tai¹, Hirofumi Daiguji¹, Jean-Jacques Delaunay¹

¹The University of Tokyo (Japan), ²National Taiwan University (Taiwan)

Bloch surface wave (BSW) platform opens up new possibilities in the design of sensors and components for photonic circuits owing to its superior properties such as low propagation losses that enable long optical communication length and large surface sensing area. Here, we report a BSW platform realizing well-coupled BSW in a wide spectral range from the NIR to the MIR with possible applications in light guiding and sensing.

12:00 : Invited talk

The Wonderful World of Flat Bands

Sergej Flach

Institute for Basic Science (Korea)

Certain lattice wave systems in translationally invariant settings have spectral bands that are strictly flat or independent of momentum, arising from either internal symmetries or fine-tuned coupling. These flat bands display remarkable strongly interacting phases of matter. Flat bands have now been observed in a variety of settings, ranging from electronic systems to ultracold atomic gases and photonic devices. I will review the design and implementation of flat bands and chart future directions of this exciting field.

12:20 : Invited talk

Acoustic metamaterial beam with a perfect bandgap based on a quadruple-mode resonator array

Motonobu Tomoda, Kentaro Fujita, Osamu Matsuda, Oliver B. Wright

Hokkaido University (Japan)

We introduce a metamaterial in the form of a rectangular cross-section beam made of a single isotropic material that can simultaneously suppress all elastic-wave polarizations, i.e. compressional, in-plane shear, flexural and torsional waves, over a range of frequencies around 1 kHz. This is experimentally achieved by machining replicas of a unit cell based on a planar resonator with interconnected ribs in an aluminum beam. Our method opens the way to various applications in easily-manufacturable vibration isolators and novel acoustic wave control.

10:20 - 12:30 — Allan Boardman Room

Session 1A4

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

10:20 : Keynote talk**Semiconductor Nanocrystal Optoelectronics: Pushing the Limits****Volkan Demir Hilmi***Bilkent University (Turkey)*

In this talk, we will introduce the emerging field of semiconductor nanocrystal optoelectronics, with most recent examples of their photonic structures and optoelectronic devices employing such atomically flat, tightly confined, quasi 2 dimensional CQWs, also popularly nick named "nanoplatelets".

10:50 : Invited talk**Metamaterial Absorber-enhanced Light-harvesting and Light-energy conversion****Peng Yu¹, Wenhao Wang¹, Zhiming Wang¹, Alexander Govorov²***¹University of Electronic Science and Technology (China), ²Ohio University (USA)*

In this talk, we will present an overview of metamaterial absorber-enhanced light-harvestings and their use in the light-energy conversion devices, such as hot electron generation for photochemistry and photothermal for bolometry.

11:10 : Invited talk**Plasmonic nanostructures for solar fuels generation****Alberto Naldoni***Palacky University (Czech Republic)*

The development of large-scale solar energy conversion technologies for fuels and chemicals is currently hindered by low yields and scarce product selectivity. Plasmonic nanostructures have emerged as a promising building block to enhance light-matter interaction in solar fuel devices and to trigger alternative reaction pathways. In this talk, I will summarize our recent results about the integration of plasmonic nanoantennas and nanoparticles in systems for solar fuels generation, discussing the fundamental mechanisms underlying the observed enhanced reactivity.

11:30 : Invited talk**Optically Active Nanostructures****Yurii K Gun'ko, Fearghal C. Donnelly, Finn Purcell-Milton, Vera A. Kuznetsova***Trinity College Dublin (Ireland)*

We present advances in the development of various optically active nanomaterials. We discuss the main approaches to induce optical activity in nanomaterials, including: the synthesis of nanomaterials in the presence of chiral ligands, post-synthetic capping of achiral nanostructures with chiral molecules, presence of chiral defects and even simple stirring of achiral anisotropic colloidal nanostructures. Most of the optically active nanostructures demonstrate strong unique circular and linear dichroism responses and very interesting structural morphologies.

11:50 : Invited talk**Measuring the magnetic dipole transition of single nanorods by Fourier microscopy****Reinaldo Chacon¹, Aymeric Leray¹, Jeongmo Kim², Khalid Lahli², Sanro Mathew¹, Alexandre Bouhelier¹, Jong-Wook Kim², Thierry Gacoin², Gérard Colas des Francs¹***¹Université Bourgogne Franche-Comté (France), ²Ecole Polytechnique (France)*

Rare-earth ions can present optical transitions with significant magnetic dipole (MD) character that can be manipulated by the crystalline or molecular hosting environment. They are of strong interest for engineering light-matter interaction at the nanoscale with numerous applications in nanophotonics. We analyze in details the optical transition in individual and single crystalline rare-earth doped nanocrystals. We measure the MD orientation and demonstrate a 100 % magnetic transition in our homemade synthesized single crystalline

nanorods.

12:10 : Invited talk

Photonic flat-band lattices and unconventional light localization

Zhigang Chen

Nankai University (China)

In this talk, we briefly summarize developments on the flat-band localization and the associated phenomena in photonic lattices, including unconventional line states and noncontractable loop states. We show that such photonic lattices offer a convenient platform for probing the underlying physics of flat-band systems, which may provide inspiration for exploring fundamentals and applications of flat-band physics in other structured media from metamaterials to other nanophotonic materials.

10:20 - 12:30 — Tatsuo Itho Room

Session 1A5

Bottom-up approaches, new fabrication routes and ENSEMBLE3

Organized by: Dorota Pawlak and Virginie Ponsinet

Chaired by: Dorota Pawlak and Virginie Ponsinet

10:20 : Invited talk

Planar nanooptics in anisotropic van der Waals crystals

Pablo Alonso-Gonzalez

University of Oviedo (Spain)

Highly anisotropic crystals have recently attracted considerable attention because of their ability to support polaritons with a variety of unique properties, such as hyperbolic dispersion, negative phase velocity, or extreme confinement. Here, we will show experimental demonstrations of the unique behaviour of PhPs in these crystals including highly directional propagation and ultra-low losses.

10:40 : Invited talk

Nanostructured optical fibers

Rafal Kasztelan¹, Dariusz Pysz¹, Adam Filipkowski¹, Alicja Anuszkiewicz², Marcin Franczyk¹, Ryszard Buczynski¹

¹University of Warsaw (Poland), ²Lukasiewicz Research Network - Institute of Microelectronics and Photonics (Poland)

The nanostructuring technique allows to fabrication of all-glass optical fibers with properties not differing from the existing solutions. Moreover, nanostructuring makes it possible to break the circular symmetry of the fiber structure and fabricate new fibers, impossible to manufacture with other methods, as well as to optimize the fiber optical parameters more extensively.

11:00 : Keynote talk

Publishing in Nature Journals

Rachel Won

Nature Photonics (United Kingdom)

This talk, although with an emphasis on Nature Photonics, will introduce you to all the Nature journals, and cover the detailed information and guidelines on scientific manuscript preparation and submission. Also presented is an overview on the editorial and peer-review processes in all Nature journals. For those who are interested in being an editor, you will get to know the selection criteria and process of Nature journals.

11:30 : Invited talk

Self-assembled chiral plasmonic metasurfaces

Matthias Pauly, Vincent Lemaire, Sriharani Sekar, Wenbing Wu, Gero Decher

Université de Strasbourg (France)

Chiral plasmonic metasurfaces are self-assembled from oriented non-chiral metallic nanowires and nanorods using Grazing Incidence Spraying combined to the Layer-by-Layer assembly approach. The resulting thin films display a very high circular dichroism over the whole visible and near-infrared wavelength range, and the optical properties are highly dependent on the superstructure which can be easily tuned by this approach.

11:50 : Invited talk

Optical Properties of Artificial Chiral Metasurfaces

Emilija Petronijevic¹, Grigore Leahu¹, Alessandro Belardini¹, Roberto Roberto Li Voti¹, Concita Sibilia¹, Tiziana Cesca², Giovanni Mattei²

¹Universita di Roma La Sapienza (Italy), ²University of Padova (Italy)

Here we investigate optical chiral properties of metasurfaces fabricated by means of nanosphere lithography. This low-cost and simple technique combined with angled evaporation of a plasmonic layer was employed to produce arrays of tilted elliptical nanoholes in Au or Ag. The elliptical shape and the in-plane tilt lead to the symmetry breaking, which further enables different coupling with circularly polarized light of opposite handedness. We investigate intrinsic and extrinsic chiral behavior in the visible and near infrared range.

12:10 : Invited talk

Scattering electromagnetic eigenstates of a two-constituent composite and their exploitation for calculating a physical field

David J Bergman, Parry Chen, Asaf Farhi

Tel Aviv University (Israel)

The spectral representation of an electric field in a two-constituent composite medium is revisited. A theory is developed for calculating the electromagnetic (EM) eigenstates of Maxwell's equations for such a composite where the magnetic permeability as well as the electric permittivity have different uniform values in the two constituents.

10:20 - 12:20 — Christian Huygens Room

Session 1A6

Symposium III: Advanced passive and active metasurfaces and zero-index optics

Organized by: Howard Lee, Yu-Jung (Yuri) Lu and Zi Jing Wong

Chaired by: Howard Lee, Yu-Jung (Yuri) Lu and Zi Jing Wong

10:20 : Invited talk

Metasurface enhanced high-sensitive IR spectroscopy

Takuo Tanaka

RIKEN and Tokushima University (Japan)

Metasurfaces consist of metal-insulator-metal structure were developed for a versatile platform of high-sensitive IR spectroscopy. A device with nano-fluidic channel allows the introduction and precise control of number of analyte molecules into the intense electromagnetic field of metamaterials, resulting in the improvement of sensitivity up to 2 orders compared to state-of-the-art plasmonic enhanced IR spectroscopies. High sensitive gas spectroscopy has also demonstrated by the use of vertically aligned MIM metasurface device.

10:40 : Invited talk

Generation of high-efficiency light sheets by dielectric metasurfaces

Dangyuan Lei

City University of Hong Kong (Hong Kong)

In this talk, I will present a few analytical and intelligent approaches for designing high-efficiency light sheets with dielectric metasurfaces, including realization of multiple light sheets by metasurface-based Fourier transformation, structured light sheets by aperture-like metasurface arrays, large-area light sheet by phase-controlled intelligent metasurface. I will also discuss the possibility of using phase-changing materials to achieve switchable metasurfaces.

11:00 : Invited talk**Metasurface design and application for compact AR device****ByoungHo Lee, Jangwoon Sung, Gun-Yeal Lee***Seoul National University (Korea)*

The principles and applications of metasurface will be given with detailed examples. Three metasurface designs that can exceed the conventional optical modulation will be given, which are complex-amplitude modulation, full-space visible light modulation, and phase-change material-based metasurface. Then, metalens eyepiece for augmented reality will be discussed.

11:20 : Invited talk**Active Metasurfaces based on Phase-Change Materials****Min Qiu***Westlake University (China)*

Metasurfaces, incorporated with phase-change materials, show extraordinary tunability and reconfigurability with a simple design. We explore their possible applications in dynamic light manipulation, emissivity control and thermal management and camouflage, etc., which hold high promise for uncovering novel physical phenomena and settling practical problems.

11:40 : Invited talk**Near-infrared Active Metasurfaces for Tunable Beam Diffraction and Dynamic Polarization Conversion****Pin Chieh Wu¹, Ruzan Sokhoyan², Ghazaleh Kafaie Shirmanesh², Wen-Hui Cheng², Ragip A. Pala², Meir Grajower², Harry Atwater²**¹*National Cheng Kung University (Taiwan)*, ²*California Institute of Technology (USA)*

In this talk, I will firstly discuss an all-dielectric active metasurface platform based on electro-optically tunable III-V multiple-quantum-well (MQW) resonators. By selectively applying an electrical bias to metasurface elements, we experimentally realize a dynamically switchable diffraction grating and beam steering. Subsequently, I will report a scheme of active polarization modulation by using indium tin oxide (ITO)-based tunable metasurfaces. By suitably biasing the metasurface structure, the polarization of reflected light can be converted to versatile states.

12:00 : Invited talk**Generating High Resolution Accelerating Optical Beams and Beam Array Based on All-dielectric metasurfaces****Binbin Yu, Lei Chen, Xu Chen, Dawei Zhang, Jing Wen***University of Shanghai for Science and Technology (China)*

We experimentally realize high-resolution Airy optical beams and beam array with long working and non-diffraction propagation distances based on our proposed highly compact all-dielectric synthetic metasurface, that integrate a cubic phase and the phase of a Fresnel holographic lens. In addition, we demonstrate that imposing the phase of a Dammann grating to the above synthetic metasurface produces an array of Airy optical beams.

10:20 - 12:30 — Augustin Fresnel Room**Session 1A7****Symposium IV: Chirality, magnetism, and magnetoelectricity: Separate phenomena and joint effects in metamaterial structures**

Organized by: Eugene Kamenetskii

Chaired by: Eugene Kamenetskii

10:20 : Invited talk**Interaction of chiral light with topological singularities in plasmonic metasurfaces****Eliav D. Epstein, Leeju Singh, Maayan Fox, Shmuel Sternklar, Yuri Gorodetski**

Ariel University (Israel)

We present simple metasurfaces with different topologies supporting plasmonic edge states at the boundary between them. Preliminary experiments and numerical simulations show the achieved mode localization in line defects and point singularities. Line singularities are shown to support dark and bright modes. Point singularities show strong localization of light which can be further modified by the varying topological order of the structure.

10:40 : Invited talk

Theory of Photoinduced Spin Polarization in Spin-Orbit-Coupled Systems

Masahito Mochizuki, Takashi Inoue, Yasuhiro Tanaka

Waseda University (Japan)

We theoretically study the photo-induction of spin polarization in electron systems with spin-orbit interactions irradiated with circularly polarized light. It is demonstrated that the rotating electric field of CPL is converted to an effective static field and induces the spin polarization perpendicular to the polarization plane. We find that the magnitude and sign of the induced spin polarization are governed by the electron filling and the Fermi-surface geometry.

11:00 : Invited talk

Time-resolving surface plasmon orbital angular momentum from pure geometric chirality

Chen-Bin Huang¹, Yanan Dai², Hrvoje Petek²

¹*National Tsing Hua University (Taiwan)*, ²*University of Pittsburgh (USA)*

We demonstrate the creation of plasmonic OAM contributed purely through the geometrical chirality of the spiral. The experiments are carried out through time-resolved two-photon photoemission electron microscopy, where the recorded movies reveal the orbiting trajectories of surface plasmon polaritons coupled by the spiral under linearly-polarized excitations.

11:20 : Invited talk

Theory of Single Molecular Near field Circular Dichroism by Photo-induced Force Microscopy

Hajime Ishihara, Hidemasa Yamane, Nobuhiko Yokoshi

Osaka Prefecture University (Japan)

We theoretically propose the near field circular dichroism (CD) of single molecules by photo-induced force microscopy (PiFM). Based on the extended discrete dipole approximation (e-DDA) incorporating the nonlocal optical response, we calculate PiFM CD map. The result indicates a greatly enhanced near field CD of single molecules.

11:40 : Invited talk

Universal energy barriers of large magnetic skyrmions

Jan Masell

RIKEN Center for Emergent Matter Science (Japan)

Magnetic skyrmions are whirls in the magnetization for which a topological winding number can be defined. Because of this topology, skyrmions are often believed to gain some superior topological protection, in particular close to the continuum limit. However, even in this limit their energy barriers remain finite for a large class of systems. Moreover, the height of the barrier is determined by higher order gradients of the continuum theory which arise, e.g., from higher order interactions.

12:00 : Keynote talk

A new spin for acoustics

Konstantin Bliokh

RIKEN (Japan)

I will show theoretical and experimental results proving that water waves possess nontrivial spin and momentum densities, which can be described within a universal relativistic field theory construction. These results provide a new twist for sound and water-surface waves, and offer new fields for applications of various vector-wave phenomena, so far mostly restricted to photonic and quantum-electron systems.

10:20 - 12:30 — Ibn Al-Haytham Room

Session 1A8

Symposium V: Phononics and acoustic metamaterials

Organized by: Jensen Li and Guoliang Huang

Chaired by: Jensen Li and Guoliang Huang

10:20 : **Invited talk****Asymmetric Elastic-wave Transmission in a Lossless Metasurface****Bing Li¹, Yabin Hu¹, Yongquan Liu², Meiyang Zhao¹, Zheng Li³**¹Northwestern Polytechnical University (China), ²Xi'an Jiaotong University (China), ³Peking University (China)

Our research presents theoretical, numerical and experimental investigations of asymmetric elastic-wave transmission in a thin plate with a lossless elastic metasurface. A theoretical framework is developed to control the asymmetric elastic-wave transmission without tailoring losses. We numerically and experimentally demonstrate that the asymmetric behavior can be realized based on higher order diffraction in a lossless structural waveguide. We further experimentally show that the novel phenomenon can be obtained within a relatively broad range of incident angles and frequency band.

10:40 : **Invited talk****Giant extraordinary transmission of longitudinal acoustic waves****Oliver B. Wright¹, Thibaut Devaux¹, Hiroya Tozawa¹, Paul H. Otsuka¹, Sylvain Mezil¹, Motonobu Tomoda¹, Osamu Matsuda¹, Eun Bok², Sam Hyeon Lee²**¹Hokkaido University (Japan), ²Yonsei University (Korea)

We demonstrate extraordinary longitudinal bulk-wave acoustic transmission inside a solid structure, acting as an acoustic meta-atom, that is connected between two tungsten blocks. By including concentric grooves of optimal dimensions, we show how the extraordinary acoustic transmission efficiency can be increased to a value exceeding 500, and how directed output beams inside solids can be produced. Applications include acoustic imaging and sensing.

11:00 : **Invited talk****Self-synchronization of Thermal phonons in a Charged Silicon Resonator System****Zhongwei Zhang¹, Yangyu Guo¹, Marc Bescond¹, Jie Chen², Sebastian Volz¹, Masahiro Nomura¹**¹The University of Tokyo (Japan), ²Tongji University (China)

Self-synchronization is an important collective behavior, in which oscillators are dynamic in a coherent state. In this work, we demonstrate that the Coulombic force coupled thermal phonons can be spontaneously synchronized, without any external driving, which is well understood from the synchronization of frequency and phase. By generating coherent thermal phonons, self-synchronization also significantly enhances thermal transport. Our findings might promote physical understanding of the emergence of coherent phonons, and also the engineering of nanoscale thermal transport.

11:20 : **Spatiotemporal effective media for acoustic waves****Xinhua Wen¹, Xinghong Zhu¹, Hong Wei Wu², Jensen Li¹**¹Hong Kong University of Science and Technology (Hong Kong), ²Anhui University of Science and Technology (China)

Based on an approach of virtualized metamaterials with software-defined impulse response, we experimentally realize an acoustic spatiotemporal effective medium in dynamically switching between two resonating configurations with a modulation frequency at least 5 times higher than the signal frequency. We also establish the effective medium formula.

11:35 : **Invited talk****Experimental investigation of defect modes in tubular phononic crystals****Frieder Lucklum**

Technical University of Denmark (Denmark)

In this contribution the use of defect modes in phononic crystals for sensing applications will be discussed on the basis of different practical realizations. As a specific example, experimental results of wave transmission through tubular phononic crystals will be compared for different materials and in the presence of different geometric defects to the ideal crystal as well as to reference measurements through a corresponding hollow tube and solid cylinder.

11:55 : Invited talk

On the use of a Helmholtz resonator to acoustically dope a Plate-type metamaterial

Matthieu Mallejac¹, Aurélien Merkel², Vincent Tournat¹, Vicent Romero-Garcia¹, Jean-Philippe Groby¹

¹Le Mans Université (France), ²Université de Lorraine (France)

In this work, we analytically, numerically, and experimentally investigate the feasibility of an acoustic analogue of the photonic doping effect, i.e., the ability of changing the effective properties of a medium by embedding a dopant. We show that a one dimensional Plate-type Acoustic Metamaterial (PAM) can be efficiently doped using a single doping impurity, e.g., a tuned Helmholtz resonator. The influence of both the location of the dopant and the losses are studied.

12:15 : Reminiscence of edge states in 1-dimensional hyperuniform acoustic materials

Svetlana Kuznetsova¹, Jean-Philippe Groby¹, Lluís M. Garcia-Raffi², Vicent Romero-Garcia¹

¹Université du Mans (France), ²Universitat Politècnica de Valencia (Spain)

Hyperuniform materials are systems made from point distributions that suppress density fluctuations on large length scales. They present wide and isotropic bandgaps despite being highly disordered and degenerate. Here, we report the localization of acoustic waves at the interface between two 1-dimensional hyperuniform materials yielding in different states. Both materials represent an air-filled acoustic waveguide with rigid diaphragms acting as scatterers. Tunability of the band structure and emergence of the edge modes provide promising applications in wave control devices.

10:20 - 12:20 — Gaston Floquet Room

Session 1A9

Acoustic and seismic metamaterials

10:20 : A homogenization model for acoustic metasurfaces and design sensitivity analysis for topology optimization

Yuki Noguchi¹, Takayuki Yamada²

¹Tokyo Institute of Technology (Japan), ²Kyoto University (Japan)

In this paper, an efficient numerical analysis model for acoustic metasurfaces is provided based on a homogenization method. Considering its application to the optimal design of acoustic metasurfaces by topology optimization, we discuss the design sensitivity called the topological derivative. Numerical examples are provided to confirm the validity of the proposed model and sensitivity.

10:35 : Fluid-like Elastic Reflective Metasurface

JooHwan Oh, YeJeong Shin

Ulsan National Institute of Science and Technology (Korea)

Elastic mode conversion has been considered as a unique characteristic of elastic waves that cannot be avoided. In this paper, a fluid-like elastic reflective metasurface that can break this coupling is proposed with numerical and experimental supports. In other word, only longitudinal wave is reflected for the obliquely incident longitudinal wave, i.e., the surface behaves as fluid boundary. We believe that our research may provide a new way in elastic metasurface technologies.

10:50 : Deep-subwavelength holey acoustic second-order topological insulators for spoof surface wave

Zhiwang Zhang¹, Ying Cheng¹, Dajian Wu², Desheng Ding³, Xiaojun Liu¹

¹Nanjing University (China), ²Nanjing Normal University (China), ³Southeast University (China)

We illustrate a holey HOTI capable to sustain deeply confined corner states fifty times smaller than the

wavelength. Not only do we experimentally observe a remarkable resilience of these surface confined acoustic states against defects, but do also demonstrate topologically protected sound in three different frequency regimes. Concerning this matter, our findings will thus have the capability to push forward exciting applications for robust acoustic imaging way beyond the diffraction limit.

11:05 : Design and Manufacturing of Monolithic Mechanical Metastructure with Ultrawide Bandgap for Low Frequency Vibration and Noise Control

F. Muhammad, C. W. Lim

City University of Hong Kong (Hong Kong)

The present study proposes a novel 3-D monolithic mechanical metastructure with capability to induce ultrawide three-dimensional bandgap with relative bandwidth or gap-to-mid-gap ratio 171.5%. The bandgap is induced and discussed by principle of mode separation that utilizes the locally resonant global and local modes to open ultrawide bandgap. The wave attenuation inside the bandgap frequencies are demonstrated by developing finite array model and performing numerical frequency response study. The numerical findings are corroborated through experiment test on 3-D printed prototype.

11:20 : Sonic semimetals with intersecting nodal rings

Liyang Zheng, Johan Christensen

University Carlos III de Madrid (Spain)

Nodal rings, protected by bulk topology and symmetries, arise from the crossing of the valence and conduction bands. In this work, we experimentally demonstrate three-dimensional sonic semimetals that exhibit intersecting nodal rings, and further investigate their topological origin and the corresponding exciting wave properties. The nontrivial topology characterized by winding number due to nonsymmorphic symmetry guarantees the appearance of zero-energy surface waves. Novel transport phenomena, such as negative reflection and focusing, are also discussed.

11:35 : Coupled Scholte modes supported by soft elastic plates in water

Bethany Staples, Thomas Graham, Alastair Hibbins, Roy Sambles

University of Exeter (United Kingdom)

Unlike many materials, such as aluminium, the transverse wave speed in soft solids are less than that of longitudinal sound in water. This leads to a localized surface wave, the Scholte wave, having a speed well below that of sound in water. Here we use a thin acrylic plate and demonstrate that the two Scholte waves couple across the plate by using pulsed ultrasound excitation to record the dispersion of both the slow antisymmetric and faster symmetric coupled waves.

11:50 : Numerical and conceptual design of vibroacoustic metamaterial solutions for structural vibration reduction in launcher components

Daria Manushyna¹, Heiko Atzrodt¹, Marvin Droste¹, Niels Deschauer²

¹Fraunhofer Institute for Structural Durability and System Reliability LBF (Germany), ²MT Aerospace AG (Germany)

This work deals with numerical and conceptual design of vibroacoustic metamaterial solutions for structural vibration reduction in launcher components. The adjustable and compact design of vibroacoustic metamaterials based on a local resonance effect leads to an attractive and flexible solution especially for lightweight thin-walled structures in space industry. An inverse unit cell modelling approach is used to for the numerical design of a demonstrator including multiple tuned vibration absorbers (TVA).

12:05 : Multiple Scattering Theory in the study of Non-Hermitian Sonic Second Order Topological Insulators

Maria Rosendo Lopez¹, Zhiwang Zhang¹, Daniel Torrent², Johan Chistensen¹

¹Universidad Carlos III de Madrid (Spain), ²Universitat Jaume I (Spain)

Here, we make use of the Multiple Scattering Theory to calculate the topological corner states of both Hermitian and Non Hermitian Sonic Second Order Topological Insulators. Our findings reveal that the sound is trapped in the corners of the Concentric Square Crystal considered, based on an inner SC made up of a topological non-trivial region enclosed by a topological trivial region. This approach allows us to compute the spectral dependence of corner states with defects, showing its robustness.

10:20 - 12:35 — Lawrence Bragg Room**Session 1A10****Plasmonics: Fundamentals and Applications**

Organized by: Hong Wei

Chaired by: Hong Wei

10:20 : Invited talk**From passive to active manipulation of the polarization states of electromagnetic waves by plasmonic metastructures****Ruwen Peng, Xiang Xiong, Ren-Hao Fan, Mu Wang***Nanjing University (China)*

In this talk, we focus on both passive and active manipulation of the polarization states of light with 2D/3D plasmonic metastructures.

10:40 : Invited talk**Colloidal self-assembly route towards efficient designing of nanophotonic architectures****Swagato Sarkar¹, Joby Joseph¹, Tobias A.F. Koenig²**¹*Indian Institute of Technology Delhi (India)*, ²*Leibniz Institute of Polymer Research Dresden e. V. (Germany)*

We report a plasmonic grating of Au nanoparticle chains on a TiO₂ layer to study the dispersion relation through simple spectroscopic methods. Compared to Au nanobars, the experimentally observed range of hybridized guided-modes can now be extended to modes along the nanoparticle chain lines. Fabrication of such hybrid-architectures over macroscopic areas through successful confluence of top-down and bottom up approaches like interference lithography and colloidal self-assembly respectively has resulted in polarization-dependent sensitivity enhancement.

11:00 : Invited talk**Multipole Engineering in Silicon Mie Resonator with Cap Layer****Junichi Takahara, Ikuto Hotta***Osaka University (Japan)*

We study the mechanism of spectral narrowing of resonant scattering from a silicon Mie resonator by adding a cap layer. Multipole decomposition analysis reveals that lossy materials cause spectral red-shift of electric dipole only and induces the narrowing of Mie resonance.

11:20 : Invited talk**Nanoantennas Made of Gold and Silver Nanowires****Jianfang Wang, Xiaolu Zhuo***The Chinese University of Hong Kong (Hong Kong)*

Wet-chemistry methods are developed to grow high-aspect-ratio gold and silver nanorods in high purity. These nanorods possess higher-order plasmon modes. Moreover, their longitudinal dipolar plasmon mode can be synthetically varied from the visible to the mid-infrared region. The broadside-antenna and color-routing functions are demonstrated with these nanorods, where light of different colors is directed to different directions. Moreover, these nanorods are used to enhance the vibrational signals of organic molecules in the mid-infrared region with remarkable enhancement factors.

11:40 : Invited talk**Plasmonic chirality of one-dimensional nanostructures: the role of lattice resonance****Wei Zhang***Institute of Applied Physics and Computational Mathematics (China)*

We perform systematic studies of one-dimensional (1D) chains of twisted nanorod dimers, focusing on the collective effect. Our studies reveal that the interplay between local structure/near field interaction and collective effect/far field interaction leads to quite different optical activity than that of the local nanostructures. In particular, it is found that the one-dimensional arrays of achiral objects show chiral responses due to the collective effect. Our studies provide useful guidance for the design of sensors based on optical activity.

12:00 : Invited talk

Plasmonic colour printing for information encryption

Zhang-Kai Zhou, Jiancai Xue

Sun Yat-sen University (China)

Plasmonic structures can selectively absorb or reflect light, making them can provide on-demand structural colours for building image devices. Based on our studies about plasmonic printing of interference colours, herein we introduce a new nano-steganography technique which is based on the optical inference of a plasmonic system. By carefully controlling the structural perturbation in such system, information encryption with near-perfect hiding and the property of countersurveillance can be obtained. Also, Information encryption based switchable metasurface imaging will be introduced.

12:20 : Electron Dynamics in Plasmons

Hue T. B. Do¹, Wen Jun Ding², Zackaria Mahfoud², Lin Wu², Michel Bosman¹

*¹National University of Singapore (Singapore), ²A*STAR (Singapore)*

We show that the Particle-in-Cell (PIC) simulation method can be used to robustly describe plasmon resonances, with a unique emphasis on the motion of the electrons in the time domain. Laser-excited and electron-beam-excited plasmons are studied to obtain the femtosecond time-scale dynamics of electrons in plasmons, including the plasmon dephasing, the electron kinetics during damping and the evolution of plasmons and electrons during interaction with an electron beam.

10:20 - 12:20 — Rene Descartes Room

Session 1A11

Novel Topological Photonic Materials

Organized by: Matthias Saba and Sang Soon Oh

Chaired by: Matthias Saba and Sang Soon Oh

10:20 : Invited talk

Topological Singular Points in Photonic Crystals with Broken Symmetry

Masaya Notomi

Tokyo Institute of Technology (Japan)

We demonstrate that it is possible to deterministically generate topologically-protected bound states in the continuum (BIC) by breaking C₆ symmetry of triangular-lattice photonic crystals. Furthermore, we found a variety of pair-creation and annihilation processes of circularly-polarized states, which are another type of singular points, by symmetry breaking. We also clarify that there is an intriguing conservation rule of two different topological charges, which account for vectorial nature of these singularities.

10:40 : Invited talk

Hidden symmetry enforced nexus points of nodal lines in layer-stacked dielectric photonic crystals

Ruo-Yang Zhang¹, Zhongfei Xiong², Yuntian Chen², Che Ting Chan¹

¹Hong Kong University of Science and Technology (Hong Kong), ²Huazhong University of Science and Technology (China)

In an AB-layer-stacked photonic crystal consisting of anisotropic dielectrics, we discover that the unique photonic band connectivity leads to triply degenerate nexus points of two nodal rings and a Kramers-like nodal line. The emergence and intersection of the line nodes are guaranteed by a hidden symmetry of Maxwell's equations. The bands with a constant k_z and the iso-frequency surfaces nearby the nexus point both disperse as a spin-1 Dirac-like cone, indicating exotic transport features of light at nexus point.

11:00 : Invited talk

Three-dimensional metamaterials and two-dimensional photonic crystals for topological photonic phase

Minkyung Kim, Junsuk Rho

Pohang University of Science and Technology (Korea)

My recent work on 3D topological semimetals based on metamaterials and 2D topological insulators based on photonic crystals will be presented in this talk. In the first part, topological semimetals whose topological phases arising from effective optical properties, hyperbolicity and chirality, will be covered. In the latter part, exploration of 2D nontrivial topology characterized by quantum Hall phase and Zak phase will be presented.

11:20 : Invited talk

Nodal link in double diamond photonic crystal

Haedong Park¹, Stephan Wong¹, Xiao Zhang², Sang Soon Oh¹

¹Cardiff University (United Kingdom), ²Sun Yat-sen University (China)

We demonstrate nodal links in momentum space using a dielectric double diamond structure. We also characterize the topological natures of the nodal links. These topological natures are summarized into non-Abelian charges.

11:40 : Invited talk

Moiré superlattice induced giant gauge field and Landau levels in bilayer Metacrystal

Wenhui Wang¹, Wenlong Gao², Yuanjiang Xiang³, Shuang Zhang²

¹University of Fribourg (Switzerland), ²University of Birmingham (United Kingdom), ³Hunan University (China)

We report the first experimental observations of the Landau level flat bands and the associated eigen states in photonics bi-layer metacrystals. The moiré pattern formed by the lattice constant mis-match in the metacrystals is shown to create huge effective gauge field resulting in the photonic flat-band Landau levels. The photonic Landau levels are measured by the microwave near field scanning system and agrees with the theoretical and numerical predictions excellently.

12:00 : Invited talk

Topological photonics: Mistaken paradigms and new opportunities

Aitzol Garcia-Etxarri¹, Maria Blanco de Paz¹, Chiara Devescovi¹, Matt Proctor², Paloma Arroyo Huidobro³, Barry Bradlyn⁴, Maia Garcia Vergniory¹, Dario Bercioux¹

¹Donostia International Physics Center (Spain), ²Imperial College London (United Kingdom), ³Instituto Superior Tecnico (Portugal), ⁴University of Illinois at Urbana-Champaign (USA)

In this work, through the application of the method of "Topological Quantum Chemistry"(TQC) to photonic crystals and the numerical calculation of Wilson loops and different topological invariants we will introduce a variety of novel topological effects in 2D and 3D photonic crystals. For instance, we will present the first instance of fragile topology in a photonic system, higher order photonic TI sustaining topologically protected corner states and our latest advances in the design of 3D topological photonic crystals.

Lunch

12:30 - 14:00

14:00 - 15:25 — Victor Veselago Room

Session 1A12

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

14:00 : Invited talk

Tailoring the emission and the photodynamics of quantum emitters with high index dielectric nanostructures

Peter Wiecha¹, Mélodie Humbert¹, Clément Majorel¹, Nicolas Mallet¹, Bruno Masenelli², Gérard Colas des Francs³, Franck Fournel⁴, Vincent Larrey⁴, Aurélie Lecestre¹, Guilhem Larrieu¹, Arnaud Arbouet¹, Christian Girard¹, Vincent Paillard¹, Aurelien Cuche¹

¹Université de Toulouse (France), ²Université de Lyon (France), ³Université Bourgogne-Franche Comté (France)

ce), ⁴ *Université Grenoble Alpes (France)*

We show both experimentally and theoretically that the photodynamics of several electric and/or magnetic quantum emitters can be controlled by high index dielectric nanostructures made of Silicon.

14:20 : Tailoring the plasmonic properties of metals: The case of substoichiometric titanium nitride

A. Catellani, A. Calzolari

CNR-NANO Istituto Nanoscienze (Italy)

We present a first principles study of the composition-plasmonic properties interplay in sub-stoichiometric TiNx crystals, which conjugate superior structural properties and tunable optical response. We investigate the stability of defective TiNx over a large range of vacancy concentrations, and we describe the microscopic character and the plasmonic properties of these systems. Our results indicate a clear trend in the modification of the plasmonic activity of TiNx as a function of nitrogen concentration.

14:35 : Invited talk

Ultrafast all-optical magnetic writing in transparent dielectrics

Andrzej Stupakiewicz

University of Bialystok (Poland)

We demonstrated the all-optical photo-magnetic recording in Co-ions doped iron garnet films using a time-resolved magneto-optical spectroscopy and single-shot ultrafast imaging of magnetic domains. Ultrashort laser pulses at near-infrared range it is possible to write and rewrite magnetic bits with a frequency of up to 20 GHz, with the maximum repetition rate being defined by the frequency of ferromagnetic resonance in the field of photoinduced magnetic anisotropy.

14:55 : Keynote talk

Strainoptronics: A New Degree of Freedom for 2D Material Device Engineering

Rishi Maiti, Hamed Dalir, Volker J. Sorger

George Washington University (USA)

Here we introduce "strainoptronics the local strain engineering of 2D materials for novel optoelectronic components. We exemplarily demonstrate heterogeneously integrating 2D materials in photonic circuits thus realizing a photodetector featuring a strong photoresponse (responsivity 0.5 A/W) operating at 1550 nm in silicon photonics.

14:00 - 15:20 — Allan Boardman Room

Session 1A13

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

14:00 : Invited talk

Broadband light beaming from a helical traveling-wave nanoantenna

Mengjia Wang, Roland Salut, Miguel Suarez, Nicolas Martin, Thierry Grosjean

FEMTO-ST Institute (France)

Nanoantennas have attracted much interest for their ability to directionally radiate light, with important implications in the optical detection of nanoscale objects. Based on the control of the resonant or dispersive properties of nanostructures, nanoantenna directivity is strongly wavelength-dependent. Here, we show that invariant light beaming can be obtained over a broad spectral range from an individual helical traveling-wave nanoantenna. This plasmonic antenna results from the extension to optics of the low-frequency helical antenna operating in the "axial mode".

14:20 : Invited talk

Active optical antennas driven by hexagonal boron nitride tunnel junctions**Kai Braun, Lukas Jakob, Frank Wackenhut, Florian Laible, Monika Fleischer, Alfred J. Meixner***University of Tuebingen (Germany)*

The conversion of an electrical signal to an optical signal with maximum bandwidth and practical on-chip integration is of fundamental interest. Here we present active optical antennas driven by inelastic electron tunneling. We have fabricated arrays of vertical coupled antennas in metal-insulator-metal arrangements based on electrically connected gold nano rods (GNR) and hexagonal boron nitride as insulating layer. By tuning the aspect ratio of the GNRs we shift the localized plasmon resonance of the antennas and adjust the emission wavelength

14:40 : Invited talk**Plasmonic and Photonic Catalysis****Emiliano Cortes***Ludwig-Maximilians-Universitat (Germany)*

Optical modes engineering in metallic and dielectric nanoparticles could open new paths for assisting chemical transformations using sunlight. In recent years, we have investigated these phenomena at the single nanoparticle level in order to unravel the mechanisms inducing catalytic transformations at these illuminated interfaces. Gaining a nanoscopic insight of these processes and their interplay could aid in the rational design of novel plasmonic and photonic photocatalysts.

15:00 : Invited talk**Modeling plasmonic hot-electron generation and their role in photocatalysis****Lucas Vazquez Besteiro¹, Zhiming M. Wang², Alexander O. Govorov³***¹Universidad de Vigo (Spain), ²University of Electronic Science and Technology of China (China), ³Ohio University (USA)*

Plasmonic nanoparticles can excite high-energy electrons capable of traversing into neighboring materials and driving secondary processes in photosensing and photocatalytic applications. In this talk we will describe strategies to enhance the excitation of these high-energy carriers, and discuss their utility in several technological contexts. Our discussion will be supported by results from a hybrid theoretical model combining a quantum-mechanical description of the electronic states and a classical electrodynamic calculation of the collective plasmonic modes of large plasmonic nanocrystals.

14:00 - 15:20 — Tatsuo Itho Room**Session 1A14****Bottom-up approaches, new fabrication routes and ENSEMBLE3**

Organized by: Dorota Pawlak and Virginie Ponsinet

Chaired by: Dorota Pawlak and Virginie Ponsinet

14:00 : Invited talk**Block copolymer directed nanoarchitectures for the design of novel optical materials****Alberto Alvarez Fernandez¹, Barry Reid¹, Guillaume Fleury², Virginie Ponsinet², Stefan Guldin¹***¹University College London (United Kingdom), ²Université de Bordeaux (France)*

Several types of block copolymers have been used as templates to produce well-defined dielectric and metallic nanostructures in 3D or 2D, with high degree of order and tunability. Resulting structures have shown interesting optical properties such as high-index metasurfaces or antireflective coatings.

14:20 : Invited talk**Photonic nanomaterials by self-assembly of block copolymers****Tapio Niemi***Tampere University (Finland)*

Both the scientific community and the industry are actively investigating innovative means to overcome

the fundamental limitations of conventional nanolithography. One emerging technique is based on directed self-assembly of block copolymers. Besides the possibility to realize extremely small feature sizes, the self-assembled polymers enable fabrication of exotic nanostructures, which are challenging or even impossible to fabricate by other methods. I shall demonstrate few applications for etch masks, refractive index tuning and preparation of multi-material nanostructures.

14:40 : Invited talk

Non-fading Plasmonic Color Printing through Laser Processing of Semicontinuous Metal films

Piotr Nyga¹, Sarah N. Chowdhury², Zhaxylyk Kudyshev², Esteban Garcia², Alexei S. Lagutchev², Alexander V. Kildishev², Vladimir M. Shalaev², Alexandra Boltasseva²

¹Military University of Technology (Poland), ²Purdue University (USA)

We report non-fading generation of vibrant colors through a femtosecond laser post-fabrication processing of a plasmonic semicontinuous metal film deposited on a metallic mirror coated with a sub-wavelength-thick dielectric spacer. Long term stability of color is obtained through structures' overcoating with a dielectric layer. Local changes induced to nanostructures of semicontinuous film are controlled by the femtosecond laser parameters, especially fluence. Wide range of vibrant colors in reflection mode from blue to green, to red can be easily obtained.

15:00 : Invited talk

Fabrication of amorphous photonic materials by colloidal self-assembly

Joshua Ricouvier, Romain Pierrat, Remi Carminati, Patrick Tabeling, Pavel Yazhgur

ESPCI Paris (France)

Recently discovered well-engineered disordered photonic band gap materials can potentially enable the production of energy-efficient displays or freeform wave-guides. In our research, we propose a colloidal self-assembly approach to produce such amorphous materials. By drying bidispense 2D emulsions, we produce a foam-like material with extremely low long-range density fluctuations (hyperuniformity) and strong short-range order, which open a disordered PBG. The obtained results suggest that photonic foams are potentially excellent candidate for disordered 3D PBG materials.

14:00 - 15:20 — Christian Huygens Room

Session 1A15

Symposium III: Advanced passive and active metasurfaces and zero-index optics

Organized by: Howard Lee, Yu-Jung (Yuri) Lu and Zi Jing Wong

Chaired by: Howard Lee, Yu-Jung (Yuri) Lu and Zi Jing Wong

14:00 : Invited talk

Bound states in the continuum in metasurfaces with dipolar meta-atoms

Diego R. Abujetas¹, Juan J. Saenz², Jose A. Sanchez-Gil¹

¹Instituto de Estructura de la Materia (CSIC) (Spain), ²Donostia International Physics Center (Spain)

We explore the emergence of bound states in the continuum (BICs) in metasurfaces consisting of dipolar meta-atoms. We derive a coupled dipole theoretical formulation to describe the optical properties of a periodic array consisting of several electric/magnetic dipoles per unit cell, thereby investigating robust symmetry-protected BICs through different mechanisms in various kinds of arrays of interest throughout the electromagnetic spectrum: high-refractive-index disks in the GHz, metallic rod dimers in the THz, and silicon nanodisks in the visible.

14:20 : Invited talk

Nonlinear interactions of guided plasmons in graphene metasurfaces

Joel Douglas Cox

University of Southern Denmark (Denmark)

We explore nonlocal and quantum finite-size effects in the propagating plasmons supported by 1D graphene

nanoribbons and their associated nonlinear response, revealing a strong dependence on such phenomena for narrow ribbons excited by optical fields carrying large in-plane momenta. Our findings suggest new possibilities to engineer nonlinear interactions among guided plasmons in graphene-based nonlinear metasurfaces.

14:40 : Invited talk

Silicon based metastructures and metasurfaces

Ahmed Mahmoud

American University in Cairo (Egypt)

Throughout the past two decades, the concept of metamaterials introduced itself as a powerful tool for achieving non-conventional light-matter interaction in numerous scenarios. Despite the research efforts and the concurrent discoveries, the difficulty of fabrication and the incompatibility of the structures to conventional CMOS environments limit the influence of such efforts. In this work we explore various tunable metastructures and metasurfaces within our research efforts towards achieving the functionalities and phenomena of metamaterials within conventional environments and materials.

15:00 : Invited talk

Single-Emitter Near-Field Excited Quantum Dynamics in Near-Zero-Index Materials

Frank Bello¹, Nuttawut Kongsuwan², John F. Donegan¹, Ortwin Hess¹

¹*Trinity College Dublin (Ireland)*, ²*Quantum Technology Foundation (Thailand)*

We investigate the quantum dynamics from single quantum emitters (QE) excited using near-field light from a plasmonic transducer. Results show the ability to strongly drive a QE using a plasmonic near field with an ultrafast single-photon emission rate of 101-102 fs⁻¹ for emitters embedded in near-zero index (NZI) media. QEs embedded in NZI media also demonstrate excellent enhancement of spontaneous emission compared to those in free space ($\approx 10^7$). Comparisons are made using QEs embedded in other common materials such as Silicon.

14:00 - 15:15 — Augustin Fresnel Room

Session 1A16

Symposium IV: Chirality, magnetism, and magnetoelectricity: Separate phenomena and joint effects in metamaterial structures

Organized by: Eugene Kamenetskii

Chaired by: Eugene Kamenetskii

14:00 : Invited talk

3D FDTD-LLG modelling of magnetisation dynamics in thin film ferromagnetic structures

Feodor Ogrin

University of Exeter (United Kingdom)

Here I propose a model which uses 3D finite-difference-time-domain (FDTD) approach together with Landau-Lifshits-Gilbert (LLG) formalism to find the exact solutions for magnetisation dynamics in thin film ferromagnetic structures. Two case studies are demonstrated, in which the model is validated against analytical and experimental methods.

14:20 : Invited talk

Realization of Dynamic Electromagnonic Microwave Crystal via Magnon-Photon Coupling in Artificial Multiferroic Heterostructure

Alexander A. Serga¹, Alexey B. Ustinov², Andrey A. Nikitin², Andrei V. Drozdovskii², Alexander A. Semenov², Dmytro A. Bozhko³, Burkard Hillebrands¹, Erkki Lähderanta⁴, Boris A. Kalinikos²

¹*Technische Universität Kaiserslautern (Germany)*, ²*St. Petersburg Electrotechnical University (Russia)*,

³*University of Colorado at Colorado Springs (USA)*, ⁴*LUT-University (Finland)*

Magnon transport in time-dependent periodic spatial potentials-dynamic magnonic crystals-paves a way to energy-efficient data transfer and information processing. Voltage control of magnon currents promises to be

fast and low energy-consuming. It can be achieved in artificial multiferroic heterostructures, where the strong coupling of magnons and microwave photons constitutes new quasiparticles called electromagnons. Here, we report on the experimental realization of a voltage-controlled dynamic electromagnonic crystal operating at microwave frequencies.

14:40 : Invited talk

Geometrical symmetry breaking in nanomagnets

Oleksandr V. Pylypovskyi

Helmholtz-Zentrum Dresden (Germany)

Three-dimensional ferro- and antiferromagnetic nanoarchitectures possess a special interplay between their geometrical (topological) properties and the magnetic order parameter. The emergent chiral and anisotropic responses extend the intrinsic material properties and pave the way to novel functionalities of spintronic and spin-orbitronic devices.

15:00 : The role of plasmonic excitations in the far-field configurational chirality

leeju Singh¹, Shumuel Sternklar², Yuri Gorodestki¹

¹Ariel University (Israel), ²Ariel University (Israel)

Broken mirror symmetry of chiral structures imposes a lack of mirror symmetry in the scattering profile. When an energy dissipation channel is introduced in the system, an overall optical activity arises. Plasmonic nanostructures, therefore, are ideal platform to induce optical activity by means of constitutional or configurational chirality. We experimentally investigate mechanism of the plasmonically-induced configurational chirality in a periodic monoclinic hole array with a broken mirror symmetry.

14:00 - 15:20 — Ibn Al-Haytham Room

Session 1A17

Symposium V: Phononics and acoustic metamaterials

Organized by: Jensen Li and Guoliang Huang

Chaired by: Jensen Li and Guoliang Huang

14:00 : Invited talk

Taking inspiration from polymer chemistry: Tacticity in chiral phononic crystals

Andrea E. Bergamini¹, Marco Miniaci², Domenico Tallarico¹, Gwenael Hannema¹, Bart Van Damme¹, Armin Zemp¹

¹Empa, Materials Science and Technology (Switzerland), ²IEMN (France)

In this contribution, we will discuss a more complete picture of the constraints imposed by engineering considerations to the design of phononic materials, how the exploitation of chirality and tacticity as design feature is helping us tackling some of these issues, as well as answering some of the interesting physical questions that we encountered along the way.

14:20 : Invited talk

Valley-protected Topological Lamb Waves in Asymmetric Pillared Metamaterials

Bernard Bonello¹, Wei Wang¹, Bahram Djafari-Rouhani², Yan Pennec², Yabin Jin³

¹Sorbonne Université (France), ²Université de Lille - IEMN (France), ³Tongji University (China)

We present a numerical study of the valley-protected topological propagation of zero-order antisymmetric (A0), symmetric (S0), and shear horizontal (SH0) Lamb waves at different domain walls between topologically distinct asymmetric double-sided pillared phononic crystals. The topological phase transition is obtained by imposing two large space inversion symmetry breaking perturbations on the height of some pillars in the unit cell. We demonstrate the unidirectional transport of A0, S0, and SH0 modes at different domain walls in straight or Z-shaped guides.

14:40 : Invited talk

fiber-integrated microcavities for efficient generation of coherent acoustic phonons

Martin Esmann¹, Omar Ortiz², Florian Pastier³, Anne Rodriguez², Priya Priya², Aristide Lemaître², Carmen Gomez-Carbonell², Isabelle Sagnes², Abdelmounaim Harouri², Pascale Senellart², Valérian Giesz³, Norberto Daniel Lanzillotti-Kimura²

¹ *Université Paris-Saclay (Germany)*, ² *Université Paris-Saclay (France)*, ³ *Quandela SAS (France)*

Coherent phonon generation by picosecond optical pump-probe spectroscopy is an important experimental tool for studying acoustic properties at the nanoscale. In this work, we integrate semiconductor micropillar cavities confining near-infrared light and 18 GHz acoustic phonons with single-mode fibers. This approach solves a major challenge of existing pump-probe experiments using mechanical delay lines: maintaining the spatiotemporal overlap of pump and probe beams on a micro-object to achieve well-controlled, reproducible mode matching conditions.

15:00 : Invited talk**Lightweight three-dimensional metamaterials for omnidirectional attenuation of mechanical waves and vibrations**

Anastasiia O. Krushynska, Monica Acuautla

University of Groningen (The Netherlands)

Tailored composites with frequency band gaps are promising for mechanical wave control and vibration mitigation. Despite advances in their developments, the generation of omni-directional broadband band gaps remains challenging. We propose a class of lightweight phononic structures addressing this challenge. Omnidirectional performance is achieved by simultaneously activating several wave scattering mechanisms. Numerically estimated strong attenuation at ultrasound frequencies is validated in transmission experiments on polymeric 3D-printed samples.

14:00 - 15:30 — Gaston Floquet Room**Session 1A18****Plasmonics and nano-optics****14:00 : CdZnO nanoparticle coating on GaAs for IR field enhancement**

Eduardo Martinez Castellano¹, Julen Tamayo-Arriola¹, Miguel Montes Bajo¹, Oleksii Klymov², Carmen Martinez-Tomas², Said Agouram², Elias Munoz¹, Vicente Munoz-Sanjose², Adrian Hierro¹

¹ *Universidad Politécnica de Madrid (Spain)*, ² *Universitat de València (Spain)*

In this work, we study the plasmonic response of CdZnO nanoparticles grown on GaAs in the mid-IR range of the spectrum. We measured the transmittance of the system and evaluated the field-enhancement profile of the system, identifying two plasmonic modes. Finally, we briefly discussed the potential applicability of this NP coating in photonic devices, once demonstrated the penetration of the electrical field within the substrate supporting the NPs.

14:15 : Demonstration and Tuning of Tamm Plasmons at the Interface with Metasurfaces

Oleksandr Buchnev¹, Alexandr Belosludtsev², Victor Reshetnyak³, Dean R. Evans⁴, Vassili A. Fedotov¹

¹ *University of Southampton (United Kingdom)*, ² *Center for Physical Sciences and Technology (Lithuania)*, ³ *Taras Shevchenko National University of Kyiv (Ukraine)*, ⁴ *Air Force Research Laboratory (USA)*

We demonstrate experimentally that Tamm plasmons can be supported by a dielectric mirror interfaced with a metasurface, a discontinuous metal film patterned on the sub-wavelength scale. Not only do Tamm plasmons survive the nano-patterning of the metal film, but they also become sensitive to external perturbations as a result. In particular, by depositing a nematic liquid crystal on the outer side of the metasurface we were able to red-shift a Tamm plasmon by 35 nm.

14:30 : Low-melting-point borophosphate glass as a matrix for NP-doped, luminescent composites produced using the NPDD method

Rafal Nowaczynski¹, Marcin Gajc², Hancza Surma², Piotr Paszke³, Kamil Szlachetko³, Piotr Piotrowski³, Dorota Anna Pawlak²

¹ *Warsaw University of Technology (Poland)*, ² *Institute of Electronic Materials Technology (Poland)*, ³ *University*

of Warsaw (Poland)

NanoParticle Direct Doping is a method developed in the Institute of Electronic Materials Technology in Warsaw that allows fabrication of volumetric composites based on glass matrices doped with various kinds of nanoparticles, including metallic plasmonic NPs, Quantum Dots and rare-earth ions. It allows us to obtain photoluminescent materials co-doped with different types of NPs, varying in size and composition, in which it is possible to observe effects such as plasmonic enhancement of the excitonic emission or radiative energy transfer.

14:45 : Optical Probing of Plasmonic Hot Electron Occupancies

Zsuzsanna Papa¹, Judit Budai², Shirly Espinoza³, Mateusz Rebarz³, Martin Zahradnik³, Péter Dombi¹
¹Wigner Research Center for Physics (Hungary), ²ELI-ALPS (Hungary), ³ELI Beamlines (Czech Republic)

We discuss the in-depth distribution and time evolution of hot electrons generated upon the excitation of surface plasmon polaritons. We applied ellipsometry to measure the dielectric function of plasmonic systems and to determine their electron distribution. Applying cw illumination, the spatial location of hot electrons can be deduced. Pump-probe approach with <100fs resolution enabled us to measure electron distributions belonging to stages of plasmon decay when energetic electrons are generated, scattered among each other and interact with the lattice.

15:00 : Plasmons in nanostructured and corrugated 2D materials

Bruno Majérus¹, Peter Vancso², Levente Tapasztó², Luc Henrard¹

¹University of Namur (Belgium), ²Hungarian Academy of Sciences (Hungary)

In this work we theoretically investigate plasmonic excitations in 2D materials. We suggest that localized plasmons in corrugated graphene increase the surface enhanced Raman spectroscopy (SERS) response of given molecules. We also show that mirror twin boundaries (MTB) in transition metal dichalcogenides (TMDs) can sustain plasmon due to their local metallic behavior as recently experimentally highlighted.

15:15 : Quantum Dot Photoluminescence Enhancement in GaAs Nanopillar Oligomers Driven by Collective Magnetic Modes

Damir Yagudin¹, Maria Kroychuk¹, Maxim Rakhlin², Aidar Galimov², Alexey Toropov², Tatiana Shubina², Grigorii Klimko², Alexander Shorokhov¹, Andrey Fedyanin¹

¹Lomonosov Moscow State University (Russia), ²Ioffe Institute (Russia)

Resonant nanostructures made of high-index dielectrics are an auspicious platform for controlling of light at the nanoscale. In this paper, we study a GaAs nanopillar oligomer system with embedded InAs quantum dots. Our experiments and calculations show that excitation of Mie type resonances in such structures leads to the enhancement of the photoluminescence from quantum emitters due to a spectral overlap of the emitter absorption or emission spectra with the collective magnetic resonance of the cluster.

14:00 - 15:20 — Lawrence Bragg Room

Session 1A19

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

14:00 : Invited talk

Latent symmetries and their application in Physics

Malte Rontgen

University of Hamburg (Germany)

I will give an overview on latent symmetries, a new class of symmetries in systems described by matrix equations. Although these symmetries are not apparent from a geometric inspection of the system, they have powerful effects such as inducing symmetries of the eigenstates' components or even degeneracies in the eigenvalue spectrum. Their study thus allows to gain knowledge about the system's structure that remains hidden from a direct observation.

14:20 : Invited talk**Mimicking Magnetic Localized Surface Plasmons With High-Index Dielectrics: Enhancing Magnetic Resonance Imaging Signal-To-Noise Ratio****Carlo Rizza, Elia Palange, Angelo Galante, Marcello Alecci***University of L'Aquila (Italy)*

We show that surface waves supported by a negative permeability sphere (magnetic localized surface plasmons, MLSPs) can be well reproduced by the electromagnetic field scattered by a low-loss high-index dielectric sphere. In the magnetic resonance imaging (MRI) context, we suggest the use of these spoof MLSPs to increase MRI efficiency. More precisely, we show that the MLSPs, coupled to a standard radio-frequency surface coil, are able to enhance the MRI signal-to-noise ratio with respect to standard setups.

14:40 : Invited talk**Full-Wave Formalism for Soliton Propagation in Nonlinear Photonic Crystals****Vakhtang Jandieri¹, Ramaz Khomeriki², Douglas Werner³, Jamal Berakdar⁴, Daniel Erni¹**¹*University of Duisburg-Essen (Germany)*, ²*Tbilisi State University (Georgia)*, ³*The Pennsylvania State University (USA)*, ⁴*Martin-Luther Universität (Germany)*

The propagation of the temporal solitons in Kerr-type photonic crystal waveguides is investigated theoretically and numerically. An expression describing the evolution of the envelope of the soliton, which is based on full-wave modal analysis taking into account all space-harmonics, is rigorously obtained. The nonlinear coefficient is derived, for the first time, based on a modification of the refractive indices for each space-harmonic due to the Kerr-type nonlinearity.

15:00 : Invited talk**All dielectric 3D periodic nanoresonators for phase and polarization control of SH light****Giuseppe Marino¹, Carlo Gigli¹, Davide Rocco², Luca Carletti², Isabelle Sagne³, Stephan Suffit¹, Costantino De Angelis², Giuseppe Leo¹**¹*Université de Paris (France)*, ²*University of Brescia (Italy)*, ³*Universite Paris-Saclay (France)*

We demonstrate numerically and experimentally enhanced efficiency and polarization control of nonlinear mixing via 3D all-dielectric periodic nanostructures. The 3D spatial periodicity allows diffractive interferences of SH light for the phase control, while the meta-atom geometry allows the excitation of Mie-modes for polarization control.

14:00 - 15:15 — Rene Descartes Room**Session 1A20****Novel Topological Photonic Materials**

Organized by: Matthias Saba and Sang Soon Oh

Chaired by: Matthias Saba and Sang Soon Oh

14:00 : Topological scattering in Photonic Weyl metamaterial**Wenlong Gao¹, Hua Cheng¹, Yangang Bi², Hongbo Sun³, Jianguo Tian⁴, Shuqi Chen⁴, Shuang Zhang¹**¹*University of Birmingham (Germany)*, ²*University of Birmingham (United Kingdom)*, ³*Jilin University (China)*, ⁴*Nankai University (China)*

Weyl points are the sources or drains of Berry flux. Here we experimentally observe the topological signature in the scattering of the photonic Weyl points. Specifically, we discover a helical phase distribution in the angled-resolved reflected waves, representing a winding phase in the momentum space, which leads to an angular momentum beam in the reflection. Our findings pave way for potential device level applications to angular momentum beams with photonic Weyl systems.

14:15 : Topological Kagome Lattice Laser**Yongkang Gong, Stephan Wong, Anthony J. Bennett, Diana L. Huffaker, Sang Soon Oh***Cardiff University (United Kingdom)*

Discovery of new types of photonic topological platforms facilitates the development of novel topological lasers that are potentially immune to cavity defects with high lasing efficiency. We propose a topologically protected laser cavity based on broadband nontrivial edge states in semiconductor photonic crystals with Kagome lattice, and investigate the robustness and dynamics of the lasing cavity.

14:30 : Topological nanophotonics with time-reversal-invariant plasmonic lattices

Paloma A. Huidobro

University of Lisbon (Portugal)

Topological nanophotonics offers a promising path towards the robust control of photons at the nanoscale by exploiting topologically protected boundary modes that are immune to a range of defects and imperfections. I will focus on 1D and 2D realisations of topological boundary modes in arrays of plasmonic nanoparticles.

14:45 : Transport and localization of polaritons in Semi-Dirac honeycomb lattices

Bastian Real¹, Omar Jamadi¹, Marijana Milicevic², Nicolas Pernet², Tomoki Ozawa³, Gilles Montambaux², Isabel Sagnes², Aristide Lamaitre², Luc Le Gratiet², Abdelmounaim Harouri², Sylvain Ravets², Jacqueline Bloch², Alberto Amo¹

¹ *Université de Lille (France)*, ² *Université Paris-Saclay (France)*, ³ *Tohoku University (Japan)*

Strain strongly affects the transport and localisation properties of graphene. For a critical uniaxial compression, graphene shows a semi-Dirac cone with massless and massive dispersions along perpendicular directions. Here we implement strained polariton honeycomb lattices to evidence the highly anisotropic transport of polaritons and to observe directional vacancy states with chiral symmetry. Our work paves the way for the study of transport and localisation in chiral lattices with exotic Dirac dispersions.

15:00 : Vorticial Phase Mirror in Photonic Weyl metamaterial

Wenlong Gao¹, Hua Cheng¹, Yangang Bi¹, Wenwei Liu², Zhancheng Li², Qinghua Guo¹, Yang Yang¹, Oubo You¹, Jing Feng³, Hongbo Sun³, Jianguo Tian², Shuqi Chen², Shuang Zhang¹

¹ *University of Birmingham (United Kingdom)*, ² *Nankai University (China)*, ³ *Jilin University (China)*

Weyl points are the sources or drains of Berry flux. Here we experimentally observe the topological signature in the scattering of the photonic Weyl points. Specifically, we discover a helical phase distribution in the angled-resolved reflected waves, representing a winding phase in the momentum space, which leads to an angular momentum beam in the reflection. Our findings pave way for potential device level applications to angular momentum beams with photonic Weyl systems.

Coffee Break and Exhibit Inspection

Session 1P2

Poster session II

15:20 - 16:00

P1: THz Metamaterial Device Design with SRRs

Brinta Chowdhury, Abdullah Eroglu

(USA)

THz device design method for periodic structures using single and double gap split ring resonators (SRRs) is introduced. The method involves implementation of split ring resonators with application of network parameters. The closed form relations are developed and verified with electromagnetic and circuit simulators. The prototype then has been built and measured. The results are compared, and agreement has been observed between analytical, simulation and measurement results.

P2: Design, Simulation and Fabrication of Ultra Wide Band Microstrip Patch Antenna for Wireless Application

Suleiman Babani, Zainab Yunusa, Umar Musa, Abubakar Sani Ali

Bayero University Kano (Nigeria)

Ultra wideband (UWB) technology is one of the most widely technologies used in wireless communication system to support the application that required high data rate and high speed. The purpose of this work

is to design and fabricate ultra-wideband microstrip patch antenna for wireless application with optimum performance such as wide bandwidth, good matching impedance, small antenna size exhibits and its E and H-plane radiation patterns are stable over the UWB frequency range.

P3: Focus Beam System Biaxial Cross-Polarization Approximation Technique

Nicholass Alan O’Gorman

Air Force Institute of Technology (USA)

When testing non-isotropic metamaterials, the axes of the sample being measured, and the antenna being used need to align in order to calculate the material parameters. If the axes of the sample are unknown or the sample was not aligned properly, the resulting data will not result in accurate measurements. This paper discusses a method for the focus beam system to calculate the material parameters for a biaxial sample without knowing the axis orientation.

P4: Spherical symmetric pure and hybrid electromagnetic anapoles based on cartesian electric-current multipole expansion

Vishal Vashista¹, Andriy Serebryannikov¹, Matti Kaivola¹, Maciej Krawczyk²

¹Aalto University (Finland), ²Adam Mickiewicz University (Poland)

In this work, we have proposed a plasmonic realization of fundamental anapole using three nanorods structure and studied its properties. With our design, we are able to suppress the far-field radiation by a factor of 1000 compared to a dipole source with the same integrated amplitude of the electric current density.

P5: Broadband vectorial ultra-flat optics with up to 99 % experimental efficiency in the visible

Arturo Burguete-Lopez, Fedor Getman, Makarenko Maksim, Andrea Fratolocchi

KAUST (Saudi Arabia)

We propose a methodology that allows the production of high efficiency (up to 99 % in the visible) ultra-flat (down to 50 nm thick) optics for vectorial light control and for arbitrarily defined broadband input-output responses of a desired wavefront shape. Experimentally, we show basic transmission/reflection components such as polarizer beam splitters and dichroic mirrors can be manufactured with over 90 % efficiency across the visible and present the basis for a two sub-pixel flat optics display.

P6: Ultra-flat optics design platform for a high-efficient wavefront engineering

Maksim Makarenko, Fedor Getman, Arturo Burguete-Lopez, Andrea Fratolocchi

KAUST (Saudi Arabia)

In this work we propose a universal design platform for the development of high-efficient wavefront engineering structures. We demonstrate the efficiency of this approach by designing a series of common optical devices with an efficiency exceeding 99 %.

P7: Chiral Nanophotonics with Atomically Thin Semiconductors

T. V. Raziman, Rasmus H. Godiksen, Shaojun Wang, Moos Muller, Jaime Gomez Rivas, Alberto G. Curto

Eindhoven University of Technology (The Netherlands)

Spin-valley polarization in atomically thin semiconductors opens a new perspective to explore optical chirality. Here we summarize our progress on the exploitation of 2D semiconductors as sources and probes for chiral nano-optics. We show how competing optical transitions in few-layer materials allow a high degree of circular polarization. Second, in order to enhance polarization, we design achiral nanophotonic resonators that satisfy the conditions needed for improving chiral light emission as a path towards efficient sources of spin-valley-polarized light.

P8: Fabrication and Simulation of Photonic Crystals based on Silicon Nanopillars

Elena Lopez-Aymerich, Roger Lera-Leri, Zoilo Nunez-Lasus, Maria Dimaki, Winnie E. Svendsen, Sergi Hernandez, Daniel Navarro-Urrios, Mauricio Moreno, Florenci Serras, Albert Romano-Rodriguez

University of Barcelona (Spain)

In this paper we show successful nanofabrication of photonic crystals based on up to 3 μm long silicon nanopillars. Different nanopillars with 60 to 100 nm radius, a triangular distribution and pitch of 500 nm have been fabricated on silicon or Silicon-On-Insulator substrates. Simulations of the electromagnetic field distribution inside these devices have been carried out to ensure the correct behaviour of the samples as photonic crystals, waveguides and resonators for their use in sensing applications.

P9: Thermally tunable invisibility at terahertz: different mechanisms in one structure**Andriy E. Serebryannikov***Adam Mickiewicz University (Poland)*

Thermally tunable invisibility at terahertz frequencies is revisited with the focus on the contribution of different mechanisms achievable in different frequency and permittivity ranges. The mechanisms based on localized surface plasmon resonances and volume-mode resonances in high-index dielectric shells and others may co-exist for one scatter, enabling on-off switchable functionality.

P10: field characterization of nanometric gap-antennas for strong coupling**Maria Monserrat Alvarez Ortiz, Pawel Wozniak, Niek F. van Hulst***The Institute of Photonic Sciences (ICFO) (Spain)*

The field enhancement at the nanoscale is a powerful tool to achieve the strong coupling regime to control and maximize the emission of single-photon sources. Nanoantennas show strong optical response due to the excitation of localized surface plasmon resonances. Particularly strong resonances can be achieved by exploiting coupled resonators forming gap nanoantennas. Here, we investigate the properties of plasmonic dimer antennas of sub 10 nm separation.

P11: Surface plasmon-polariton in a hybrid structure of hyperbolic metasurface and phase-changing material**Olga Kharitonova¹, Igor Bychkov¹, Vladimir Shavrov², Dmitry Kuzmin¹***¹Chelyabinsk State University (Russia), ²Kotelnikov Institute of Radio-engineering and Electronics of RAS (Russia)*

In this work we theoretically investigate surface plasmon-polariton (SPP) in a hybrid structure of hyperbolic metasurface based on graphene and phase-changing material (VO₂). Introducing the PCM into hyperbolic metasurface leads to additional tunability of optical properties of the structure, which may be of great interest for numerous practical applications.

P12: Optical and photocatalytic properties of rare earth doped ZnS Nanocrystalline**Amel Tounsi, Djahida Talantikite-Touati, Roumaissa Khalfi, Hamid Merzouk***University of Bejaia (Algeria)*

New nanosized materials play a fundamental role in various industrial applications thanks their unique and functional properties. Moreover, in recent years, a great effort has been made to the design and control fabrication of nanostructured semiconductors such as zinc sulphide. So, much attention has been according in research to co-doped ZnS for improve the quality of ZnS thin films. We present in this work the preparation and characterization of ZnS and lanthanides doped ZnS prepared by co-precipitation method.

P13: Phase-shifted ellipsometric surface plasmon resonance sensor**Ibrahim Watad, Ibrahim Abdulhalim***Ben Gurion University of the Negev (Israel)*

We have demonstrated both experimentally and theoretically simple approaches to extract the light polarization changes induced by surface plasmon resonance [1-6]. Here, we present an ellipsometric approach based on phase-shifted ellipsometry with a design based on Kretschmann-Raether configuration both in the angular and spectral modes [5-6].

P14: Plasmonic Nanocavity Metasurface Based on Laser-Structured Silver Surface and Silver Nanoprisms for the Enhancement of Adenosine Nucleotide Photoluminescence**Oleg Yeshchenko¹, Vladislav Kudrya¹, Anastasiia Tomchuk¹, Igor Dmitruk¹, Nataliya Berezovska¹, Petro Teselko¹, Sergii Golovynskyi², Bin Xue², Junle Qu²***¹Taras Shevchenko National University of Kyiv (Ukraine), ²Shenzhen University (China)*

A photoluminescence spectroscopy and imaging of biomolecules at room temperature realized in the plasmonic cavity metasurface based on the Ag laser-induced periodic surface structure (LIPSS) and Ag triangular nanoprisms studied in the present work. The strong plasmon enhancement of photoluminescence of dAMP deposited on such metasurfaces observed. The plasmonic cavity metasurfaces consisting of metal LIPSS and metal nonspherical nanoparticles showed to be crucial for the highly sensitive detection and imaging of biomolecules at room temperature without any dye labels.

P15: Enhanced Electromagnetic Properties of Golden nanoparticle by hybrid film of Graphene in visi-

ble Frequency**Mohammad A. Abdollahi, Golamreza Moradi, Amir Nader Askarpour***Amirkabir University of Technology (Iran)*

This paper want to consider of Golden nanoparticle (NP) that covered by graphene layer. This made that NP excited in lower frequency. This excitation express in Surface Plasmon. So it should considered the properties of SPP resonance. Scattering Cross Section and Absorption Cross Section examined by FEM method in Comsol and results compared with calculation of Mie Expansion in Matlab application. At least the losses of the proposed structure checked

P16: Simulation of plasmon enhanced electric field and its impact on vibrational spectra**Tanguy Colleu-Banse, Vincent Liégeois, Luc Henrard***University of Namur (Belgium)*

Raman and infrared spectroscopy allow for an unambiguous identification of microscopic objects. Surface-Enhanced Spectroscopies improves its cross section by a coupling with plasmon excitations. However, the interpretation of resulting spectra is challenging. This works focuses on the enhancement of the local electromagnetic field associated with metallic nanodisks plasmon excitation. Simulation are obtained by Discrete Dipole Approximation (DDA). This local response is then combined with Time Dependent DFT (TD-DFT) to investigate its effect on the vibrational spectrum of hydrogen cyanide.

P17: Two-Directional Optical Binding Near One-Dimensional Photonic Crystal**Nataliia Kostina, Alexander S. Shalin***ITMO University (Russia)*

Here we consider optical forces acting on a pair of particles near 1D photonic crystal. The surface wave of such structure (BSW) provides stable optical binding for both polarizations of incident light. The distances between the stable positions of the particles are shown to be less than incident wavelength due to Bloch surface wave dispersion. These results demonstrate possibility of flexible optical manipulation of the arrays of nanoparticles.

P18: Gold nanostructures devices on flexible substrate for strain optical monitoring**Florian Lamaze¹, Abdelhamid Hmima¹, Nicolas Bercu², Thomas Maurer¹, Louis Giraudet², Julien Proust¹***¹University of Technology of Troyes (France), ²University of Reims Champagne-Ardennes (France)*

Gold nanoparticles dimers (GNDs) can be used as sensitive optical sensors for the detection of local nano-deformation. There is a growing interest for sensor technologies based on polydimethylsiloxane (PDMS) as flexible substrate, since it allows reversible strain. In this context, the present work tackles the development of organized gold nanopatterns on PDMS to improve the sensitivity of strain monitoring at nanoscale.

P19: Design of Ultra-Broadband and Compact Beamsplitter using Subwavelength Grating-based Multimode Interference Coupler**Kamalodin Arik, Mahmood Akbari, Amin Khavasi***Sharif University of Technology (Iran)*

An ultra-broadband and compact beam splitter is proposed using inherent anisotropy and dispersion property of a subwavelength grating (SWG)-based all dielectric metamaterial. These properties are used for designing a multimode interference (MMI) splitter. The designed 3dB SWG MMI splitter possess a bandwidth of more than 800nm, maintaining insertion loss below 1dB and a zero power imbalance. The overall footprint of the proposed device is highly small.

16:00 - 17:10 — Victor Veselago Room**Session 1A21****Plenary Session II****16:00 : Plenary talk****Metasurfaces for Light Management in Semiconductor Thin films**

Deirdre M. O'Carroll*Rutgers University (USA)*

In this talk, recent work on improving light trapping and light extraction in organic semiconductor thin films using plasmonic metasurfaces will be presented. Numerous optical phenomena, such as absorption induced scattering, out-of-plane waveguiding and morphology-dependent surface plasmon outcoupling, are identified due to exciton-plasmon coupling between the organic semiconductor and the metasurface.

16:35 : Plenary talk**Structuring Light and Dark with Metaoptics****Federico Capasso***Harvard University (USA)*

Metasurfaces can generate arbitrary vector beams. I will discuss devices that enable light's spin and orbital angular momentum (OAM) to evolve, simultaneously, from one state to another along the propagation direction and polarizing elements that virtually rotate their orientation as a function of the propagation distance. Advances in high OAM lasing will be reported along with the design and realization of two-dimensional phase and polarization singularities.

Break**17:10 - 17:30****17:30 - 19:05 — Victor Veselago Room****Session 1A22****Symposium II: New trends in nanophotonics and advanced materials**

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

17:30 : Invited talk**Plasmon dephasing and macroscopic polarization conversion in single nanoporous particles****Pritam Khan, Grace Brennan, Syed A. M. Tofail, Ning Liu, Christophe Silien***University of Limerick (Ireland)*

Plasmon damping strongly affects their temporal and spectral behaviour. Coupled and mesoscopic plasmonic systems having complex spectra generally impede damping analysis. As such, we report that polarimetric dark-field microscopy with an incident circular polarized laser is sensitive to the plasmon dephasing and unambiguously records the change in damping induced upon molecule adsorption. These results offer new perspectives in molecule sensing and materials tunability for light polarization conversion at sub-microscopic scale.

17:50 : Invited talk**Metasurface enabled ultrafast polarization and on-chip light shaping****Amit Agrawal***National Institute of Standards and Technology (USA)*

We demonstrate the versatility of dielectric metasurfaces to (i) shape the temporal evolution of ultrafast optical pulses, and (ii) discuss their applications towards creating integrated photonic interfaces with quantum systems.

18:10 : Invited talk**Low-Loss On-Chip Surface Grating Couplers Engineered Using Subwavelength-Structured Metamaterials****Daniel Benedikovic¹, Xavier Le Roux¹, Carlos Alonso-Ramos¹, Cécilia Dupré², Bertrand Szlag², Pavel Cheben³, Daivid Fowler², Sylvain Guerber², Eric Cassan¹, Delphine Marris-Morini¹, Frédéric Boeuf⁴, Laurent Vivien¹**¹Université Paris-Saclay (France), ²University Grenoble Alpes (France), ³National Research Council Canada

(Canada), ⁴STMicroelectronics (France)

The presence of low-loss optical interfaces is arguably one of the key factors to succeed for silicon photonics. Here, we report on our latest advances in subwavelength-structured and metamaterial-engineered fiber-chip grating couplers with a L-shaped waveguide profiles. Grating couplers are made in silicon-on-insulator platform and facilitate robust sub-decibel coupling of light with device features lying well in the range of available fabrication technologies.

18:30 : Invited talk

Enhanced forward scattering from magnetic scatterers: design and synthesis of resonant Huygens' sources

Philippe Barois, Ashod Aradian, Alexandre Baron, Maria Letizia De Marco, Romain Dezert, Glenna Drisko, Etienne Duguet, Rajam Elancheliyan, Tom Jatteau, Laurent Lermusiaux, Olivier Mondain-Monval, Virginie Ponsinet, Serge Ravaine, Philippe Richetti, Mona Treguer-Delapierre
Université de Bordeaux (France)

We present optical studies of nano-resonators (meta-atoms) designed for a specific control of their angular scattering in visible light. Directional scattering is obtained by adjusting the series of odd and even scattering modes. The architecture of the meta-atoms is designed to enhance the magnetic dipolar scattering which appears as the major contribution to the odd modes. Several types of meta-atoms are investigated. We show that the bottom-up approach enables multiple synthetic strategies and we compare their optical efficiencies.

18:50 : Photon bunching and antibunching of two quantum emitters near a coated sphere

Tiago J. Arruda¹, Romain Bachelard², John Weiner³, Sebastian Slama⁴, Philippe W. Courteille³

¹Federal University of Alfnas (Brazil), ²Federal University of São Carlos (Brazil), ³University of São Paulo (Brazil), ⁴University of Tübingen (Germany)

We theoretically study the cooperative decay rate and the dipole-dipole interaction associated with two dipole emitters close to a coated nanosphere. Considering a plasmonic nanoshell containing a gain medium inside the core, the second-order correlation function for the two emitters shows the possibility of tuning the photon emission, selecting either photon bunching or antibunching as a function of the polarization and position of the sphere. This result opens vistas to applications involving tunable single-photon sources in engineered artificial media.

17:30 - 18:50 — Allan Boardman Room

Session 1A23

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

17:30 : Invited talk

Plasmonics of thermoresponsive nanocomposite metasurfaces

Michele Magnozzi¹, Yannic Brasse², Tobias A. F. Koenig², Francesco Bisio³, Eva Bittrich², Andreas Fery², Maurizio Canepa¹

¹Università di Genova (Italy), ²Leibniz-Institut für Polymerforschung Dresden e.V. (Germany), ³CNR-SPIN (Italy)

Gold/PNIPAM core-shell nanoparticles (NPs) possess proven photothermal and sensing capabilities. In this contribution we report in situ spectroscopic ellipsometry (SE) measurements to determine the complex, temperature dependent properties of ordered lattices of such NPs. The approach proposed here is instrumental to the analysis and design of functional hybrid metasurfaces with plasmonic functionalities, including particle-to-film coupled systems.

17:50 : Invited talk

Ultrafast Spectroscopy and Imaging of Quantum Optical and Plasmonic Processes**Gary Wiederrecht***Argonne National Laboratory (USA)*

There is a strong need for the ability to spatially resolve photoinduced processes in nanoscale structures, and to temporally resolve the evolution of energy flow in these structures. These materials can be particularly challenging for imaging due to issues with scattering and small sample volumes. Novel approaches to the time-resolved spectroscopy and imaging of quantum optical and plasmonic processes are described here.

18:10 : Invited talk**Temporal Plasmonics of metallic nanoparticle dimers: The Fano and Rabi regimes****Oscar Rodrigo Avalos-Ovando¹, Lucas V. Besteiro², Zhiming Wang³, Alexander O. Govorov¹***¹Ohio University (USA), ²Universidad de Vigo (Spain), ³University of Electronic Science and Technology of China (China)*

We study signature temporal responses of the Fano and Rabi regimes in dimers of plasmonic nanoparticles. In their optical spectra, the Fano system possesses a peculiar line-shape, while the Rabi system exhibits a characteristic splitting. We find a difference in temporal dynamics as well: Fano systems show at most one temporal beat after a pulsed excitation, whereas Rabi systems have a significant number of beats. Remarkably, both regimes show coherent time dynamics with non-trivial and characteristic relaxation behaviors.

18:30 : Invited talk**Photon Down-Conversion via Nonadiabatic Dynamics in Molecular Polaritons****Juan B. Perez-Sanchez, Joel Yuen-Zhou***University of California San Diego (USA)*

We theoretically studied the photoisomerization of thiacyanine iodide molecule embedded in an optical microcavity. Our model consisted of the reaction coordinate and two electronic states coupled to a single cavity mode. We showed that an electronic excitation at cis configuration is followed by the generation of two photons upon isomerization. Our finding suggests a new mechanism to achieve photon down-conversion by exploiting the emergent molecular dynamics in polaritonic architectures.

17:30 - 18:55 — Tatsuo Itho Room**Session 1A24****Bottom-up approaches, new fabrication routes and ENSEMBLE3**

Organized by: Dorota Pawlak and Virginie Ponsinet

Chaired by: Dorota Pawlak and Virginie Ponsinet

17:30 : Keynote talk**Metamaterials that travel faster than light: putting the squeeze on photons****John B. Pendry¹, Paloma Huidobro², Emanuele Galiffi¹***¹Imperial College London (United Kingdom), ²Instituto Superior Tecnico-University of Lisbon (Portugal)*

'Nothing can travel faster than light' is not a correct statement. Many things can and do. Think of a wave breaking at an angle on the sea shore. The point of impact travels along the beach very fast if the angle is a shallow one and can travel infinitely quickly as the angle tends to zero. I shall speak about metamaterials in which the structure moves with a velocity close to or faster than light giving rise to phenomena not seen in static structures. The structures naturally break time reversal invariance giving rise to effects for photons that resemble electrons in a magnetic field. In another realisation the metamaterial grabs hold of the field lines of incident radiation and squeezes then into a tightly formed pulse forming a supercontinuum of intense radiation.

18:00 : Invited talk**Combining top-down and bottom-up techniques to fabricate metamaterials****Kay Dietrich¹, Matthias Zilk², Martin Steglich², Thomas Siefke², Uwe Hübner¹, Andreas Tünnermann³,**

Ernst-Bernhard Kley², Carsten Rockstuhl⁴, Thomas Pertsch²

¹Leibniz Institute of Photonic Technology (Germany), ²Friedrich Schiller University Jena (Germany), ³Fraunhofer Institute for Applied Optics and Precision Engineering (Germany), ⁴Karlsruhe Institute of Technology (Germany)

By combining top-down and bottom-up nanotechnologies, we fabricate an isotropic metamaterial with resonant electric and magnetic response. Our approach is based on realizing a large number of nano-scatterers by fast character projection electron-beam lithography and their subsequent randomized embedding into a liquid matrix, which can later be applied to any other surface and solidified. Our approach unlocks novel opportunities to fabricate nanomaterials with a complex optical response in the bulk but also on top of arbitrarily shaped optical elements.

18:20 : Directed assembly of hybrid colloids for optics

Maeva Lafitte¹, Rajam Elanchelian¹, Romain Dezert¹, Quentin Flamant¹, Oren Regev², Philippe Barois¹, Alexandre Baron¹, Olivier Mondain Monval¹, Virginie Ponsinet¹

¹Université de Bordeaux (France), ²Ben Gurion Univ Negev (Israel)

Colloidal assemblies of gold nanoparticles were synthesized using an emulsion-based formulation route. The synthesis technique involved emulsification followed by controlled ripening of an aqueous suspension of gold nanoparticles in an oil phase. The structural control of the as-synthesized clusters is demonstrated using different techniques. Their optical resonant properties are determined by spectroscopic polarized multi-angle static light scattering. The study evidences strong optical magnetic dipolar resonances and strongly forward scattering patterns, both being tunable by monitoring the cluster inner structure.

18:35 : Invited talk**Dispersion, tunability and active properties of chosen structures of hyperbolic metamaterials**

Janusz Parka, Alessandro Pianelli, Karol Sielezin, Rafał Kowrdziej, Michał Dudek, Marek Olifierczuk
Military University of Technology (Poland)

Numerical simulations of hyperbolic metamaterial properties metal-dielectric type structures were described. Propagation of electromagnetic wave from VIS and NIR range through hyperbolic metamaterial structure have been analyzed. The characteristics describing components of permittivity tensor in these materials were presented. Relations between properties of different type of hyperbolic metamaterial are discussed.

17:30 - 18:50 — Christian Huygens Room**Session 1A25****Symposium III: Advanced passive and active metasurfaces and zero-index optics**

Organized by: Howard Lee, Yu-Jung (Yuri) Lu and Zi Jing Wong

Chaired by: Howard Lee, Yu-Jung (Yuri) Lu and Zi Jing Wong

17:30 : Invited talk**Enhanced Hot Carrier Effects Using Ultra-Thin Metal films, Alloys, And Index Near-Zero Substrates**

Jeremy Munday

University of California, Davis (USA)

Bulk metals are good reflectors of light, however, ultra-thin films and nanostructures composed of these same materials can create highly absorptive systems. Here we show that ~100% absorption is achievable provided that the ultra-thin metals are deposited on an index near zero substrate. We further demonstrate that this absorption can be used to generate photocurrent through the transfer of energy from the photons to the free electrons in the metal, enabling new designs for efficiency hot carrier optoelectronic devices.

17:50 : Invited talk**Nonlinear Doping of Epsilon-Near-Zero Media**

Ehsan Nahvi¹, Inigo Liberal², Nader Engheta¹

¹University of Pennsylvania (USA), ²Public University of Navarra (Spain)

We theoretically explore the possibility of obtaining enhanced magnetic nonlinearity by doping a linear epsilon near-zero (ENZ) host with a nonlinear dielectric inclusion. As an application of this concept, we exploit the enhanced nonlinearity in the context of a nonlinear absorber where we deploy a nonlinear doped ENZ slab as the spacer layer in a Salisbury screen. In addition to the enhanced nonlinearity, such absorbers are shown to exhibit highly tunable characteristics, such as bandwidth control.

18:10 : Invited talk

A new approach to meter-scale and durable all-dielectric meta-optics: gains and challenges

Eyal Feigenbaum, J. H. Yoo, N. J. Ray, H. T. Nguyen, M. A. Johnson, S. Baxamusa, S. Elhadj
Lawrence Livermore National Laboratory (USA)

We present an alternative approach to dielectric meta-surfaces based on resonant elements which has far less limitations on scalability and durability. The process is based on laser raster-scan of a thin metal film on a glass, followed by dry-etching and removal of the metal mask. Since the air-glass volumetric ratio mixing approach is limited by the depth of the layer, we have developed approaches to "boost" the attainable phase response, to be discussed here.

18:30 : Invited talk

Metaphotonic Computational Image Sensors

Arka Majumdar

University of Washington (USA)

We demonstrated full color imaging using metasurface and computational imaging techniques.

17:30 - 18:40 — Augustin Fresnel Room

Session 1A26

Symposium IV: Chirality, magnetism, and magnetoelectricity: Separate phenomena and joint effects in metamaterial structures

Organized by: Eugene Kamenetskii

Chaired by: Eugene Kamenetskii

17:30 : Invited talk

Chiral Flat Optics: from chiral photonic crystals to spin-selective metasurfaces

Behrooz Semnani, Jeremy Flannery, Rubayet Al Maruf, Michal Bajcsy

University of Waterloo (Canada)

This work presents the design, fabrication and characterization of a chiral photonic crystal structure which exhibits extreme optical activity around the high-symmetry points in k-space. We also extend the design to chiral focusing metasurfaces with unique functionalities. The underlying physics is guided-mode-resonance combined via weakly completed resonators to introduce both a desired phase profile and reflectivity in a monolithic dielectric structure.

17:50 : Invited talk

Shaping Optical Chirality on the Nanoscale with Photonic Molecules from High Refractive Index Building Blocks

Bjoern Reinhard

Boston University (USA)

Nanostructures, including nanowires, nanodisks and nanoparticles from high refractive index materials can support both magnetic and electric resonances. Combination of these basic building blocks into oligomers or clusters, so called photonic molecules, facilitates an engineering of electromagnetic fields with unique properties. In this presentation, we will discuss design criteria for generating enhanced optical chirality in high refractive index photonic molecules assembled from structurally well-defined silicon nanodisks. Experimental evidence for switchable chiroptical hot spot formation is presented.

18:10 : Keynote talk**Optical and photochemical properties of chiral plasmonic nanostructures****Alexander Govorov¹, Lucas V. Besteiro²**¹Ohio University (USA), ²UESTC and INRS (China)

Chiral photochemical reactions at the molecular level have proven to be a challenging task since chiral molecule species possess tiny chiroptical signals. In contrast, plasmonic nanocrystals offer very strong circular dichroism. We propose taking advantage of this property, introducing a novel mechanism driving surface photochemistry in a chiral plasmonic nanocrystal. This mechanism is based on the generation of hot electrons and leads to plasmon-assisted chiral growth.

17:30 - 18:55 — Ibn Al-Haytham Room**Session 1A27****Symposium V: Phononics and acoustic metamaterials**

Organized by: Jensen Li and Guoliang Huang

Chaired by: Jensen Li and Guoliang Huang

17:30 : Invited talk**Diffraction-free propagation of Gaussian sound beam through layered water-steel structure****Arkadii Krokhn¹, Yurii Zubov¹, Bahram Djafari-Rouhani², Yuqi Jin¹, Mathew Sofiel¹, Ezekiel Walker³, Arup Neogi¹**¹University of North Texas (USA), ²Institut d'Electronique, de Microelectronique et Nanotechnologie (France), ³Echonovus Inc. (USA)

A finite-width acoustic beam propagating in a homogeneous medium spreads with distance. The spreading occurs due to the Fourier components which make nonzero angles with the direction of propagation. The rate of spreading is defined by the diffraction coefficient. We consider propagation of Gaussian sound beam through a periodic steel-water layered structure and demonstrate that for certain frequencies the diffraction coefficient vanishes. In the experiment the nonspreading propagation of sound was observed at distances about 1 m.

17:50 : Invited talk**Double negativity in bubble metamaterials****Maxime Lanoy¹, Geoffroy Lerosey¹, Fabrice Lemoult¹, Valentin Leroy², Arnaud Tourin¹, John Page³**¹Institut Langevin, ESPCI Paris, Université PSL (France), ²Université de Paris (France), ³University of Manitoba (Canada)

I will show how double negativity can be achieved in a metamaterial populated solely with monopolar subwavelength resonators, such as air bubbles in water. A disordered set of identical bubbles in water is known to exhibit a gap above their individual resonance in the frequency range of which only the effective compressibility is negative. But introducing pair-wise spatial correlations leads to the formation of a transparency window, which is associated with negative values of both compressibility and density.

18:10 : Higher-order topological sound transport in synthetic dimensions**Hui Chen¹, Hongkuan Zhang², Qian Wu¹, Yu Huang², Huy Nguyen¹, Emil Prodan³, Xiaoming Zhou², Guoliang Huang¹**¹University of Missouri (USA), ²Beijing Institute of Technology (China), ³Yeshiva University (USA)

We formulate the theoretical principles and for the first time manufacture acoustic crystals composed of arrays of acoustic cavities strongly coupled through modulated channels to evidence two-dimensional (2D) dynamic topological pumping. We delineate the generated four-dimensional (4D) quantum Hall effects by calculating second Chern numbers and physically demonstrate robustness against the imperfections. Synthetic dimensions could open up a new platform to explore any continuous orbit in higher-order topological matter in dimensions four and higher.

18:25 : Active Metamaterials with Odd Micropolar Elasticity**Yangyang Chen¹, Xiaopeng Li¹, Colin Scheibner², Vincenzo Vitelli², Guoliang Huang¹**¹University of Missouri (USA), ²The University of Chicago (USA)

Materials made from active, living, or robotic components can display emergent properties arising from local sensing and computation. Here, we realize a freestanding active metabeam with piezoelectric elements and electronic feed-forward control that gives rise to an odd micropolar elasticity absent in energy-conserving media. Our continuum approach, built on symmetries and conservation laws, could be exploited to design others systems such as synthetic biofilaments and membranes with feed-forward control loops.

18:40 : Microtwist homogenization of 3D Pyrochlore lattices on Mechanical Polarization**Rongyu Xia¹, Hui Chen¹, Zheng Li², Guoliang Huang¹**¹University of Missouri (USA), ²Peking University (China)

We develop a 3D microtwist effective theory of Pyrochlore lattices to capture the Parity (P) asymmetric zero modes by which polarization emerges or fades on a macroscopic scale. By mapping three periodic zero modes to macroscopic degrees of freedom, the 3D microtwist theory ends up being a kinematically enriched theory. The 3D microtwist elasticity is formulated by using two-scale asymptotic approach and its constitutive and balance equations are derived for a fairly generic isostatic lattice: the 3D Pyrochlore lattice.

17:30 - 18:45 — Gaston Floquet Room**Session 1A28****Metamaterials and metasurfaces****17:30 : Ultra-wideband waveguide embedded graphene-based terahertz absorber****James Campion¹, Nikolaos Xenidis¹, Roman Ivanov², Joachim Oberhammer¹, Irina Hussainova², Dmitry Lioubtchenko³**¹KTH Royal Institute of Technology (Sweden), ²Tallinn University of Technology (Estonia), ³KTH Royal Institute of Technology (Sweden) and Institute of High-Pressure Physics, PAS (Poland)

A novel type of absorber material is developed for the frequency range of 67-500 GHz. The absorber is based on graphene augmented inorganic nanofibers deposited inside a metallic waveguide cassette, allowing them to be utilised in standard waveguide systems. The material's microstructures result in a low level of reflectance (< -15 dB) and good absorbance (*textgreater* 20 dB) from 110-500 GHz, making them highly suited for wideband terahertz applications.

17:45 : Multipolar analysis of random all-dielectric nanoresonator arrays**Maria Bancerek, Krzysztof M. Czajkowski, Olga Kochanowska, Tomasz J. Antosiewicz**

University of Warsaw (Poland)

The optical properties of bottom-up amorphous arrays with stochastic distribution of high-index dielectric antennas stem from an interplay of the single-particle response, mutual coupling and substrate-mediated effects. We develop a theoretical framework that incorporates electric and magnetic dipolar interactions to prove that interparticle coupling even in random arrays is important. Our results show how to utilize intra-array coupling to maximize array sensitivity for refractometric sensing purposes.

18:00 : Ge₂Sb₂Te₅-based, ultrathin, all-dielectric tunable mid-wavelength infrared perfect absorber**Roy Avrahamy, Amiel Avraham Ishaaya, Mark Auslender**

Ben-Gurion University of the Negev (Israel)

GST-225 (Ge₂Sb₂Te₅) phase change material was recently experimentally reported to exhibit measurable photoconductivity, a well-defined bandgap, and reconfigurable continuous partial crystallization. Sequentially, here we propose an ultrathin, all-dielectric, metamaterial-design based on an asymmetrical optical micro-/nanocavity, enclosing a 10nm thick GST-225 photoactive layer, inversely optimized for perfect tunable absorption in the mid-wavelength infrared. The perfect absorption, solely in GST-225, can be spectrally-tuned actively (thermally/electrically/optically) by varying the crystallinity, and geometrically using the design parameters, which is highly application beneficial.

18:15 : Linear-to-circular polarization conversion using time-dependent metamaterials**Victor Pacheco Peña¹, Nader Engheta²**¹Newcastle University (United Kingdom), ²University of Pennsylvania (USA)

In this communication we explore a mechanism to achieve an arbitrary conversion of the polarization of electromagnetic waves by using time-dependent metamaterials for real-time polarization rotation and frequency conversion.

18:30 : Symbolic regression in nano-optics: characterization of dispersive materials as a case study**Claudio Canales¹, Alexandre Vial², Demetrio Macias²**¹Universidad de Santiago de Chile (Chile), ²University of Technology of Troyes (France)

In this contribution we propose a Symbolic Regression (SR) scheme, based on Genetic Programming (GP), to retrieve the closed expression that represents the dispersion model of a given material. As starting point, we consider an isotropic, homogeneous, dispersive dielectric material to illustrate the operational principles of our approach. Then, we discuss its possibilities and limitations when used to solve this kind of inverse problem.

17:30 - 19:00 — Lawrence Bragg Room**Session 1A29****Plasmonics: Fundamentals and Applications**

Organized by: Hong Wei

Chaired by: Hong Wei

17:30 : Invited talk**Plasmon-Exciton Coupling: Light-forbidden Transitions and Quasichiral Interactions****Antonio I. Fernandez-Dominguez***Universidad Autonoma de Madrid (Spain)*

We present two plasmon-exciton coupling phenomena emerging due to the confined nature of surface plasmon (SP) resonances in nanocavities. First, we will investigate the impact of light-forbidden transitions have in the population dynamics and far-field spectrum of nanoparticle-on-a-mirror SPs and three-level quantum emitters (QEs). Second, we will present a combined classical and quantum electrodynamics description of the interactions between two circularly-polarized QEs held above a SP waveguide. We will establish the conditions for non-reciprocal, chiral, coupling between them.

17:50 : Invited talk**Observing strong coupling in individual plasmonic cavities****Gilad Haran***Weizmann Institute of Science (Israel)*

We utilize individual plasmonic bowties as cavities to couple to a small number of quantum emitters. Devices with one to several semiconductor quantum dots demonstrate vacuum Rabi splitting in light scattering spectra and in electron energy loss spectra. Data analysis shows that our systems are within or at the onset of the strong coupling regime. Photoluminescence studies demonstrate antibunching from single quantum dots within plasmonic cavities, paving the way to cavity QED studies at room temperature.

18:10 : Invited talk**Quantitative characterization of second order nonlinear light conversion from inorganic and organic nano(micro)-structures****Ning Liu¹, Matthew Gleeson¹, Zhe Li¹, Sarah Guerin¹, Syed A. M. Tofail¹, Pritam Khan¹, Hongxing Xu², Christophe Silien¹**¹University of Limerick (Ireland), ²Wuhan University (China)

Second harmonic generation and sum frequency generation are the nonlinear optical processes of doubling or summing the frequency of input light by passing it through non-centrosymmetric crystalline materials.

Quantitative modelling of their nonlinear processes is of great importance for optimizing nano(micro)crystal based nonlinear photonic devices for applications in frequency conversion, multiplexed signal transmission and noninvasive sensing. Here we demonstrate the quantitative analyses of optical nonlinear conversion in inorganic and organic nano(micro)structures with known and unknown second order susceptibility tensor.

18:30 : THz plasmons in AlGaIn/GaN grating gate structures at 4K and 300K

Pavlo Sai, Maciej Sakowicz, Kamil Stelmaszczyk, Dmytro B. But, Mateusz Słowikowski, Maciej Filipiak, Maksym Dub, Pawel Prystawko, Grzegorz Cywinski, Sergey Romyantsev, Wojciech Knap

CENTERA Laboratories (Poland)

Terahertz plasmon resonances were studied at 4.2K and 300K in GaN-based grating gate structures using two THz spectroscopy techniques: Fourier-Transform Infrared Spectroscopy and Time Domain Spectroscopy. Gratings of different periods were coupled to the two-dimensional electron gas in AlGaIn/GaN in order to investigate the dispersion law of 2D-plasmons. The plasmon frequency was tuned by the gate voltage both, at 4.2K and 300K. Observation of the tunable plasmons at room temperatures opens the way for high temperature THz plasmonic devices.

18:45 : Deciphering Mode-Coupling Mechanism in Extraordinary Optical Transmission via Signature Fano Resonance for Active Tuning

Meng-Ju Yu, Jimmy Xu

Brown University (USA)

We report on an investigation that aims at deciphering the mode-coupling mechanisms in extraordinary optical transmission (EOT). The EOT phenomenon, being extraordinary, is also intriguing in the interplays between its modes or mode-couplings. Understanding the coupling mechanism would pave the way to the next advances such as active- or on-command tunable EOT. We conduct computational experiments to decipher the coupling mechanism through the Fano resonance signature.

17:30 - 18:50 — Rene Descartes Room

Session 1A30

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

17:30 : Invited talk

Sparse Array as Metamaterial for Higher-Order Modes Suppression in an Accelerator Cavity

Ning Zhou¹, Terry Smith², Geoff Waldschmidt², Alireza Nassiri², Thomas T. Y. Wong¹

¹ *Illinois Institute of Technology (USA)*, ² *Argonne National Laboratory (USA)*

A metamaterial formed by sparsely populated array of metallic rods exhibiting photonic bandgap (PBG) properties is employed to fill a cavity resonator for future particle accelerator applications. Beginning with an array on a triangular lattice, the optimization process leads to a star-shape array. The cavity-waveguide assembly was fabricated with copper and cold tested for resonance characteristics. A return loss of over 20 dB at the designed resonance frequency of 11.41 GHz was measured.

17:50 : Invited talk

Acoustic Graphene Plasmons under a DC bias

Michael Sammon, Tony Low

University of Minnesota (USA)

It has been shown that in a graphene/insulator/metal structure, the plasmons acquire a linear dispersion whose sound velocity is very near the Fermi velocity in graphene. I show that a DC current causes a redshift of the acoustic plasmons travelling upstream of the direction in which electrons flow. As the redshift of the upstream branch causes acoustic plasmon to approach the particle-hole continuum, the spectral weight of this plasmon branch is substantially reduced, resulting in a near unidirectional plasmon.

18:10 : Invited talk

Nanostructured Complex Metal Oxides for Infrared Plasmonics**Nicholas Charipar, Raymond Auyeung, Heungsoo Kim, Kristin Charipar, Alberto Piqué***U.S. Naval Research Laboratory (USA)*

Complex metal oxides, such as indium tin oxide (ITO) and vanadium dioxide (VO₂), are of interest for plasmonics because ITO is an optically transparent electrical conductor while VO₂ exhibits a metal-insulator transition. The growth of ITO and VO₂ thin films via pulsed laser deposition is discussed. Laser processing techniques are investigated as a means to produce ordered nanostructures in these thin films. The infrared properties of both the as-grown thin films and resulting nanostructured complex metal oxides are characterized.

18:30 : Invited talk**field-Ready Quantum Technologies based on Nanowire Sources****Khaled Mnaymneh¹, Robin L Williams¹, David B Northeast¹, Jeongwan Jin¹, Sofiane Haffouz¹, Patrick Laferrière², Edith Yeung², Lambert Giner², Henri Morin², Jean Lapointe³, Geoffrey C Aers³, Philip J Poole³, Dan Dalacu³**¹*Emerging Technologies, National Research Council Canada (Canada)*, ²*University of Ottawa (Canada)*,³*National Research Council Canada (Canada)*

Quantum technologies based on semiconductor quantum dots placed in photonic nanowires are discussed. Epitaxial growth strategies of the embedded dots targeting field-appropriate wavelengths are presented. The tapered shape of the nanowires allows for direct and evanescent coupling to bulk and integrated optics, respectively. This enables a high degree of photonic integration aimed at deploying plug-and-play quantum systems to the field for applications in secure communications, sensing and metrology.

18:00 - 19:00 — Tutorial Room**Session 1A31****Conference Tutorials I**

Organized by: Ishwar Aggarwal

18:00 : Tutorial**Metasurface Flat Optics: from components to mass manufacturing to systems****Federico Capasso***Harvard University (USA)*

Flat optics based on metasurfaces has emerged in recent years as a promising alternative to refractive and Fresnel optics in many applications, due to the smaller footprint, mass-manufacturing using the same technology of semiconductor chips, easier control of aberrations and multifunctionality. I will cover recent advances in components and show how they have led to breakthroughs in cameras and other systems such as ultra compact spectrometers.

Wednesday 21st July, 2021

09:00 - 11:00 — Victor Veselago Room

Session 2A1

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

09:00 : **Invited talk**

Nanophotonics and femtosecond magnetism in all-dielectric metasurfaces

Alexander Chernov¹, Mikhail Kozhaev¹, Daria Ignatyeva¹, Andrey Voronov¹, Dolendra Karki², Miguel Levy², Vladimir Belotelov¹

¹Russian Quantum Center (Russia), ²Michigan Technological University (USA)

Light manipulation in magnetic nanostructured materials attracts much attention in the context of data processing, spintronic and light modulation applications. In this work we demonstrate a subwavelength light localization within the magnetic dielectric metasurface made of bismuth-substituted iron garnet leading to light intensity modulation and an efficient magnon excitation. The advanced light control and selective spin manipulation are achieved due to appearance of various types of excited modes in the nanostructured surfaces.

09:20 : **Invited talk**

Free-carriers nonlinearities in semiconductor plasmonics

Federico De Luca¹, Michele Ortolani², Cristian Ciraci¹

¹Istituto Italiano di Tecnologia (IIT) (Italy), ²Sapienza University of Rome (Italy)

We study free-carriers nonlinearities in highly doped semiconductors. We develop a theoretical model based on the hydrodynamic description of free-electrons expanding nonlinear terms up to the third-order. Because of small carrier densities in semiconductors compared to noble metals, hydrodynamic effects result strongly amplified. We show that contrarily to noble metals, in fact, free-electron nonlinearities in doped semiconductors can be several orders of magnitude larger than crystalline lattice nonlinearities.

09:40 : **Invited talk**

Surface waves from flexural and compressional resonances of beams

Agnes Maurel¹, Kim Pham², Sébastien Guenneau³, Jean-Jacques Marigo⁴

¹Institut Langevin/ESPCI (France), ²Institut Polytechnique de Paris (France), ³Imperial College (United Kingdom), ⁴Ecole Polytechnique (France)

We present a model describing the propagation of elastic waves in a soil substrate supporting an array of cylindrical beams experiencing flexural and compressional resonances. The resulting surface waves are of two types: in the sagittal plane, hybridized Rayleigh waves which can propagate except within bandgaps resulting from a complex interplay between flexural and compressional resonances. anti-plane shear wave, decoupled from the hybridized Rayleigh wave, which is the elastic analogue of electromagnetic spoof plasmon polaritons.

10:00 : **Invited talk**

A computational technique for the scattering problem of inhomogeneous metasurfaces

Bo O. Zhu

Nanjing University (China)

Metasurfaces have attracted much attentions recently because of the enriched electromagnetic wave control abilities. So far, the full wave analysis of metasurfaces, especially that of inhomogeneous ones, is not well supported by the popular commercial softwares. In this paper, a computational technique is proposed for the scattering problems of metasurfaces. Compared with the popular numerical methods, the finite element method, the proposed approach is easy to implement since it does not need meshing and basis functions.

10:20 : Invited talk

Optical Chirality Generation and Amplification in Biomolecule-Assembled Plasmonic Nanostructures
Dangyuan Lei

City University of Hong Kong (Hong Kong)

I will discuss the generation and amplification of optical chirality in achiral metal nanostructures assembled with chiral biomolecules. The first part of this talk focuses on how an individual plasmonic nanostructure interacts with chiral biomolecules, enabling switching the molecular chirality and probing the molecular conformation evolution. The second part presents how to assemble one-dimensional plasmonic nanoparticle chains with chiral biomolecules to induce plasmonic circular dichroism response in the NIR-UV region and their assembly configuration dependence.

10:40 : Invited talk

Characterization of mid-IR photonic crystal slabs using angle-resolved FT-IR spectroscopy

Takashi Kuroda¹, Siti Chalimah¹, Yuanzhao Yao¹, Naoki Ikeda¹, Yoshimasa Sugimoto¹, Rei Hashimoto², Tsutomu Kakuno², Kei Kaneko², Shinji Saito², Kazuaki Sakoda¹

¹National Institute for Materials Science (Japan), ²Toshiba Corp. (Japan)

Fourier transform (FT) spectroscopy is a versatile technique to study the infrared (IR) optical response of solid-, liquid-, and gas-phase samples. Here we design and construct a high-precision angle-resolved reflectance setup compatible with a commercial FT-IR spectrometer. As a demonstration of the capability of the reflectance setup, we measure the angle-dependent mid-infrared spectra of two-dimensional photonic crystal slabs, and determine the in-plane photonic band dispersion in the vicinity of gamma point in momentum space.

09:00 - 11:00 — Allan Boardman Room

Session 2A2

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

09:00 : Keynote talk

Dielectric metasurfaces for flat optics: wavefront engineering and future applications

Junsuk Rho

Pohang University of Science and Technology (POSTECH) (Korea)

Miniaturization is a main stream in modern technology, but reduction of conventional optical components accompanies performance degradation that limits the minimum feature size of optical devices. Metasurfaces that consist of ultrathin subwavelength antenna arrays can be a promising solution because metasurfaces provide an effective way of wavefront engineering without constraints on the device size. Electromagnetic responses of individual building blocks are determined by its geometric configurations, and many kinds of antennas have been explored to clarify the capability of metasurfaces, thereby, it has been verified that dielectric antennas can control amplitude, phase, and even both of them simultaneously.

09:30 : Invited talk

Microscopic Study on Circular Dichroism Localized in Materials to Analyze Hierarchical Chirality in Various Scales

Tetsuya Narushima

Institute for Molecular Science (Japan)

Circular dichroism (CD) has been used to analyze chirality in molecules. These years, it has been reported that spatially extended chiral systems, such as crystals, molecular assemblies and nanostructures, also show CD activity. In order to explore properties of the CD activity and its physical origin, we developed a CD microscope and visualized spatial CD distribution observed in the chiral systems of metallic nanostructures,

biological cells, and so forth.

09:50 : Invited talk

Gap-SPPs of Closely Coupled Nanowire Dimers Visualized by SERS Imaging

Sang-Min Park, Zee Hwan Kim

Seoul National University (Korea)

We report a visualization of gap-SPPs of a AgNW dimer. A dimer loaded with a monolayer of molecules is locally excited to launch the SPPs, and the SERS maps of the molecules are acquired to visualize gap-SPPs. The SPP maps reveal that the dimers with a few nm of gap can propagate up to $\sim 8 \mu\text{m}$. They also show oscillating components with periods of 400 \sim 800 nm, arising from the beating between a monopole-monopole and a dipole-dipole gap-SPPs.

10:10 : Invited talk

Luminescence Manipulation from Upconversion Nanorods via Plasmonic Nanogratings and Shear-Induced Alignment

Huilin He, Anshi Chu, Zhen Yin, Jianxun Liu, Ke Li, Jiawei Wang, Dan Luo, Yanjun Liu

Southern University of Science and Technology (China)

Greatly enhanced upconversion luminescence was demonstrated by integrating the core-shell upconversion nanorods with the Ag nanogratings. We also demonstrated a facile polypropylene-aided shear-driven method to obtain large-scale orientationally ordered upconversion nanorods, showing a liquid crystalline nematic phase.

10:30 : Aluminum fractal nanoantennas based on Cayley trees

Thomas Simon¹, Xiaoyan Li², Dmitry Kholpin¹, Jérôme Martin¹, Mathieu Kociak², Davy Gerard¹

¹Universite de Technologie de Troyes (France), ²Université Paris Sud (France)

We propose and demonstrate optical antennas based on the Cayley tree branching geometry. Using electron energy loss spectroscopy, we unveil the complex mode structure associated with these antennas. A broadband optical response, spanning from the UV to the mid-IR, is evidenced.

10:45 : Strong Coupling in Metallo-dielectric Hybrid Metasurfaces

Ajith Padyana Ravishankar, Srinivasan Anand

KTH Royal Institute of Technology (Sweden)

The presented work involves designing a hybrid metasurfaces by incorporating merits of two different types of resonator system (metal and dielectric) and investigating novel optical features emerging from the metasurface. The metasurface design consists of a high-index resonator array on top of thick metal film with a spacer layer. Simulation studies show that a strong coupling can be achieved between the anapole mode in the disk and a surface plasmon polariton (SPP) mode at metal dielectric interface.

09:00 - 10:55 — Tatsuo Itho Room

Session 2A3

Bottom-up approaches, new fabrication routes and ENSEMBLE3

Organized by: Dorota Pawlak and Virginie Ponsinet

Chaired by: Dorota Pawlak and Virginie Ponsinet

09:00 : Reversible strain-tuning of quantum optical emission in WSe₂ monolayers

Javier Martin-Sanchez¹, Oliver Iff², Davide Tedeschi³, Magdalena Moczala-Dusanowska², Sefaattin Tongay⁴, Kentaro Yumigeta⁴, Javier Taboada-Gutiérrez¹, Matteo Savaresi³, Armando Rastelli⁵, Pablo Alonso-Gonzalez¹, Sven Höfling², Rinaldo Trotta³, Christian Schneider²

¹University of Oviedo (Spain), ²Universität Würzburg (Germany), ³Sapienza University of Rome (Italy), ⁴Arizona State University (USA), ⁵Johannes Kepler University Linz (Austria)

The future development of ultra-compact two-dimensional (2D) photonic technologies for quantum information

processing relies on our ability to tailor the optical properties of single photon sources in 2D nanomaterials. In this talk, we will present hybrid 2D-piezoelectric devices for the reversible manipulation of the emission energy of quantum emitters in wrinkled WSe₂ semiconductor monolayers. Our results show a record tuning range of about 15 meV while preserving a high single photon purity.

09:15 : Invited talk

Mid-submicrometre pixelation of InGaN micro-LED displays with high integration capabilities for AR-glasses

Jun Hee Choi, Jinjoo Park, Kiho Kong, Joo Hun Han, Jung Hun Park, Nakhyun Kim, Eunsung Lee, Joosung Kim, Dong Chul Shin, Younghwan Park, Jaikwang Shin

Samsung Advanced Institute of Technology (Korea)

InGaN-based blue light-emitting diodes (LEDs), with their high efficiency and brightness, are entering the display industry. However, a significant gap remains between the expectation of highly efficient light sources and their experimental realization into tiny pixels for ultrahigh-density displays for augmented reality (AR). Here, we report using tailored ion implantation (TIIP) to fabricate highly-efficient, electrically-driven pixelated InGaN microLEDs (μ LEDs) at the mid-submicron scale (line/space of 0.5/0.5 μ m). Moreover, we demonstrate high-density TFT and QD C/F integration technologies.

09:35 : Keynote talk

Nanophotonics with Two-Dimensional Materials

Javier Garcia de Abajo

ICFO – Institut de Ciències Fotoniques (Spain)

In this presentation, we will overview the general characteristics of the optical response of these materials, which can be understood in terms of simple theoretical models. We will also cover more sophisticated descriptions, aiming at exploring genuinely quantum-mechanical effects. We will further overview recent advances in the fields of ultrafast optical response and nonlinear optics, as well as the potential application of these materials in light modulation, quantum-optics, and optical sensing.

10:05 : Keynote talk

3D Laser Nanoprinting of 3D Metamaterials

Martin Wegener

Karlsruhe Institute of Technology (KIT) (Germany)

We review our recent progress concerning 3D laser nanoprinting of 3D metamaterials. In the context of this special session, we emphasize advances in 3D additive manufacturing with respect to printing speed.

10:35 : Invited talk

Self-Organized Metamaterials and Nanoparticles for Nonlinear Optics and Hot Electron Processes

Luke H. Nicholls, Francesco Lotti, Yunlu Jiang, Mazhar E. Nasir, Anatoly V. Zayats

King's College London (United Kingdom)

Metamaterials offer a way to control and utilize light on the nanoscale, which cannot be achieved with conventional materials. However, many metamaterial constructions require time consuming fabrication methods that result in small active area arrays. The metallic nanorod metamaterial, with its self-organised fabrication method, allows 2-inch wafer size metamaterial arrays to be constructed both quickly and cheaply. In this talk the applications of this metamaterial will be discussed from chemical sensing to nonlinear optics and intensity dependent filters.

09:00 - 10:00 — Christian Huygens Room

Session 2A4

Symposium III: Advanced passive and active metasurfaces and zero-index optics

Organized by: Howard Lee, Yu-Jung (Yuri) Lu and Zi Jing Wong

Chaired by: Howard Lee, Yu-Jung (Yuri) Lu and Zi Jing Wong

09:00 : Invited talk**Electrically Tunable Metasurface for Complex Amplitude Modulation****Sangjun Han¹, Seyoon Kim², Shinho Kim¹, Tony Low³, Victor Brar², Min Seok Jang¹**¹*Korea Advanced Institute of Science and Technology (KAIST) (Korea)*, ²*University of Wisconsin-Madison (USA)*, ³*University of Minnesota (USA)*

Tunable plasmonic modes offered by graphene provide new opportunities to create electro-optically active devices with novel characteristics that have thus far been impossible to be realized by using conventional media. In this talk, we introduce dynamic complex amplitude modulation with graphene-based metasurfaces.

09:20 : Invited talk**Plasmon-assisted multipolar terahertz absorption spectroscopy in graphene****Andrea Marini¹, Alessandro Ciattoni², Claudio Conti³**¹*University of L'Aquila (Italy)*, ²*CNR-SPIN (Italy)*, ³*Consiglio Nazionale delle Ricerche (Italy)*

We explore plasmon-enhanced absorption spectroscopy showing that multipolar rotational transitions of molecules in proximity of localized graphene structures can be accessed thanks to terahertz plasmons. In particular we consider H₂⁺, demonstrating that graphene micro-rings provide a giant field localization enabling the enhancement of the absorption cross-section by 8 orders of magnitude.

09:40 : Invited talk**Metaphotonics meet fibers: a novel pathway towards boosting in-coupling efficiencies and single-fiber optical trapping****Markus A. Schmidt¹, Henrik Schneidewind¹, Uwe Huebner², Matthias Zeisberger³, Malte Plidschun³, Jisoo Kim³, Oleh Yermakov⁴, Yuri Kivshar⁴, Andrey Bogdanov⁴, Haoran Ren⁵, Stefan A. Maier⁶**¹*Leibniz Institute of Photonic Technology (Germany)*, ²*Leibniz Institute of Photonic Technology (Germany)*, ³*Leibniz Institute of Photonics Technology (Germany)*, ⁴*ITMO University (Russia)*, ⁵*Macquarie University (Australia)*, ⁶*Ludwig-Maximilian University of Munich (Germany)*

Here we present that interfacing optical fiber with nanostructures defines a novel class of fiber-integrated devices - nanostructure-interfaced fiber - allowing to reach new domains of applications for Fiber Optics research. Through 3D nanoprinting and modified electron-beam lithography, we integrate high NA-meta-lenses and dielectric ring gratings onto the end faces of single mode fibers. These devices allows for efficient light incoupling at angles as large as 80° and for trapping polymer microbeads with one single-mode fiber device.

10:00 - 11:00 — Christian Huygens Room**Session 2A5****Non-Hermitian Photonics: From Plasmonics, and Metamaterials, to Topological and Quantum systems**

Organized by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

Chaired by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

10:00 : Invited talk**Nonequilibrium dynamics and thermalization in open quantum many-body systems****Yuto Ashida, Masahito Ueda***The University of Tokyo (Japan)*

I will present two distinct theoretical formalisms to analyze out-of-equilibrium open many-body dynamics with an arbitrary number of quantum jumps. first, I discuss propagation of correlations beyond the Lieb-Robinson bound, which originates from the nonorthogonality of non-Hermitian eigenstates. Second, I show that a generic nonintegrable many-body system subject to continuous observation thermalizes itself at a single-trajectory level. This finding provides a way to efficiently solve a many-body Lindblad master equation.

10:20 : Invited talk**Localization and universality in non-Hermitian many-body systems**

Ryusuke Hamazaki¹, Kohei Kawabata², Naoto Kura², Masahito Ueda²

¹RIKEN (Japan), ²University of Tokyo (Japan)

We show that novel and rich physics concerning localization and universality appears in non-Hermitian quantum many-body systems. As a first topic, we analyze non-Hermitian systems with asymmetric hopping in the presence of interaction and disorder. We demonstrate that a novel real-complex transition occurs upon many-body localization. As a second topic, we discuss universality classes of spectral statistics in non-Hermitian random matrices. We find two new universality classes characterized by transposition symmetry, which is distinct from time-reversal symmetry.

10:40 : Invited talk

Inversely Designed Clustering in Non-Hermitian Disorder for Controlled Wave Delocalization

Sunkyoo Yu, Xianji Piao, Namkyoo Park

Seoul National University (Korea)

Here we explore wave localization in non-Hermitian disordered potentials. By employing the inversely designed optical platform having the target nearest-neighbor correlation, we investigate the effect of clustering on localization. At the specific level of gain or loss, anomalous delocalization of light is observed for highly disordered patterns, which originates from the clustering-induced alteration of the parity-time symmetry breaking condition. The comparison with other delocalization processes in non-Hermitian systems will also be discussed.

09:00 - 11:00 — Augustin Fresnel Room

Session 2A6

Symposium IV: Chirality, magnetism, and magnetoelectricity: Separate phenomena and joint effects in metamaterial structures

Organized by: Eugene Kamenetskii

Chaired by: Eugene Kamenetskii

09:00 : Invited talk

Generation of superconducting vortices by angular momentum of light

Takehito Yokoyama

Tokyo Institute of Technology (Japan)

We investigate a superconducting state irradiated by a laser beam with spin and orbital angular momentum. It is shown that superconducting vortices are generated by the laser beam due to heating effect and transfer of angular momentum of light. Possible experiments to verify our prediction are also discussed.

09:20 : Invited talk

Chirality of Multipolar Lattice Resonances in Plasmonic Crystal Excited by Vortex Beams

Keiji Sasaki, Hiroki Kitajim, Kyosuke Sakai

Hokkaido University (Japan)

We demonstrate that dipolar and quadrupolar lattice resonances in finite-sized, square-lattice arrays of metal nanodisks can be excited by circularly polarized vortex beams carrying spin and orbital angular momenta. The simulation results show that spatial distributions of the lattice resonances in the plasmonic crystals exhibit characteristic patterns with the chirality that conserve the chirality of the incident light and the excited plasmonic fields of individual nanodisks.

09:40 : Invited talk

Optical vortex induced structured materials via two-photon-absorption

Takashige Omatsu

Chiba University (Japan)

We report on that irradiation of picosecond optical vortex pulses with orbital angular momentum allows us to form unique helical or flower-shaped structures via two photon absorption. Such structures reflect not only the

spatial intensity profile, wavefront and polarization of the irradiated optical vortex field but also the nonlinear spatial modulational instability of optical vortex in materials.

10:00 : Invited talk

High Harmonic Generation in Quantum Spin Liquids: Analogy to Graphene, Semiconductors, and Superconductors

Minoru Kanega¹, Tatsuhiko N. Ikeda², Masahiro Sato¹

¹Ibaraki University (Japan), ²University of Tokyo (Japan)

High harmonic generation (HHG) in matter has been extensively studied, and metallic systems have long been its central target in solids. Recently, researchers have begun to extend the targets of HHG to other materials. We theoretically study HHG in magnetic insulators, especially, focusing on quantum spin liquids (QSLs). We show that HHG in QSLs may exhibit their characteristics and the emergence of their even-order harmonics is controlled with a static external field in a class of magnets.

10:20 : Invited talk

Evolution of the Hybridization Processes in Heavy Fermion CeCoIn₅

Zhengxing Wei, Yupeng Liu, Jingbo Qi

University of Electronic Science and Technology of China (China)

We investigate the quasiparticle dynamics in the heavy fermion CeCoIn₅ using ultrafast pump-probe spectroscopy. Our results indicate that this material system undergoes hybridization fluctuations before the establishment of heavy electron coherence, as the temperature decreases from ~ 120 K (T₊) to ~ 55 K (T^{*}). Such observation urges to develop new microscopic theory different from the conventional single impurity Kondo model for understanding the hybridization process.

10:40 : Invited talk

Spin current generation due to Stern-Gerlach-like effects

Mamoru Matsuo

University of Chinese Academy of Sciences (China)

Spin current is a key concept in spintronics. It is known that the spin current is generated in media with the presence of spin dependent potentials, such as a strong spin-orbit coupling and spin-vorticity coupling. In this talk, we focus on the Stern-Gerlach-like spin transport phenomena driven by spin-vorticity coupling in elastic materials as well as by the transverse spin of the surface plasmon polariton.

09:00 - 10:40 — Ibn Al-Haytham Room

Session 2A7

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

09:00 : Invited talk

Plasmonic Probe for Electrochemical Reaction at Metal Nanostructures

Kei Murakoshi

Hokkaido University (Japan)

Single layer graphene can be used to determine the electronic structures of plasmonic nano-electrodes for photoelectrochemical energy conversions. We have revealed the relationship between the photoenergy conversion ability and electrochemical potential of the Fermi level of the plasmonic structure. The present electrochemical Raman measurements were proved to provide detailed understanding on the plasmon-induced charge transfer process for further developments on the ability.

09:20 : Invited talk

Directional emission from colloidal quantum dots using dielectric metasurfaces

Yeonsang Park, Hyochul Kim, Jeong Yub Lee, Woong Ko, Kideok Bae, Kyung-Sang Cho

Samsung Advanced Institute of Technology (Korea)

We present a dielectric metasurface deflector composed of two TiO₂ nanoposts whose design was adapted from the optical Yagi-Uda nanoantenna. The deflector array was formed and integrated with a colloidal quantum-dot resonant cavity using a dry etching method. To demonstrate the deflection performance, we measured the far-field image of the photoluminescence from colloidal quantum dots. We have demonstrated that the quantum dot emission transmitted through deflector arrays was deflected by 20°.

09:40 : Invited talk

Metasurface-Enabled 3-Dimensional Structured-Light Imaging

Seunghoon Han, Jang-Woo You, Byunghoon Na, Jeong Yub Lee, Namseop Kwon, Jaekwan Kim, Hyeonsoo Park, Narae Han, Yongsung Kim, Hyuck Choo

Samsung Electronics (Korea)

We have demonstrated compact structured-light projectors for 3D-depth imaging. The projectors are made by integrating metastructures directly on a DFB LD and a VCSEL array. The metastructures are subwavelength dielectric nanostructures that allow very precise, versatile wavefront manipulation. They serve as a single-element solution that provides precise control of phases and amplitudes, and the nanoscale size allows a wide field-of-view in a smallest form-factor possible.

10:00 : Invited talk

Phase Mapping of the SPP-Coupled Nanoparticles by Angle-Resolved Cathodoluminescence

Takumi Sannomiya¹, Andrea Konecna², Taeko Matsukata¹, Zac Thollar¹, Takayuki Okamoto³, F. Javier Garcia de Abajo², Naoki Yamamoto¹

¹ *Tokyo Institute of Technology (Japan)*, ² *ICFO-Institut de Ciències Fotoniques (Spain)*, ³ *RIKEN (Japan)*

Nanoscale gaps between metals can strongly confine electromagnetic fields that enable efficient electromagnetic energy conversion and coupling to nanophotonic structures. In particular, the gap formed by depositing a metallic particle on a metallic substrate produces coupling of localized particle plasmons to propagating surface plasmon polaritons. Here we demonstrate the experimental visualization of the phase associated with the plasmonic field of metallic particle-surface composites through nanoscopically and spectroscopically resolved cathodoluminescence using a scanning transmission electron microscope.

10:20 : Invited talk

On-chip super-compact photonic quantum CNOT gate

Xifeng Ren¹, Ming Zhang², Lantian Feng¹, Daoxin Dai²

¹ *University of Science and Technology of China (China)*, ² *Centre for Optical and Electromagnetic Research (China)*

We demonstrate a super-compact integrated quantum CNOT gate on a silicon chip by using the idea of symmetry breaking of a 6-channel waveguide superlattice. The present path-encoded quantum CNOT gate is implemented with a footprint of 4.8x4.45 μm² as well as a high process fidelity of ~0.925 and a low excess loss of <0.2 dB. The footprint is shrunk very significantly by ~10,000 times compared to those previous integrated quantum CNOT gates based on dielectric waveguides.

09:00 - 10:55 — Gaston Floquet Room

Session 2A8

Parity-Time and quasi-normal modes in Photonics, Plasmonics, Acoustics

Organized by: Anatole Lupu and Henri Benisty

Chaired by: Anatole Lupu and Henri Benisty

09:00 : Invited talk

Non-conservative optics with dielectric metasurfaces

Andrey A. Sukhorukov

The Australian National University (Australia)

We introduce a general approach for tailored non-conservative transformations of polarization states of light

based on scattering and interference from specially engineered nano-resonators in ultra-thin dielectric metasurfaces. We present theoretical and experimental results demonstrating the fundamental aspects and potential applications of such metasurfaces. These include an optimal monitoring of deviations from a selected polarization, transformation of any input two-photon quantum polarization-entangled state to an arbitrary target state, and discrimination between a set of objects with different polarization characteristics.

09:20 : Invited talk

Revealing the missing dimension at an exceptional point

Renmin Ma

Peking University (China)

The radiation of electromagnetic and mechanical waves depends not only on the intrinsic properties of the emitter but also on the surrounding environment. We show experimentally that this scenario breaks down at a non-Hermitian degeneracy known as an exceptional point. We find a chirality-reversal phenomenon in a ring cavity where the radiation field reveals the missing dimension of the Hilbert space, known as the Jordan vector.

09:40 : Hiding Parity-time Symmetry by Transformation Optics

Xinghong Zhu¹, Hongfei Wang², Danyuan Lei², John Brian Pendry³, Jensen Li¹

¹The Hong Kong University of Science and Technology (Hong Kong), ²City University of Hong Kong (Hong Kong), ³Imperial College London (United Kingdom)

We investigate how exceptional points can be constructed by using transformation optics (TO). By transforming a seed structure with conventional PT symmetry, we theoretically generate non-Hermitian daughter systems which do not possess PT symmetry in the usual sense but with inherited exceptional points and PT-phase transitions.

09:55 : Invited talk

Restricted Hilbert transform for a non-Hermitian management of electromagnetic fields

Waqas Ahmed¹, Muriel Botey², Ramon Herrero², Ying Wu¹, Kestutis Staliunas²

¹King Abdullah University of Science and Technology (KAUST) (Saudi Arabia), ²Universitat Politècnica de Catalunya (UPC) (Spain)

Spatial symmetry breaking in non-Hermitian systems allows tailoring wave propagation. Inspired by such property, we propose a local Hilbert Transform to control the field flows in non-Hermitian systems, restricting the complex refractive index within practical limits. We propose an iterative procedure, and show, how such procedure limits the space of complex refractive index within desired ranges. The proposed method provides a flexible way to systematically design locally PT-symmetric systems realizable with a limited collection of realistic materials.

10:15 : Invited talk

Control of Light Reflection and Transmission with Active Metasurface

Vasily V. Klimov

Lebedev Physical Institute (Russia)

New approach to all-optical control of light propagation with active metasurfaces is suggested. This approach is based on modification of losses in meta-atoms with an external light beam. The efficiency and quickness of the response of our metasurface are achieved by using complicated structures of eigen-oscillations in meta-atoms consisting of lossy and gain nanoparticles of an identical shape. The approach suggested paves the way for developing new active metasurfaces with a fast and substantial response to control light beam.

10:35 : Invited talk

A universal form of one-dimensional complex potentials featuring spectral singularities

Vladimir Konotop, Dmitry A. Zezyulin

Universidade de Lisboa (Portugal)

A one-dimensional complex potential in the Schrodinger equation features spectral singularities if and only if it has a universal form. Respective solutions have a universal form too. This allows one to construct complex potentials enabling either coherent perfect absorption or to lasing, or to both, at any prescribed wavelength. The described potentials allow for deformations leading to bound states in continuum or to exceptional points. We also describe potentials resulting in two or in three spectral singularities at desirable wavelengths.

09:00 - 10:45 — Lawrence Bragg Room

Session 2A9

Plasmonics: Fundamentals and Applications

Organized by: Hong Wei

Chaired by: Hong Wei

09:00 : **Invited talk****Light Assisted Synthesis of Novel Plasmonic Nanostructures for Single-molecule Surface-enhanced Raman Scattering and Nanofocusing****Zhipeng Li***Capital Normal University (China)*

The introduction of light during the synthesis process brings additional controllable freedom that enables new possibilities to tailor the physical parameters of nanomaterials. We have developed a two-step photo-reduction approach to generate highly reproducible single-molecule Surface-enhanced Raman Scattering (SM-SERS) materials. We also show that photochemical synthesis can create silver nano-needles with high aspect ratio and ultra-smooth surface. These nano-needles are excellent nanofocusing waveguides with divergent effective refractive index for plasmon propagating modes.

09:20 : **Invited talk****CQED in hybrid nanophotonic structures****Ying Gu, Zhiyuan Qian, Fan Zhang, Zhichao Li, Qi Zhang, Qihuang Gong***Peking University (China)*

We first propose an approach that combines a photonic crystal and metallic nanoparticle structure to create nanocavities with both strong local-field intensity and high helicity, where both strong and weak couplings with unidirectional propagation are obtained. Then, we propose the mechanism of edge state-led mode coupling under topological protection. Based on this mechanism, in topological photonic structure containing a resonant plasmon nanoantenna, an obvious absorption reduction in the spontaneous emission spectra appears.

09:40 : **Plasmonics in near-zero-index media****Ben Johns, Joy Mitra***Indian Institute of Science Education and Research (India)*

We investigate surface plasmon polaritons (SPPs) at the interface of a metal and a near-zero index (NZI) medium. Interestingly, these SPPs can be excited directly by free-space radiation as their dispersion lies above the free-space light line, in contrast to the case of conventional metal-dielectric interfaces. We explore unusual phenomena such as impedance matching of a NZI medium with free space and perfect electromagnetic absorption, and present designs for active and passive plasmonic waveguides incorporating NZI media.

09:55 : **Invited talk****The Coulomb Blockade in Plasmonics and How to Optically Lift it****Thomas A Klar, Dmitry Sivun***Johannes Kepler University Linz (Austria)*

We propose that a Coulomb blockade can suppress the tunneling current in quantum plasmonics in case of very small nanoparticles. Hence, a redshift is sustained even for sub-nanometer approach. This holds up to moderate fields that do not surpass the Coulomb blockade. Only for stronger optical fields, the Coulomb blockade is lifted and a charge transfer plasmon can be formed.

10:15 : **Measuring the temperature of plasmonic systems in ultrafast pump-probe experiments****Maria Sygletou¹, Marzia Ferrera¹, Giuseppe Della Valle², Michele Magnozzi¹, Daniele Catone³, Patrick O'Keeffe³, Alessandra Paladini³, Francesco Toschi³, Stefania Benedetti⁴, Piero Torelli⁴, Lorenzo Mattera¹, Maurizio Canepa¹, Francesco Bisio⁵***¹ University di Genova (Italy), ² IFN-CNR - Politecnico di Milano (Italy), ³ CNR-ISM (Italy), ⁴ CNR-Istituto Officina dei Materiali (Italy), ⁵ CNR-SPIN (Italy)*

After light-metallic nanoparticles (NPs) interaction, the system subcomponents, such as electron gas, lattice

and environment, gradually return to equilibrium by means of complex dynamic relaxation processes. Up to now, the absence of direct experimental method for measuring the dynamic temperature evolution of each system subcomponents was a major hurdle in understanding such processes. In this work, we discuss different methods for directly measuring the ultrafast evolution of the electronic temperature in metallic NPs, impulsively excited by ultrafast radiation pulses.

10:30 : Tunable Optical Response of Plasmonic Au Nanoparticles Embedded in Ta-doped TiO₂ Transparent Conductive films

Cristina Mancarella¹, Maria Sygletou², Beatrice Roberta Bricchi¹, Francesco Bisio³, Andrea Li Bassi¹
¹Politecnico di Milano (Italy), ²University of Genova (Italy), ³Consiglio Nazionale delle Ricerche-CNR-SPIN (Italy)

Localized Surface Plasmon Resonances (LSPR) of Au nanoparticles can be tailored in the visible range through nanoparticle geometry. However modulation is limited by the fixed carrier density of metals. Embedding Au nanoparticles in Transparent Conductive Oxides (TCOs) is an original approach to widen LSPR tunability by acting directly on the surrounding host. In this contribution we show that the easily modifiable permittivity of the TCO matrix succeeds as an additional degree of freedom in tuning properties.

09:00 - 10:45 — Rene Descartes Room

Session 2A10

Metamaterial enabled new devices and applications

Organized by: Weiren Zhu, Ciyuan Qiu and Fajun Xiao

Chaired by: Weiren Zhu, Ciyuan Qiu and Fajun Xiao

09:00 : Invited talk

Metamirror for generation and control of Airy beams

Rui Feng¹, Badreddine Ratni², Jianjia Yi³, André de Lustrac⁴, Hailin Zhang¹, Shah Nawaz Burokur²
¹Xidian University (China), ²Université Paris Nanterre (France), ³Xi'an Jiaotong University (China), ⁴Université Paris-Saclay (France)

Due to their intriguing diffraction-free, self-bending, and self-healing properties, Airy beams have attracted enormous research interests. An electronically reconfigurable metamirror is proposed to generate Airy beams on a wide frequency range by designing two distinct coding states with opposite reflective phase of 0° and 180° as digital bits of "0" and "1" states. Both numerical simulations and experimental measurements are performed to verify the Airy beam properties from 9 GHz to 12 GHz.

09:20 : Invited talk

Planar Vortex Beam Generator for Circularly Polarized Incidence Based on FSS

Yuxiang Wang¹, Kuang Zhang¹, Yueyi Yuan¹, Badreddine Ratni², Shah Nawaz Burokur², Qun Wu¹
¹Harbin Institute of Technology (China), ²Université de Paris Nanterre (France)

A new technique to design a low-profile planar vortex beam generator is proposed based on microwave frequency selective surface (FSS). Each unit cell is composed of a stack of patches and grids separated by thin dielectric substrates. A simple equivalent circuit model, composed of transmission lines coupled together with shunt capacitors and inductors, is presented to analyze this structure. The prototype of the proposed planar OAM generator operating in X-band is designed, fabricated and experimentally characterized.

09:40 : Invited talk

Plasmon Nano-tweezers towards sub 10 nm particles trapping

Fajun Xiao, Jianlin Zhao

Northwestern Polytechnical University (China)

In this work, we demonstrate the radially polarized beam can excited an annular potential well in a plasmonic coaxial aperture for stably trapping the 5 nm particles under a relatively low optical power. Furthermore, we demonstrate the plasmon mode residing in the closely spaced silver coated fiber tip and gold film can

produces a sharp quasi-harmonic potential well, capable of stably trapping 2 nm quantum dot beneath the tip apex with the laser power as low as 3.7 mW.

10:00 : Programmable absorbing metasurface for active scattering manipulation

Linda Shao, Weiren Zhu

Shanghai Jiao Tong University (China)

We present a programmable absorbing metasurface for active scattering manipulation. A metasurface unit with switchable perfect absorption and perfect reflection is achieved by incorporating a PIN diode into a typical metamaterial absorber, where the absorption and reflection functions can be switched by biased voltages. The simulated field results shows that the scattering properties can be controlled by changing the voltage distribution on PIN diodes on the metasurface.

10:15 : Metamaterial Absorbers for Infrared Stealth Applications

Jinglan Zou, Cong Quan, Zhihong Zhu, Xiaodong Yuan, Jianfa Zhang

National University of Defense Technology (China)

We explore selective metamaterial absorbers for infrared stealth applications. Several different designs will be discussed. We show that with multi-band absorption, the designed metamaterials can suppress the scattering near infrared signals used by laser-guided missiles, and the mid-infrared emission by matching the atmospheric absorption band. High-temperature metals for metamaterial designs, such as tungsten and molybdenum, will be discussed.

10:30 : Superscattering and superdirective emission via mode stacking in subwavelength meta-atoms

Alex W. Powell, Alastair Hibbins, John Roy Sambles

University of Exeter (United Kingdom)

Designing a subwavelength meta-atom so that multiple resonances occur at the same frequency can vastly enhance its interaction with electromagnetic radiation, as well as its directivity. However, experimentally demonstrating this effect has previously proven difficult. We demonstrate that such mode stacking, leading to superscattering and superdirectivity of emission, can be readily achieved through the careful structuring and arrangement of core-shell dielectric spheres. This work has applications in diverse fields such as antenna design, imaging and optoelectronics.

Coffee Break and Exhibit Inspection

Session 2P1

Poster session III

11:00 - 11:40

P1: Single-walled carbon nanotube phase shifters for low THz frequencies

Aleksandra Przewłoka¹, Serguei Smirnov², Aleksandra Krajewska¹, Joachim Oberhammer², Mikhail Khodzitsky³, Dmitri Lioubtchenko¹

¹CENTERA Laboratories (Poland), ²KTH Royal Institute of Technology (Sweden), ³ITMO University (Russia)

In this work single-walled carbon nanotube length dependence on phase tuning properties of dielectric rod waveguide is experimentally studied in ultra-wide frequency band of 0.1-0.5 THz.

P2: Structural Morphological and Optical Studies Of La-doped ZnS Thin films synthesized by Chemical Bath Deposition Technique in Acidic medium.

Hamid Merzouk, Djahida Talantikit

Algeria University (Algeria)

La-doped ZnS thin films have been produced on glass substrates using a chemical bath deposition route. X ray diffraction, atomic force microscopy, scanning electronic Microscopy, spectrophotometer and photoluminescence characterizations are used to study structural, morphological and optical properties of our samples. The lanthanum concentration was varied from 0 % to 10 % Spectroscopic analyses have shown significant improvement of optical transmittance reaching 90 % for films doped at 1 % in the visible but For all others the transmission do not exceed 80 %

P3: Gold Nanoparticle Arrays on Flexible Substrate for Stress Measurements**William d'Orsonnens¹, Florian Lamaze¹, Abdelhamid Hmima¹, Julien Proust¹, Aymeric Leray², Thomas Maurer¹, Eric finot²**¹University of Technology of Troyes (France), ²University of Burgundy (France)

Gold nanoparticles, and gold nanoparticles arrays, have widely been used as sensors, especially in the biology field due to their plasmonic properties. They present strong coupling in the visible range which makes them easy to observe. We therefore aim to use nanoparticles arrays to create a mechanical strain test sensor based on Fano effect. To do so we developed techniques to deposit gold nanoparticles on flexible substrates.

P4: Scattering properties of non-reciprocal systems with gain/loss**Hamed Ghaemidzicheh***Lancaster University (United Kingdom)*

We develop a transfer matrix description for non-reciprocal media with gain and loss, and show how topological signatures in these systems can be detected in transport experiments. This includes signatures of the non-Hermitian skin effect, for which we clarify the role of scattering boundary conditions.

P5: Exciton diffusion and annihilation in nanophotonic landscapes**T. V. Raziman, C. Peter Visser, Alberto G. Curto***TU/e - Eindhoven University of Technology (The Netherlands)*

Excitonic emitters in semiconductors exhibit diffusion and annihilation. Conventional nanophotonics improves light emission by providing enhancements of excitation, emission efficiency, and collection, but neglects exciton dynamics. We exploit exciton dynamics for improving emission, going beyond the localized Purcell effect. We present guidelines to benefit from diffusion and to ameliorate the effects of annihilation. We identify the dominant mechanisms for enhancement for limits of diffusion and annihilation. Controlling exciton dynamics has direct implications for light-emitting devices based on excitonic nanomaterials.

P6: Tunable and robust long-range coherent dipole interactions mediated by Weyl bound states**Inaki Garcia-Elcano¹, Alejandro Gonzalez-Tudela², Jorge Bravo-Abad¹**¹Universidad Autonoma de Madrid (UAM) (Spain), ²Instituto de fisica Fundamental (IFF-CSIC) (Spain)

The interaction between a quantum emitter and a Weyl photonic environment is investigated. We show that a photon-atom bound state is formed at the proximities of the Weyl frequency. Remarkably, the photonic part of such light-matter hybrid state features a tunable and robust power law spatial confinement. When considering more than one emitter couple to the bath we demonstrate that the emergent Weyl bound state can be harnessed to mediate long-range interactions between emitters.

P7: Accurate Circuit Model for Periodic Array of Square Patches**Saeed Zolfaghary, Amin Khavasi, Behzad Rejaei***Sharif University of Technology (Iran)*

A novel circuit model for the grid impedance of electrically dense arrays of subwavelength metallic square patches is derived using analytical formulas for arrays of metallic square holes. By comparison with full-wave simulations, we show that the model can predict the reflection and transmission of frequency selective surface with high accuracy.

P8: Designing Scattering Properties of Metasurfaces**James Capers, Simon Horsley, Alastair Hibbins***University of Exeter (United Kingdom)*

Metasurfaces have been shown to enhance the Local Density of Optical States (LDOS), providing large improvements in the power emission of dipole emitters. Many experimental works have focused on the properties of metasurfaces composed of square arrays of scatterers. In this work, we develop a method of designing metasurfaces to produce the desired scattering properties, focusing on the application to antenna design. Using this method, we demonstrate a three-fold improvement in the LDOS compared to a square array.

P9: Equivalent Circuit Model for Charactering Hysteresis of Perovskite Solar Cells**Ting Xu¹, Zishuai Wang², Wei E. I. Sha¹**¹Zhejiang University (China), ²The University of Hong Kong (China)

Equivalent circuit model is a useful tool to characterize device physics of perovskite solar cells (PVSCs). Based on detailed balance theory and dynamics of carriers and ions, an equivalent circuit model is proposed

for modeling current-density voltage characteristics of PVSCs. The dynamic hysteresis phenomena are reproduced and the connections between the device properties and circuit elements are established. Evidently, the circuit model offers systematic descriptions to the working mechanisms of PVSCs, which is beneficial to design and performance optimization.

P10: New Plasmonic System for Visible Light-Driven Hydrogen Evolution Reaction

Hiro Minamimoto, Daiki Sato, Kei Murakoshi

Hokkaido University (Japan)

Excitation of the localized surface plasmon resonance leads to the generation of the reaction active species. The excited species makes it possible to trigger the efficient multi electron transfer reactions. In this study, plasmon-induced hydrogen evolution reactions have been achieved by the introduction of the plasmonic metal nanostructures into the p-type GaP semiconductor electrode. Through the photoelectrochemical measurements, the unique molecular process has been confirmed at the present plasmonic photoconversion electrode, resulting in the unique reaction selectivity.

P11: Enhanced light generation due to hybridization of lattice and gap plasmon modes in periodic MIM tunnel junction

Saurabh Kishen, Jinal Tapar, Naresh Kumar Emani

Indian Institute of Technology Hyderabad (India)

A promising approach to realize electrically excited on-chip nanoscale optical sources is through inelastic electron tunneling. Its practical implementation, however, suffers from low electron-to-photon transduction efficiencies. Here, we investigate the enhancement of light generation in a periodic Ag-SiO₂-Ag tunnel junction due to inelastic electron tunneling. By efficiently coupling lattice resonance with gap plasmon mode, we achieve an enhancement in the local density of optical states by three orders of magnitude and a radiative efficiency 30% higher than the uncoupled structure.

P12: Large area fabrication of plasmonic nanostructures using nanoimprint lithography with contact transfer

Jun Hyun Kim, Doo-In Kim, Myung Yung Jeong

Pusan National University (Korea)

In this study, the nanoimprint lithography (NIL) process was proposed and demonstrated to fabricate the plasmonic nanostructures. An embossing of a PMMA substrate and the transfer of a deposited metallic nanostructures were implemented simultaneously using a thermal-NIL for the cost-effective large area fabrication of the plasmonic nanostructures.

P13: Fabrication of 1-D Photonic crystals to enhance thermochromic properties of VO₂ nanostructures

Dipti Umed Singh¹, Omkar Bohite¹, Remya Naryanan²

¹Indian Institute of Science Education and Research (India), ²Savitribai Phule Pune University (India)

Effect of 1-D photonic crystals on optical transmission of VO₂ is studied by depositing VO₂ thin films on Distributed Bragg Reflectors (DBR) in the infrared (IR) spectrum. Monoclinic VO₂ nanoparticles were first synthesized by solution processed method. By combining VO₂ films on DBR structure, the average optical transmission approaches to zero in the IR region in and above the critical temperature. which could be the positional design for VO₂ nanoparticles based hybrid Photonic absorbers for various smart window applications

P14: Spectral singularities and non-reciprocal light scattering in 2D PT-symmetric metamaterials

Jinal Kiran Tapar, Saurabh Kishen, Saurabh Kishen, Naresh Kumar Emani

Indian Institute of Technology (India)

We consider vertically stacked GaInP PT-symmetric resonators that constitute meta-atoms for the 2D active metamaterial. From numerical simulations and scattering theory formalism, we show that these metasurfaces support zero-width resonances, i.e. spectral singularities (SS). By tuning the coupling between gain and loss resonators, we demonstrate the spectral singularities are robust over a wide range of parameter variations. We also show that vertically stacked GaInP resonators can exhibit broadband unidirectional invisibility and strong directional scattering.

P15: Small signal Frequency Response of a slotted photonic crystal based Raman laser

Akash Kumar Pradhan, Mrinal Sen*Indian Institute of Technology (ISM) (India)*

Small signal frequency response of a silicon nanocrystal based slotted photonic crystal Raman laser has been evaluated theoretically by solving the coupled mode equations for laser in steady state. Modulation bandwidth in the order of ~ 144 GHz has been achieved in this type of laser due to the high Raman gain coefficient of silicon nanocrystal.

P16: Fluorescent Microdiamonds Conjugated with Hollow Gold Nanoparticles as Photothermal fiducial Markers**Shan-Jen Kuo¹, Sih-Wei Chang¹, Yuen-Yung Hui², Oliver Y. Chen², Yen-Wei Chen², Ching-Che Lin¹, Dehui Wan³, Huan-Cheng Chang², Hsuen-Li Chen¹**¹*National Taiwan University (Taiwan)*, ²*Academia Sinica (Taiwan)*, ³*National Tsing Hua University (Taiwan)*

In this study, we characterized fluorescent microdiamonds (FMDs) and explored their potential use as fiducial markers for image guided photothermal therapy. To exploit the excellent thermal conductivity of diamonds, we decorated the surface functionalized FMDs with hollow gold nanoparticles (HGNs). The dual functions fluorescent and thermal imaging of the HGN-FMDs were performed when applying the distinct excitation laser wavelengths of the HGNs and the FMDs, respectively.

P17: Sub-wavelength imaging of few-cycle nanoplasmonic near-fields using interferometric time-resolved photoemission electron microscopy**Alexander Gliserin¹, Soo Hoon Chew¹, Sungho Choi², Dong Eon Kim², Seungchul Kim¹**¹*Pusan National University (Korea)*, ²*Pohang University of Science and Technology (Korea)*

We present a first implementation of a novel perturbative autocorrelation technique for direct sampling of ultrafast electric fields with a photoemission electron microscope. Unlike conventional nonlinear autocorrelation, this method preserves the phase, providing a full characterization of the electric field without additional spectral information. This allows direct observation of sub-wavelength electric near-fields at plasmonic nanostructures on the shortest time scales, which we demonstrate at a monolayer of self-assembled 40-nm gold nanospheres with sub-100-nm spatial resolution.

P18: 3D Metastructure Design for Noise Suppression of Audible Frequency Band**Sung-Sil Cho, Ic-Pyo Hong***Kongju National University (Korea)*

We fabricated a simple acoustic spectrum measurement system for measuring acoustic transmission loss and proposed a new acoustic metastructure that can reduce noise. To validate the performance of the designed system, the previously studied acoustic metastructure was generated and the acoustic transmission loss was measured and compared. In addition, the novel acoustic metastructure was proposed for noise reduction. The simulation results show that the proposed acoustic metastructure has a loss of 67dB at 478Hz and a bandgap of 448-546Hz.

P19: Synthesis of gold nanoparticles using α -amino acids**Aleksandra Maria Figat, Bartosz Bartosewicz, Malwina Liszewska, Bartłomiej J. Jankiewicz***Military University of Technology (Poland)*

Various α -amino acids have been used as reducing and stabilizing agents in the synthesis of gold nanoparticles following the Turkevich protocol. The shape, size distribution, stability and optical properties of synthesized nanoparticles were characterized by SEM, DCS, PALS technique and UV-vis spectroscopy. The differences in chemical structure of α -amino acids strongly affect their reactivity and influence the shape, size distribution and stability of synthesized gold nanoparticles.

11:40 - 12:30 — Victor Veselago Room**Session 2A11****Symposium II: New trends in nanophotonics and advanced materials**

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

11:40 : Keynote talk

Commercializing Metaphotonics

Hyuck Choo

Samsung Electronics Co., Ltd. (Korea)

According to a market research organization, the metaphotonics market is expected to exceed \$10B in 10 years, and its growth be fueled by the rapidly spreading IoT and related services powered by the 5G network. But, what are the successful commercialization examples of metamaterials that we know? In this presentation, I will share some of our recent results and on-going commercialization efforts for metaphotonics utilizing Si-based IC-processing technologies. A few areas of metamaterial applications include safer autonomous driving; custom-tailored point-of-care health monitoring and diagnosis; and improving light-capturing efficiency of sub-micron-pixel CMOS imagers. I look forward to share and discuss our findings and vision for the coming years.

12:10 : Invited talk

Bi-physical stealth metamaterials for waterborne objects

Yi Zhou¹, Jian Chen¹, Rui Chen¹, Wenjie Chen¹, Liu Liu², Zheng Fan³, Yungui Ma¹

¹Zhejiang University (China), ²South China Normal University (China), ³Nanyang Technological University (Singapore)

In this work, a bi-physical ultrathin metamaterial that could simultaneously make underwater objects invisible for magnetic field and acoustic wave is demonstrated and validated. This coat could shield the magnetic field with the field disturbance ratio less than 0.5% over a broad frequency range (10-250 kHz) and strongly attenuate ultrasonic waves with a nearly unity absorptivity meanwhile. This metamaterial may shed light on building multifunctional devices for various waterborne applications.

11:40 - 12:40 — Allan Boardman Room

Session 2A12

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

11:40 : Invited talk

Polarization state generator obtained by self-assembled plasmonic nanoparticles

Yann Battie¹, Jie Gao², Emilie Pouget², Reiko Oda², Laurent Broch¹, Matthias Pauly³, Aotmane En Naciri¹

¹Université de Lorraine (France), ²Université de Bordeaux (France), ³ICS (UPR22-CNRS) (France)

In this work, we explore the optical properties of gold nanoparticles assembled into chiral structure. This structure exhibits a complex multianisotropy which comes from the dipolar interaction between NPs. We show that these nanostructures are good candidates for new polarization devices.

12:00 : Invited talk

Precisely positioned atomistic quantum emitters in monolayer MoS₂

Alexander Holleitner

Technische Universität München (Germany)

For photonic and plasmonic hybrid circuits, it is essential to place single photon emitters with the highest lateral position possible to allow for a controlled exploitation of the local electromagnetic fields. We demonstrate the deterministic generation of single defect emitters in a monolayer MoS₂ van der Waals heterostructure with a lateral creation accuracy of ~9 nm. Our work paves way towards the controlled generation of atomistic single photon emitters in monolayer 2D materials embedded in optoelectronic quantum devices.

12:20 : Invited talk

Overcoming Limits in Nano-Optical Simulations, Design and Experiments Using Deep Learning**Peter R. Wiecha, Guilhem Larrieu, Aurélie Lecestre, Otto L. Muskens***LAAS-CNRS Toulouse France (France)*

Tremendous research efforts have been put into the field of nano-optics, leading to applications like flat optics or negative index metamaterials. However, there are physical and/or methodological constraints, hard to overcome. For instance, the optical diffraction limit is a difficult obstacle in microscopy or optical information storage. Inverse design of nanostructures is another example for a difficult tasks. We show how very hard to solve problems can be tackled efficiently using methods of artificial intelligence and specifically deep learning.

11:40 - 12:20 — Tatsuo Itho Room**Session 2A13****Bottom-up approaches, new fabrication routes and ENSEMBLE3**

Organized by: Dorota Pawlak and Virginie Ponsinet

Chaired by: Dorota Pawlak and Virginie Ponsinet

11:40 : Invited talk**Organic Polymeric-carbon-nitride-based Metamaterials: A Novel Platform for Nano-optical Device Innovation****Daniel Clarke¹, Julya Siena², Corinna Kaspar³, Wolfram Pernice³, Markus Antonietti², Ortwin Hess¹**¹*Trinity College Dublin (United Kingdom)*, ²*Max Planck Institute of Colloids and Interfaces (Germany)*, ³*University of Munster (Germany)*

In this contribution, we present a combined theoretical and experimental investigation of pCN-coated, gold double-fishnet metamaterials. Our full-dimensionality, finite-difference time-domain calculations are complemented with state-of-the-art device fabrication procedures which make possible the realization of such pCN-based metamaterials, and we compare our theoretical predictions directly with experimental measurements.

12:00 : Invited talk**Gold nanoparticles: from plasmonic field enhancement and luminescence to nanofabrication****Céline Molinaro¹, Sylvie Marguet¹, Ludovic Douillard¹, Fabrice Charra¹, Farid Kameche², Olivier Soppera², Dandan Ge³, Renaud Bachelot³, Céline Fiorini-Debuisschert¹**¹*Université Paris-Saclay (France)*, ²*Université de Haute-Alsace (France)*, ³*Université de Technologie de Troyes (France)*

The two-photon-excited luminescence of gold nanoparticles is studied in detail. A phenomenological model is proposed which reveals the importance of field enhancement at the plasmonic NP resonances for both plasmon enhanced absorption and plasmon enhanced emission, with surface effects also playing an important role. Beyond these fundamental aspects, we show that plasmonic hot spots can also interestingly be taken into profit for the realization of advanced hybrid nanostructures for photonics

11:40 - 12:40 — Christian Huygens Room**Session 2A14****Non-Hermitian Photonics: From Plasmonics, and Metamaterials, to Topological and Quantum systems**

Organized by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

Chaired by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

11:40 : Invited talk**Stability optimization of random networks with added non-Hermitian nodes from a parity-time symmetry perspective****Henri Benisty***Institut d'Optique Graduate School (France)*

A network being described by a real symmetric coupling matrix between its elements, we look at the addition of gain and loss elements coupled to the network, to maximize the so-called unbroken phase of the parity-time symmetry approach now familiar in optics: we minimize imaginary parts of eigenvalues. We explore with Fourier tool how coupling should be arranged and attempt to find rules similar to those established for "stability-optimized-circuits" in the context of neural networks.

12:00 : Invited talk**Non-Hermitian media with global and local unidirectionality: theory and applications****Muriel Botey, Waqas W. Ahmed, Zeki Hyaran, Judith Medina, Hamza Kurt, Ramon Herrero, Kestutis Staliunas***Universitat Politècnica de Catalunya (UPC) (Spain)*

Structured media provide the momentum compensation for the scattering of waves, shaping the the propagation of light. Yet only the interplay between both the real and imaginary modulations introduces unidirectionality in the light management. A generalized Hilbert transform allows tailoring the two quadratures of the complex permittivity to design periodic or disordered non-Hermitian media, holding either global or local unidirectionality following arbitrary vector fields. Moreover, the method allows restricting the permittivity within realistic values rendering it suitable for applications.

12:20 : Invited talk**Ultrathin Acoustic Parity-Time Symmetric Metasurface Cloak****Johan Christensen¹, María Rosendo-López¹, Hao-Xiang Li², Yi-fan Zhu², Xu-dong Fan², Daniel Torrent³, Bin Liang², Jian-chun Cheng², Johan Christensen¹**¹*Universidad Carlos III de Madrid (Spain)*, ²*Nanjing University (China)*, ³*Universitat Jaume (Spain)*

By using an elaborate arrangement of gain and lossy acoustic media respecting parity-time symmetry, we built a one-way unhearability cloak able to hide objects seven times larger than the acoustic wavelength. Generally speaking, our approach has no limits in terms of working frequency, shape, or size, specifically though we demonstrate how, in principle, an object of the size of a human can be hidden from audible sound.

11:40 - 12:30 — Augustin Fresnel Room**Session 2A15****Symposium IV: Chirality, magnetism, and magnetoelectricity: Separate phenomena and joint effects in metamaterial structures**

Organized by: Eugene Kamenetskii

Chaired by: Eugene Kamenetskii

11:40 : Bi-hyperbolic Isofrequency Topologies in a Gyroelectromagnetic Medium**Vladimir Tuz¹, Volodymyr Fesenko²**¹*Jilin University (China)*, ²*National Academy of Sciences of Ukraine (Ukraine)*

Isofrequency topologies are studied for a gyroelectromagnetic medium. This medium is obtained by stacking optically thin magnetized ferrite and semiconductor layers into a unified structure. In such a structure, a bi-hyperbolic isofrequency contour appears as a simultaneous effect of both periodic arrangement of constitutive layers and external magnetic field influence. It is proposed to consider the obtained bi-hyperbolic isofrequency contour as a new topology class of the wave dispersion.

11:55 : Invited talk**Antiferromagnetism, chirality, and magneto-optical effects**

Jonathan Kipp¹, Fabian Lux¹, Kartik Samanta¹, Maximilian Merte¹, O. Gomonay², Frank Freimuth¹, Marjana Lezaic¹, Stefan Blugel¹, Yuriy Mokrousov²

¹Peter Grunberg Institut (Germany), ²University of Mainz (Germany)

We study the influence of chirality on magneto-optical phenomena in frustrated magnets. By referring to models and microscopic calculations we suggest a way to identify the sense of crystal and magnetic chirality of two-dimensional magnets from the behavior of the Hall and magneto-optical effects that they exhibit, promoting new protocols for probing these fundamental properties of matter.

12:15 : Full Manipulation of Circular Polarizations for Multiple Vortex Beams Generation based on Phase-Engineered Metasurface

Kuang Zhang¹, Yueyi Yuan¹, Badreddine Ratni², Shah Nawaz Burokur², Qun Wu¹, Patrice Genevet³

¹Harbin Institute of Technology (China), ²Université Paris Nanterre (France), ³Université Côte d'Azur (France)

In this paper, a phase-engineered non-interleaved metasurfaces is proposed for multiple vortex beam generation, which is combining versatile phase modulation methods. Different from traditional scheme based on phase interleaving for multiple functionalities, four vortex beam wavefronts carrying independent orbital angular momentum (OAM) can be created through all four circular polarization (CP) transmission channels. Theoretical simulations are conducted and effectively verified the feasibility of the proposed theory for artificial manipulation of CP manipulation in microwave region.

11:40 - 12:45 — Ibn Al-Haytham Room

Session 2A16

Exotic Meta-media - Time-dependent, Nonlocal and Other Novel Responses

Organized by: Martin McCall, Jonathan Gratus and Paul Kinsler

Chaired by: Martin McCall, Jonathan Gratus and Paul Kinsler

11:40 : Asymmetric frequency conversion with acoustic non-Hermitian space-time varying metamaterial

Xinhua Wen¹, Xinghong Zhu¹, Alvin Fan¹, Wing Yim Tam¹, Jie Zhu², Fabrice Lemoult³, Mathias fink³, Jensen Li¹

¹Hong Kong University of Science and Technology (China), ²Hong Kong Polytechnic University (China), ³PSL University (France)

We experimentally realize an acoustic non-Hermitian space-time varying metamaterial using digital virtualized resonating meta-atoms. By temporally modulating the material gain and loss, we can diminish the main band and achieve high efficiency frequency conversion at the same time due to the gain-loss balance in time domain. We also experimentally demonstrate the asymmetric amplification with such an acoustic metamaterial.

11:55 : Invited talk

Analysis of nonlocal constitutive relations to homogenize metamaterials

Fatima Zohra Goffi¹, Karim Mnasri¹, Michael Plum¹, Carsten Rockstuhl¹, Andrii Khrabustovskiy²

¹Karlsruhe Institute of Technology (Germany), ²University of Hradec Kralové (Czech Republic)

It has been appreciated that nonlocal constitutive relations are more performant to homogenize metamaterials, which are obtained by approximating a general response function of the electric field describing the response of the metamaterial. In this research, a second order Padé approximation of the response function is adopted that leads to several formulations. To decide which formulation is consistent for the homogenization, we present a checklist each constitutive relation has to pass to be admissible.

12:15 : Wave scattering by layered structures in critical conditions

Kim Pham¹, Agnès Maurel²

¹IMSIA - ENSTA Paris (France), ²Institut Langevin (France)

We derive an effective model governing the light scattering in structures alternating metal/dielectric layers

of subwavelength thicknesses. The homogenization procedure is conducted on the Maxwell equations in three dimensions, the resulting model involves effective constitutive relations which link the electric field and electric displacement as well as non-intuitive transmission conditions at the extremities of the substructure. In transverse electric polarization, the model reduces to a fully local model while in transverse magnetic polarization, classical non local terms appear.

12:30 : 4-dimensional covariant helicity states in vacuum and linear media

Robert Thompson¹, Ivan Fernandez-Corbaton¹, Martin McCall²

¹Karlsruhe Institute of Technology (Germany), ²Imperial College London (United Kingdom)

The notion of helicity in solutions to Maxwell's vacuum equations in standard 3 + 1 notation is extended to media expressed in 4-D coordinate-free notation. This allows us to significantly generalise the definition of helicity that is associated with the (anti-)self-duality of solutions to Maxwell, a generalization that permits easy extension to media in curved spacetimes.

11:40 - 12:40 — Gaston Floquet Room

Session 2A17

Advanced modeling techniques for the design of metasurface devices

Organized by: Patrice Genevet and Stéphane Lanteri

Chaired by: Patrice Genevet and Stéphane Lanteri

11:40 : Invited talk

Metasurfaces for Divergent Beams and Large-Area Metasurfaces

Daniel Andrén, Jade Martinez-Llinas, Ruggero Verre, Mikael Kall, Philippe Tassin

Chalmers University of Technology (Sweden)

We will give an overview of our recent work on the design and fabrication of metasurfaces, i.e., dense arrays of subwavelength-sized scatterers (meta-atoms) designed in shape, size, position, and orientation. First, we have developed simulation strategies for large-scale metasurfaces and metasurfaces for strongly divergent beams. Second, we have developed a facile fabrication technique based on an exposed resist to build large-scale metasurfaces. These novel computational and fabrication techniques allow us to achieve metasurfaces with unprecedented functionality.

12:00 : Invited talk

Fundamental limitations of ultra-flat Huygens metasurfaces

Carlo Gigli¹, Pierre Chavel², Philippe Lalanne³

¹Université de Paris (France), ²Université de Lyon (France), ³Université de Bordeaux (France)

There are two physical effects that are exploited nowadays to implement flat metalenses, either subwavelength guidance implementing varying propagation delays, or resonant confinement combining two resonances. We compare both approaches and identify possible FUNDAMENTAL limitations with the second approach.

12:20 : Invited talk

Contour integral methods for resonance phenomena in nano-optics

Felix Binkowski¹, Fridtjof Betz¹, Remi Colom¹, Martin Hammerschmidt², Philipp-Immanuel Schneider², Lin Zschiedrich², Sven Burger¹

¹Zuse Institute Berlin (Germany), ²JCMwave GmbH (Germany)

We review contour integral methods for the solution of nonlinear eigenvalue problems resulting from Maxwell's equations. Numerical realizations of the methods are applied to compute and analyze resonances in nano-optical systems with material dispersion.

11:40 - 12:40 — Lawrence Bragg Room**Session 2A18****Machine learning for metamaterials and metasurfaces**

Organized by: Mohamed Bakr and Willie Padilla

Chaired by: Mohamed Bakr and Willie Padilla

11:40 : Invited talk**Inverse design in nanophotonics using deep-learning****Junsuk Rho***Pohang University of Science and Technology (POSTECH) (Korea)*

Recent introduction of deep learning into nanophotonics has enabled efficient inverse design process. Once the deep learning network is trained, it allows fast inverse design for multiple design tasks. In this talk, we show several inverse designing nanophotonic structures using deep learning. We firstly discuss inverse design methods that increase the degree of freedom of design possibilities.

12:00 : Invited talk**Deep Learning Approach for the Enhanced Light-Matter Interactions in Dielectric Nanostructures****Lei Xu¹, Mohsen Rahmani², Yixuan Ma¹, Daria A. Smirnova², Khosro Zangeneh Kamali², Fu Deng¹, Yan Kei Chiang¹, Lujun Huang¹, Haoyang Zhang³, Stephen Gould², Dragomir N. Neshev¹, Andrey E Miroshnichenko¹**¹*UNSW Canberra (Australia)*, ²*The Australian National University (Australia)*, ³*The Queensland University of Technology (Australia)*

In this paper, we utilize a deep-learning approach for obtaining high-quality factor (high-Q) resonances with desired characteristics, such as linewidth, amplitude and spectral position. We exploit such high-Q resonances for the enhanced light-matter interaction in nonlinear optical metasurfaces and optomechanical vibrations, simultaneously.

12:20 : Invited talk**A generalized accurate predictor of nano-optical near fields and far-fields using a deep learning neural network****Otto L. Muskens, Peter R. Wiecha***University of Southampton (United Kingdom)*

We present a deep learning artificial neural network (ANN) capable of predicting the full near-fields and far-fields of nanostructures. The ANN captures the direct relationship between the geometry and the internal fields. It successfully addresses plasmonic antenna modes, magneto-electric resonances of high-index dielectrics, electromagnetic anapole states, Kerker effects, near-field particle interactions and chiral hot spots. This new data-driven ANN approach to nano-optical modelling enables very fast evaluations and opens new routes to inverse design of nano-optical structures and metasurfaces.

11:40 - 12:40 — Rene Descartes Room**Session 2A19****Plasmonics for single molecule detection and manipulation**

Organized by: Denis Garoli

Chaired by: Denis Garoli

11:40 : Invited talk**Light-Induced Particle Binding Assisted by Metamaterial Substrates**

Alexander Shalin¹, Natalia Kostina¹, Pavel Ginzburg²

¹*ITMO University (Russia)*, ²*Tel Aviv University (Israel)*

Here we study light-induced interaction of several dielectric particles above a hyperbolic metamaterial. It is shown, that both surface and volumetric modes of the substrate define distances between interacting particles, beyond the diffraction limit. Moreover, by varying the thickness of the metamaterial substrate it is possible to tune magnitude of the optical force and distances between the particles.

12:00 : Invited talk

Protein-Tailored Plasmonic Silver Nanorings over Graphene-Coated Nanopores for Localized Enhanced Fluorescence

Matteo Ardini¹, Giorgia Giovannini², Nicolo Maccaferri³, Xavier Zambrana-Puyalto², Gloria Panella¹, Francesco Angelucci¹, Rodolfo Ippoliti¹, Denis Garoli², Francesco De Angelis²

¹*University of L'Aquila (Italy)*, ²*Istituto Italiano di Tecnologia (Italy)*, ³*University of Luxembourg (Luxembourg)*

Engineering electromagnetic fields through plasmons provides advances in several applications. Nanodevices with improved optical properties, for instance, are obtained using the architecture of proteins and their affinity towards nanomaterials. Here, silver nanorings are synthesized on the ring protein Prx and arranged over graphene-coated nanohole arrays achieving improved and localized fluorescence. This approach represents a proof-of-concept for future nanopore-based technologies, e.g. next-generation sequencing and single-molecule detection.

12:20 : Invited talk

Plasmonic Sensing by Gold Nano-Mushroom Arrays Fabricated using a Novel Anisotropic Etching Regime

Daniel Darvill¹, Marzia Iarossi¹, Ricardo M. Abraham Ekeroth², Aliaksandr Hubarevich¹, Francesco De Angelis¹

¹*Istituto Italiano di Tecnologia (Italy)*, ²*Universidad Nacional del Centro de la Provincia de Buenos Aires (Argentina)*

We report a novel fabrication route for creating a plasmonic sensing platform based upon arrays of gold nano-mushrooms. This technique allows for a single processing step to create polystyrene mushrooms from colloidal masks produced using colloidal lithography prior to metal deposition. We show the potential of this plasmonic nanostructure as a biosensor tuned to the near infra-red with efficacy similar to those found in literature.

Lunch

12:40 - 14:00

14:00 - 15:10 — Victor Veselago Room

Session 2A20

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

14:00 : Keynote talk

Prineha Narang

Harvard University (USA)

Keynote

14:30 : Invited talk

Few-femtosecond plasmon transients probed with nm-scale sensitivity

Peter Dombi

Wigner Research Centre for Physics (Hungary)

Photoelectron probing of few-femtosecond plasmon transients on nanostructures reveals the ultrafast dynamics of localized plasmon oscillation decay with nm-scale sensitivity at plasmonic hot spots.

14:50 : Invited talk

Hyperuniform and Local Self-Uniform Solar Light Absorbers

Marian Florescu¹, Nasim Tavakoli², Richard Spalding¹, G. Gkantzounis¹, Chenglong Wan¹, Ruslan Röhrich², Evgenia Kontoleta², Femius Koenderink², Riccardo Sapienza³, Esther Alarcon Llado²

¹University of Surrey (United Kingdom), ²NWO-I Amolf (The Netherlands), ³Imperial College London (United Kingdom)

We explore the ability of hyperuniform disordered structures to improve light absorption in thin-film architectures and show that hyperuniform and local self-uniform correlations may be designed to enhance the coupling to quasi-guided modes supported by the thin film. We report a theoretical solar energy absorption of 84 % in a broadband spectral range (400-1050 nm), in a one micron-thick Si membrane, which is, to the best of our knowledge, the best value achieved in ultra-thin Si membranes.

14:00 - 15:20 — Allan Boardman Room

Session 2A21

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

14:00 : Invited talk

Degenerate ZnO nanocrystals and nanowires for Mid IR plasmonics

Bruno Masenelli¹, Mohamed K. H. Taha¹, Olivier Boisson¹, Patrice Mélinon¹, K. Masenelli-Varlot¹, Vincent Sallet², Isabelle Lefebvre³, Christophe Delerue³

¹Université de Lyon (France), ²Université de Versailles- St Quentin (France), ³Université de Lille-ISEN (France)

We investigate Ga doped degenerate ZnO nanocrystals and nanowires. We show that tunable mid IR plasmons are achieved in both cases. The dopant location and activation within the nanostructures have to be carefully controlled in order to induce efficient plasmons.

14:20 : Invited talk

Dynamic Polarization Control Using Temperature Driven Nonlinearities in Anisotropic Metamaterials

Luke H. Nicholls, Jingyi Wu, Anastasiia Zaleska, Mazhar Nasir, Anatoly V. Zayats

King's College London (United Kingdom)

By altering the permittivity of an anisotropic metamaterial, achieved by exploiting the Kerr-type nonlinearity of metals, we demonstrate control over the polarization of light. This is accomplished in a reflection configuration, allowing higher signal intensities. Moving away from femtosecond pulse control illumination, we also investigate the effect of using CW control light. This is made possible by the designed epsilon near zero (ENZ) region of the nanorod metamaterial and its related nonlocal properties.

14:40 : Invited talk

Large scale fabrication of silicon Mie nanoresonators: an alternative to gold ?

Julien Proust¹, Pierre Michel Adam¹, Redha Abdeddaim², Anne Laure Baudrion¹, Jeremie Beal¹, Frederic Bedu², Thomas Begou², Johann Berthelot², Sebastien Bidault³, Nicolas Bonod², Wajdi Chaabani⁴, Abdallah Chehaidar⁴, Stephane Chenot⁵, Anne Laure Fehrembach², Bruno Gallas⁵, Maria F. Garcia Parajo⁶, Julien Lumeau², Artur Movsesyan¹, Mathieu Mivelle⁷, Igor Ozerov², Jerome Plain¹, Raju Regmi², Herve Rigneault², Jerome Wenger², Pamina M. Winkler²

¹University of Technology of Troyes (France), ²Aix-Marseille University (France), ³ESPCI (France), ⁴SFAX University (Tunisia), ⁵Sorbone University (France), ⁶ICFO Institut de Ciencies Fotoniques (Spain), ⁷Universite

Pierre et Marie Curie (France)

High index dielectric nanoparticles have been proposed for many different applications. The multiples optical properties of such called Mie resonators allowed us to question there use as an alternative to gold. Actually, the optical resonances, coupled with a strong nearfield can open perspective in plasmonic uses. Widespread utilization in practice also requires largescale production methods for crystalline silicon nanoparticles. We demonstrate a lowcost, and largescale fabrication method of crystalline spherical silicon colloidal Mie resonators in water, using a blender.

15:00 : Invited talk

A bottom-up approach to the fabrication of optical metamaterials utilizing the self-assembly of silver dodecahedral clusters

Lucien Roach, Laurent Lermusiaux, Véronique Many, Alexandre Baron, Romain Dezert, Etienne Duguet, Virginie Ponsinet, Serge Ravaine, Philippe Richetti, Philippe Barois, Mona Tréguer-Delapierre
CNRS-Université de Bordeaux (France)

A key requirement on 'meta-atoms' comprising optical metamaterials, is a light-induced magnetic response of similar magnitude to the electric response. It is desirable to assemble such materials through bottom-up approaches, requiring the production of colloidally dispersed 'meta-atoms'. However, many of the currently realizable candidate colloidal particles present too weaker magnetic response. Hence, there is a need for novel magnetically-polarizable meta-atoms. We present silver dodecapods as a strong candidate for such a role.

14:00 - 15:25 — Tatsuo Itho Room

Session 2A22

Symposium III: Advanced passive and active metasurfaces and zero-index optics

Organized by: Howard Lee, Yu-Jung (Yuri) Lu and Zi Jing Wong

Chaired by: Howard Lee, Yu-Jung (Yuri) Lu and Zi Jing Wong

14:00 : Invited talk

Materials for nonlinear optical metasurfaces

Augustine Urbas

Air Force Research Lab (USA)

We will explore recent work in linear and nonlinear metasurfaces for optical applications including novel materials exploration.

14:20 : Invited talk

Low-loss zero-index metamaterials based on a bound state in the continuum (BiC)

Haoning Tang¹, Clayton DeVault¹, Phil Camayd-Munoz¹, Danchen Jia², Yueyang Liu³, Fan Du⁴, Olivia Mello¹, Yang Li³, Eric Mazur¹

¹Harvard University (USA), ²Zhejiang University (China), ³Tsinghua University (China), ⁴Nankai University (China)

On-chip metamaterials with a refractive index of zero can achieve perfect spatial coherence, leading to various applications in nonlinear and quantum optics. Those applications are severely hampered by the out-of-plane radiation losses as a result of the small amplitude momentum of zero-index metamaterial. Here we eliminate the out-of-plane radiation losses by realizing zero index with BiC. Our experimental results show that a BiC zero-index metamaterial has a quality factor over 15000 at the zero-index wavelength.

14:40 : Keynote talk

Tunable and Time-Modulated Flat Optics

Harry A. Atwater, Ghazaleh Shirmanesh, Prachi Thureja, Jared Sisler, Ruzan Sokhoyan, Meir Grajower
California Institute of Technology (USA)

Metasurfaces offer tremendous opportunity for photonics, namely, to manipulate amplitude, phase, and po-

larization of electromagnetic waves with arrays of subwavelength nanoantennas, enabling systems with flat optical components featuring dramatically reduced size, weight and power. Currently most metasurfaces are 'static' and have functions that are fixed at the time of fabrication. By making the system tunable or reconfigurable in its phase, amplitude and polarization response through incorporation of electro-optical effects, one can achieve real-time control of optical functions and indeed achieve multi-functional characteristics after fabrication. Despite the real-time tunability offered by the electro-optical metasurfaces, their operation thus far has been limited to the quasi-static regime, where temporal variations are slow enough that they do not result in changes in scattered beam frequency. By contrast, time modulated metasurfaces are temporally at rates high enough to generate new frequencies. Introducing time modulation to these metasurfaces opens a four-dimensional design space which can be used to overcome several fundamental limitations associated with static and quasi-static metasurfaces.

15:10 : Transdimensional photonic lattices with van der Waals metasurfaces and strong coupling to high-index thin layers

Viktoriia Babicheva

University of New Mexico (USA)

Transition metal dichalcogenides (TMDCs) from the family of van der Waals layered materials have been shown to exhibit tailorable optical properties along with strong nonlinearity, high refractive index, and anisotropy. We envision that TMDCs is a promising material platform for designing ultra-thin optical elements. We investigate a van der Waals metasurface that consist of disk-shaped nanoantennas made of TMDC material placed on top of a thin intermediate layer of high-index material such as silicon and low-index oxide substrate.

14:00 - 15:20 — Christian Huygens Room

Session 2A23

Non-Hermitian Photonics: From Plasmonics, and Metamaterials, to Topological and Quantum systems

Organized by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

Chaired by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

14:00 : Invited talk

Quantum correlations in PT-symmetric systems

Francesco Ciccarello

University of Palermo (Italy)

We study the dynamics of correlations in a paradigmatic setup to observe PT-symmetric physics: a pair of coupled oscillators, one subject to a gain one to a loss. Quantum correlations (QCs) are created, despite the system being driven only incoherently, and can survive indefinitely. We link PT-symmetry breaking to the long-time behavior of QCs, which display different scalings in the PT-broken/unbroken phase and at the exceptional point (EP). The EP in particular stands out as the most classical configuration.

14:20 : Invited talk

On the connection between parity-time symmetry and time-variant wave media

Theodoros T. Koutserimpas, Romain Fleury

Swiss Federal Institute of Technology Lausanne (Switzerland)

Parity-time symmetry can allow non-Hermitian Hamiltonians to exhibit real eigenvalues. This interesting condition of symmetries has enabled numerous studies in the fields of quantum mechanics and optics. Such theoretical and experimental studies involve a combination of material gain and loss. In this talk, we connect the phenomena that are related to parity-time symmetry with the wave phenomena that are related to temporal changes of the wave parameters of the medium, by introducing a time-transitioning matrix.

14:40 : Invited talk

Nonlinear mode competition and symmetric protected power oscillations in topological lasers

Simon Malzard¹, Henning Schomerus²

¹Imperial College London (United Kingdom), ²Lancaster University (United Kingdom)

Topological lasing in non Hermitian systems offers a new interpretations of topological states as they can be selectively amplified by distributed gain and loss. A key question is whether such topological mode selection persists when nonlinearities that stabilise these systems at their working point are included. Here we show that topological defect lasers can operate in genuinely topological states with analogues of zero modes from the linear setting, as well as a novel states displaying symmetry protected power oscillations.

15:00 : Invited talk

Emergent PT symmetry and quantum fluctuations in a double-quantum-dot circuit QED set-up

Archak Purkayastha¹, Manas Kulkarni², Yogesh Joglekar³

¹The University of Dublin (Ireland), ²Tata Institute of Fundamental Research (India), ³Indiana University Purdue University Indianapolis (IUPUI) (USA)

Open classical and quantum systems with effective parity-time (PT) symmetry have shown tremendous promise for advances in lasers, sensing, and non-reciprocal devices. However, the microscopic origin of such effective, non-Hermitian models is not well understood. In this work, by microscopically modelling a double-quantum-dot-circuit-QED set-up that is realizable in state-of-the-art experiments, we show that a non-Hermitian Hamiltonian emerges naturally, which can be controllably tuned to observe both PT-transition, as well as the effect of quantum fluctuations.

14:00 - 15:20 — Augustin Fresnel Room

Session 2A24

Symposium IV: Chirality, magnetism, and magnetoelectricity: Separate phenomena and joint effects in metamaterial structures

Organized by: Eugene Kamenetskii

Chaired by: Eugene Kamenetskii

14:00 : Invited talk

The influence of the internal domain wall structure and chirality on spin wave dynamics in periodic magnetic stripe domain patterns

Pawel Gruszecki¹, Jan Kisielewski², Andrzej Maziewski², Maciej Krawczyk¹

¹Adam Mickiewicz University (Poland), ²University of Bialystok (Poland)

The study of the Dzyaloshinskii-Moriya interaction together with perpendicular magnetocrystalline anisotropy and film's thickness influence on both the static magnetic configuration and the dynamics of spin waves in periodic aligned stripe domain patterns is provided. We analyze the sensitivity of excitation of resonant modes on polarization of the microwave field. Subsequently, we demonstrate in micromagnetic simulations the unidirectionality of spin-wave propagation along the domain walls and define the conditions for its occurrence.

14:20 : Invited talk

Synthetic chiral light for efficient control of chiral light matter interaction

David Ayuso, Ofer Neufeld, Andres Felipe Ordonez, Piero Decleva, Gavriel Lerner, Oren Cohen, Misha Ivanov, Olga Smirnova

Max-Born-Institut Berlin (Germany)

I will talk about synthetic chiral light, a new type of freely propagating optical fields that we have recently introduced. Synthetic chiral light enables the highest possible degree of control over chiral light-matter interaction. It allows us to drive a strong nonlinear response in chiral molecules of a selected handedness. As a result, we can force a particular molecular enantiomer to emit bright harmonic light while its mirror twin remains dark, reaching the ultimate efficiency limit in chiral discrimination.

14:40 : Invited talk

Superparamagnetic nanoparticles with deep UV plasmonic response

Vladimir P. Drachev¹, Ali Aliev²

¹Skolkovo Institute of Science and Technology (Russia), ²University of Texas at Dallas (USA)

Structural and magnetic characterizations prove that the single-domain and superparamagnetic properties of nanoparticles required for spin dependent plasmon oscillations. Good quality plasmon resonance in absorption is proven to be a property of isolated Co nanoparticles. The magnetic field induced aggregation of nanoparticles results in a suppression of the resonance quality. The measured magnetic moment above blocking temperature is about 0.69 of Bohr magnetons per atom in a 8.8 nm nanoparticle, which is much lower than in bulk cobalt, 1.73.

15:00 : Invited talk

Magnon-polaritons condensates with superfluidity and vortex formation

Eugene Kamenetskii

Ben Gurion University of the Negev (Israel)

We show that magnon-polaritons can be realized due to magnon condensation caused by magnetic dipole-dipole interaction. We study quantized vortices in magnon-polariton condensates arising from magnetic-dipolar-mode (MDM) oscillations in a quasi-2D ferrite disk placed in a microwave cavity. We show that it is possible to trap a magnon Bose-Einstein condensate in a ring geometry and induce rotational superflow in this system. We consider torque transfer induced by MDM oscillations.

14:00 - 15:20 — Ibn Al-Haytham Room

Session 2A25

Exotic Meta-media - Time-dependent, Nonlocal and Other Novel Responses

Organized by: Martin McCall, Jonathan Gratus and Paul Kinsler

Chaired by: Martin McCall, Jonathan Gratus and Paul Kinsler

14:00 : Invited talk

Quantum spill-out induced enhancement in surface nonlinear plasmonic response

Muhammad Khalid, Cristian Ciraci

Italian Institute of Technology (Italy)

We develop a theoretical model based on the quantum hydrodynamic description of free-electrons and present its numerical implementation to investigate surface second-order nonlinearities. The presented method is capable of dealing with realistic profiles of equilibrium density of a metal. In the spectral analysis of Na and Ag thick slabs, we observe strong resonances induced by the electron spill-out from the metal surface. These resonances can be exploited to enhance the second-harmonic conversion efficiency by several orders of magnitude.

14:20 : Invited talk

A (re-)introduction to spatial dispersion

Paul Kinsler

Lancaster University (United Kingdom)

I examine the traditional approach to spatial dispersion, and contrast this with an alternative based on a fundamental re-examination of the basic principles. I emphasise the key role of the spatial properties which generate spatially dispersive behaviour: geometry, structure, and dynamics.

14:40 : Invited talk

D and H cannot exist: Axions, topology and global charge conservation

Jonathan Gratus¹, Paul Kinsler¹, Martin W McCall²

¹Lancaster University (United Kingdom), ²Imperial College London (United Kingdom)

The excitation fields D and H cannot be directly measured and have a gauge freedom. Admitting this freedom opens many possibilities, concerning the axionic response, which have direct application. We show a simple scenario which due to topological reasons is impossible using D and H. It can be used to model periodic lattices.

ces which have non zero total charge and show that an evaporated black hold can break charge conservation. We reflect on the nature of a homogeneous axionic response.

15:00 : Invited talk

Surface waves with mixed exponential and linear localization characteristics

Tom Mackay¹, Chenzhang Zhou², Akhlesh Lakhtakia²

¹University of Edinburgh (United Kingdom), ²Pennsylvania State University (USA)

A theory underpinning new types of surface waves, guided by the planar interface of an anisotropic material and an isotropic material, has been developed. These new surface waves propagate only in one direction for each quadrant of the interface plane and their decay in the direction normal to the interface in the anisotropic partnering material is given by the product of an exponential and a linear function of distance from the interface plane.

14:00 - 15:15 — Gaston Floquet Room

Session 2A26

Parity-Time and quasi-normal modes in Photonics, Plasmonics, Acoustics

Organized by: Anatole Lupu and Henri Benisty

Chaired by: Anatole Lupu and Henri Benisty

14:00 : Invited talk

Nonorthogonality constraints in open quantum systems

Jan Wiersig

Otto-von-Guericke-Universitat (Germany)

The nonorthogonality of quasi-normal modes plays an important role in the physics of non-Hermitian systems. We demonstrate that the known nonorthogonality bound for effective Hamiltonians describing decaying systems may not be valid in quantum and wave systems with radiation due to quantum backflow. A geometric interpretation of the nonorthogonality bound is given which reveals that in this context the complex energy (or frequency) space can be seen as a surface of constant negative curvature.

14:20 : High-Q modes, quasi-modes, and pseudo-modes in dielectric nanoparticles of different shapes

Vasily V. Klimov

Lebedev Physical Institute (Russia)

Optical properties of dielectric nanoparticles of different shapes are investigated. Continuous distributions of Q factor as a function of frequency and permittivity are found, and modes, quasi-modes, and pseudo-modes of NP are extracted. The pseudo-modes are characterized by small stored energy and scattered power. Our results show that among nanoparticles of different shapes but of the same volume nanospheres have the highest Q factor in the limit of high permittivity.

14:35 : Invited talk

Optical mode transfer by encircling fixed and moving exceptional points

Pierre Berini

University of Ottawa (Canada)

We discuss recent progress on optical mode transfer in parity-time symmetric systems. We discuss structures where modes evolve in parameter space such that an exceptional point is encircled, and consider cases where the exceptional point is fixed or mobile. We also discuss prospects for realizing non-linear or active structures.

14:55 : Invited talk

Nanoantennas with Balanced Gain and Loss

Alejandro Manjavacas

University of New Mexico (USA)

We will discuss how the addition of plasmonic elements with gain to conventional metallic nanoantennas can

serve to achieve directional optical responses.

14:00 - 15:20 — Lawrence Bragg Room

Session 2A27

Machine learning for metamaterials and metasurfaces

Organized by: Mohamed Bakr and Willie Padilla

Chaired by: Mohamed Bakr and Willie Padilla

14:00 : **Invited talk**

Deep-Neural-Network Enabled Metasurface Designs

Sensong An, Clayton Fowler, Bowen Zheng, Hong Tang, Hang Li, Hualiang Zhang

University of Massachusetts Lowell (USA)

Metasurfaces are being widely investigated and adopted for their potential for integrating multiple functionalities into a single, flat optical device. A key challenge in this field is the non-intuitive design process that produces designs based on specific electromagnetic requirements. Meanwhile, deep neural network (DNN) has been proven to be an effective solution to non-intuitive design tasks. In this paper, we detail a novel approach to design metasurfaces using DNNs and demonstrate some devices achieved based on this approach.

14:20 : **Invited talk**

Deep learning the electromagnetic properties of metamaterials with neural networks

Omar Khatib, Simiao Ren, Jordan Malof, Willie Padilla

Duke University (USA)

We demonstrate use of machine learning to explore the fundamental electromagnetic properties of all-dielectric metasurfaces. Our neural network architecture enables a machine learning based approach for discovering new physical insights from data - automating the traditional ways by which physicists develop models of nature from experimental and numerical observations. Our approach is Kramers-Kronig compliant, and is not limited to the case demonstrated, but is compatible with many characterization techniques including Fourier transform spectroscopy, ellipsometry, and terahertz time domain spectroscopy.

14:40 : **Invited talk**

Accelerating the Design of Photonic Metamaterials by Artificial Intelligence

Yongmin Liu

Northeastern University (USA)

In this talk, I will discuss how to accelerate the design of novel metamaterials by deep learning, a subset of artificial intelligence (AI) that learns multilevel abstraction of data using hierarchically structured layers. Different from the conventional approaches, deep learning can produce fast and accurate designs without the need of case-by-case and time-consuming numerical calculations. The results show many exciting opportunities in the areas of optical design, integration and measurements when interfacing photonics with deep learning.

15:00 : **Invited talk**

Symphotonic Metamaterials for Photonic Computing and Machine Learning Applications

David R. Smith, Patrick Bowen, Daniel Marks, Roberto Zecca, Divya Pande

Duke University (USA)

Symphotonic metamaterials are volumetric metamaterials that multiplex large numbers of optical functions with high efficiency. Provided enough functionality can be integrated into a symphotonic device, a variety of computing algorithms can be implemented photonically, vastly reducing latency and with orders of magnitude less energy consumption. Here we introduce the symphotonic concept and describe a device that can compute a convolution optically—a critical step in many neuromorphic architectures.

14:00 - 15:20 — Rene Descartes Room**Session 2A28****Plasmonics for single molecule detection and manipulation**

Organized by: Denis Garoli

Chaired by: Denis Garoli

14:00 : Invited talk**Surface-enhanced Raman scattering signal intensity as a function of the plasmon resonance and the excitation wavelength****Sylwester Gawinkowski***Polish Academy of Sciences (Poland)*

We report on single-molecule surface-enhanced Raman spectroscopy of porphyrins, porphycenes, and phthalocyanines. Using the tunable wavelength laser and tuning it around the local plasmon resonance and electronic resonance of the molecule, we are showing the influence of that parameter on the single-molecule SERS signal strength and its detectability. The results show that steric factors play a crucial role in the detectability of single-molecules.

14:20 : Invited talk**Single-Molecule SERS Detection of DNA Bases in Oligonucleotides by Controllable Electro-plasmonic Trapping of Single Nanoparticles in Plasmonic Nanoholes****Jianan Huang, Mansoureh Z. Mousavi, Yingqi Zhao, Aliaksandr Hubarevich, Fatima Omeis, Giorgia Giovannini, Denis Garoli, Francesco De Angelis***Italian Institute of Technology (Italy)*

DNA molecules flow through nanopores too fast to allow Surface-enhanced Raman spectroscopy (SERS) detection of single bases. Here, we report an approach to control the residence time of molecules in a hot spot by an electro-plasmonic trapping effect. By directly adsorbing molecules onto a gold nanoparticle and trapping the single nanoparticle in a plasmonic nanohole up to minutes, we demonstrate single-molecule SERS detection of all four DNA bases as well as discrimination of single nucleobases in a single oligonucleotide.

14:40 : Invited talk**Transferred Nanopores Array from Nanosphere Lithography onto a Holey Membrane for Biosensing Applications****Marzia Iarossi¹, Daniel Darvill², Jian-An Huang², Ricardo M. Abraham Ekereth³, Francesco De Angelis²***¹Universita degli Studi di Genova (Italy), ²Istituto Italiano di Tecnologia (Italy), ³Universidad Nacional del Centro de la Provincia de Buenos Aires (Argentina)*

We report on an easy and low-cost strategy that combines the fabrication of an ordered array of nanopores by performing thermal annealing on a hexagonally close-packed monolayer of nanospheres and the transfer of the colloidal array onto a holey membrane. Adjusting the duration of the annealing process it is possible to control both size and shape of the interstices in the nanospheres arrays. Both optical and electrical measurements have been performed to evaluate the properties of the nanopores array.

15:00 : Invited talk**Time- and field-Resolved Response of Plasmonic Nanostructures and Their Applications to Single-Molecule Detection and Manipulation****Nicolo Maccaferri***University of Luxembourg (Luxembourg)*

Plasmonics enables a plenty of novel optical effects and functionalities. In the first part of the talk we will show that, by combining femtosecond time-domain spectroscopy and high-resolution confocal microscopy, it is possible to measure full time- and field-resolved response of single plasmonic nanoantennas. In the second part of the talk, we will show practical applications of plasmonic nanostructures to single-molecule detection, enhanced spectroscopies, nanoparticle trapping and beaming, and resonant energy transfer.

Coffee Break and Exhibit Inspection

Session 2P2

Poster session IV

15:20 - 16:00

P1: Simulation of Large Metasurfaces through Transfer Function Mask**Chenglin Xu, Mayank Bahl, Evan Heller***Synopsys, Inc. (USA)*

A more efficient and accurate approach is demonstrated to simulate metasurfaces. The new approach employs FFT-BPM to propagate the transmitted field through a transfer function mask, which is formed by using rigorous FDTD or RCWA algorithm on local nano-cells. Larger grid and step sizes can be used and the new approach is more efficient and requires less RAM, hence it can be applied to larger metasurfaces. Validations against FDTD and FD-BPM shows it remains accurate, within a reasonable approximation.

P2: Strongly Extended Many-Body Enhancement in Diamond Epsilon Near-Zero Metamaterials**Olivia L. Mello¹, Yang Li², Philip Camayd-Munoz³, Marki Loncar¹, Eric Mazur¹***¹Harvard University (USA), ²Tsinghua University (China), ³California Institute of Technology (USA)*

We demonstrate analytically and numerically that with a diamond epsilon near-zero (ENZ) metamaterial with design we experience an ultra-high cooperative enhancement over distances greater than 10 microns for both two emitters and many-body ensembles of dipoles.

P3: Dynamic plasmonic nanorod pixels**Nicholas J. Greybush¹, Kristin Charipar¹, Paul Johns¹, Stephen J. Bauman², Dennis Doyle³, Jeffrey A. Geldmeier¹, Jawad Naciri¹, Nicholas Charipar¹, Jake Fontana¹***¹U.S. Naval Research Laboratory (USA), ²University of Arkansas Fayetteville (USA), ³University of Pittsburgh (USA)*

Plasmonic metamaterials present a new paradigm for color generation, but their properties are often either static in time or suffer from slow switching speeds. However, by reversibly aligning colloidal plasmonic nanorods using electric fields, we demonstrate rapid modulation of light on microsecond time scales. Tailoring nanorod aspect ratio and composition facilitates operation across the visible through short-wave-infrared spectra. We characterize plasmonic nanorod pixels' chromaticity and luminance, and showcase spatial, spectral, and temporal control of light in exemplary display devices.

P4: Metamaterial-assisted Inductive Power Transfer using Transmission-Line Mode**Jorge V. de Almeida¹, Eduardo Costa da Silva¹, Marbey Manhaes Mosso¹, Carlos A. F. Sartori²***¹Pontifical Catholic University of Rio de Janeiro (Brazil), ²Polytechnic School of USP (Brazil)*

In this paper, a metamaterial-assisted coil for inductive power transfer system is presented. The proposed prototype exploits the transmission-line mode instead of the usual antenna one in order to minimize radiation at the resonance. By judiciously choosing the operating mode of the circuit's driver, the driver losses are reduced and the focusing mechanism of the metamaterial-based lens is improved. The theoretical results are supported by analytical and numerical evidence.

P5: Design Strategies for Highly Integrated and Efficient Metasurfaces enabling advanced Terahertz Imaging**Thomas S. Nowack, Yash D. Shah, Mitchell G. Kenney, Ivonne Escorcía Carranza, James Grant, David Cumming***University of Glasgow (United Kingdom)*

To advance Terahertz imaging towards its unique applications, drastic improvement of the optical setups in terms of efficiency and number of required components is imperative. We present our approach to design large-area, all-dielectric metasurfaces for ultra-compact, efficient imaging optics at THz frequencies, which significantly reduce reflective and alignment losses by multiplexing and superposition of optical responses. We exemplify this with two metalenses based on geometric phase that exhibit two multiplexed focal points for orthogonal polarizations and a superimposed Q-plate.

P6: Gain Properties of Axially Slotted Antenna Coated by Inhomogeneous Materials Embedded in Different Materials and Metamaterials**Saddia Ilyas¹, M. Farooq Nasir¹, Shakeel Ahmed²**¹Riphah International University (Pakistan), ²Quaid-i-Azam University (Pakistan)

The properties of axially slotted cylindrical antenna coated with inhomogeneous material placed in various metamaterials have been studied. The semi analytical technique is employed to reveal the effects of different coating layers and host medium on the gain properties of the slotted antenna. The relationship of coating thickness, inhomogeneity profile and nature of the host medium with characteristics of the slotted antenna have been discussed.

P7: Single-molecule and single-nanoparticle spectroscopy of porphycene derivatives**Abdolvahab Amirjalali¹, Katarzyna Tarnowska¹, Yoshio Hisaeda², Sylwester Gawinkowski¹**¹Polish Academy of Sciences (Poland), ²Kyushu University (Japan)

Here we have studied the SERS signal intensity as a function of concentration, temperature, and excitation wavelength for a series of porphycene derivatives. The SERS spectra were averaged over a large area of uniformly distributed gold nanorods deposited on the glass substrate. In the case of the very low concentration samples also temporal evolution of SERS signal on hot-spots was registered. Temporal evolution combined with bianaalytical method proved single-molecule detection.

P8: Effect of deposition angle on fabrication of plasmonic metal nanocones**Jiri Liska¹, Filip Ligmajer¹, Pedro V. Pinho², Lukas Kejik¹, Michal Kvapil¹, Petr Dvorak¹, Nikolaus S. Leitner³, Erik Reimhult³, Tomas Sikola¹**¹Brno University of Technology (Czech Republic), ²University of Campinas (Brazil), ³University of Natural Resources and Life Sciences (Austria)

Metal nanocones can be used to enhance light-matter interactions or for location-specific plasmonic sensing. Their fabrication often utilizes self-shading effect, which occurs during metal film evaporation into circular nanowells. We present a view on the fabrication of ordered arrays of gold nanocones using electron-beam lithography/evaporation. Lateral position of the substrate during the evaporation influences the symmetry of the fabricated nanocones and that off-axis deposition forms asymmetric structures. Our findings help to identify limits for production of wafer-scale arrays.

P9: Agile Metamaterial Absorber at W-bande**Khedidja Bouras, Abdelhadi Labiad, Mouloud Bouzouad***Laghouat University (Algeria)*

In this work we propose an SRR type absorption cell with a negative refractive index which exhibit both negative permittivity and permeability around its resonance. The agile SRR is circular shape with four branches which has switches installed. The proposed structure is a matrix 3 X 3 unit cell controllable with an external system like MEMS, where it subject boundary condition. We get the agility feature if we change the type of one or many cells in the matrix structure.

P10: Nonlinear Properties of Magnetic Metasurfaces**Anastasia Bir, Dmitriy Romanenko, Sergey Odintsov, Sergey Nikitov, Sergey Grishin***Saratov State University (Russia)*

In a paper, the experimental results of the amplitude-frequency and power characteristics of two-dimensional ferromagnetic structures are demonstrated in microwave range. It is established that the presence of a periodic structure makes it possible to control the power thresholds of nonlinear processes of spin waves.

P11: Monochromatic THz radiation of relativistic electrons from a metasurface**Daria Sergeeva¹, Alexey Tishchenko¹, Alexander Aryshev²**¹National Research Nuclear University "MEPhI"(Russia), ²KEK: High Energy Accelerator Research Organization (Japan)

We report on experimental and theoretical studies of coherent THz Smith-Purcell radiation and grating transition radiation generated by an electron bunch from a metasurface. The results are compared, and agreement is found to be marvelous for spectra, while angular distributions show discrepancy. We show that, qualitatively, the difference in angular distributions can be caused by contribution of such processes as coupling between particles or excitation of surface plasmon-polaritons.

P12: Integrated All Optical Passive Neural Network Using Silicon Metalines**Sanaz Zarei, Mahmood R. Marzban, Amin Khavasi***Sharif University of Technology (Iran)*

We propose a fully optical architecture for implementing deep neural network using nano-photonics integrated circuits. Fully optical matrix multiplications are performed using silicon-based metalines. The proposed whole-passive optical neural network is very compact and works at the speed of light, with less energy consumption than state of the art electronic counterparts. Various complex functions that is performed by digital neural networks can be implemented by our proposal at the wavelength of $1.55\mu\text{m}$.

P13: Three-Dimensional Photonic Crystal Composites with High Refractive Index Thin films**Mike P. C. Taverne¹, Xu Zheng¹, Yushao Chen¹, Katrina A. Morgan², Lifeng Chen¹, Ghadah Alzaidy², Chung-Che Huang², Ying -Lung Daniel Ho¹, Daniel W. Hewak², John G. Rarity¹**¹ *University of Bristol (United Kingdom)*, ² *University of Southampton (United Kingdom)*

We study polymer photonic crystals coated with varying thickness of high refractive index material aiming to make functional photonic devices capable of controlling light through band structure and dispersion. We observed red shifts of partial bandgaps in the near infrared region when the thickness of deposited MoS₂ films increases. A ~ 150 nm red shift of the fundamental and high order bandgaps is measured after a ~ 15 nm thick MoS₂ coating.

P14: Enhancement of Antenna Radiation through Control of the Local Environment**Leanne D. Stanfield, Alastair Hibbins, Roy Sambles***University of Exeter (United Kingdom)*

The radiated power of a dipole antenna off resonance can be enhanced by employing an iterative design process derived from the Purcell effect. The local electromagnetic environment about an emitter can be engineered by controlling the local physical environment. For a 1 cm dipole antenna radiating at 2 GHz, embedded in a dielectric structure, radiated power enhancement rates of up to 5280 times have been predicted by computational models.

P15: Anisotropic Electromagnetic Wave Propagation in Two Dimensional Ferromagnetic Metamaterials**Maria Amelchenko¹, Feodor Ogrin², Aleksey Titov¹, Sergey Grishin¹**¹ *Saratov State University (Russia)*, ² *University of Exeter (United Kingdom)*

It is well known that metamaterials are artificial structures providing electromagnetic properties that natural materials do not have. There is a class of metamaterials called ferromagnetic metamaterials possessing a negative permeability in microwave range and a permittivity that can change its sign in terahertz range. In this paper, both the anisotropic properties and dispersion characteristics of fast and slow electromagnetic waves (EMWs) propagating in two-dimensional ferromagnetic metamaterials (metasurfaces) with finite wire conductivity are studied.

P16: Chemically Modulated Hyperbolic Metamaterials**Jose L. Ocana-Pujol, Ralph Spolenak, Henning Galinski***ETH Zurich (Switzerland)*

Hyperbolic Metamaterials are multilayered optical nanomaterials that exhibit, due to their intrinsic anisotropy, optical topological transitions. Here, we propose and experimentally demonstrate an all zirconium-based hyperbolic metamaterial. We have selected ZrN/ZrO₂ as specific example, as these earth-abundant refractory materials offer unique mechanical and thermal stability in extreme environments, such as high temperatures above 2000°C. Alternating the reactive gases, oxygen and nitrogen during reactive sputtering, enables continuous growth of a chemically modulated nanomaterial.

P17: Phononic band gap of longitudinal acoustic waves in one-dimensional crystal for ultrasonic applications**Ahmed Kahlouche, Mounir Bouras, Nassim Dermeche, Abdesselam Hocini***University of Mohamed Boudiaf of M'sila (Algeria)*

Nowadays, sensor technology has attracted great interest in various domains. In this work, we have analyzed phononic band gaps of one-dimensional phononic crystal made by a stack of N bi-layers of LiNbO₃/SiO₂. The transmission spectrum of acoustic waves is calculated by using the transfer matrix method (TMM). The results clearly demonstrate the existence of phononic band gap of which the position and the width are strongly

affected by many physical parameters. Our results are useful in various applications such as acoustic barriers and sensor materials

P18: Toward investigation of reactive intermediates via Surface-enhanced Raman Spectroscopy (SERS)
Łukasz Gutowski¹, Malwina Liszewska¹, Bartosz Bartosewicz¹, Bogusław Budner¹, Jan L. Weyher², Bartłomiej J. Jankiewicz¹

¹Military University of Technology (Poland), ²Polish Academy of Sciences (Poland)

The Surface-enhanced Raman Spectroscopy (SERS) has been used to investigate the possibility of aryl monoradicals generation from thiophenols and phenylmethanethiols substituted with iodine or bromine atoms. The monolayers of radical precursors were deposited on SERS substrates, which were then immersed in methanol and irradiated for six hours with a UVC lamp. Pre- and post-reaction SERS spectra were obtained by using a portable Raman spectrometer and compared with the spectra of expected products of radicals reaction with methanol.

16:00 - 17:10 — Victor Veselago Room

Session 2A29

Plenary Session II

16:00 : Plenary talk

How Light Behaves when the Refractive Index Vanishes

Robert W. Boyd

University of Ottawa (Canada)

We describe some of the properties of light propagation through material for which the dielectric permittivity and hence the refractive index is nearly vanishing. Among other unusual optical properties, we find that such epsilon-nearzero (ENZ) materials display an extremely large nonlinear optical response, with important implications for the field of photonics.

16:35 : Plenary talk

Empowering Quantum Photonics with Nanoplasmonics and Machine Learning

Vladimir Shalaev

Purdue University (USA)

New approaches to address major challenges in quantum photonics by employing powerful ideas and concepts developed in the field of plasmonic metamaterials will be discussed.

Break

17:10 - 17:30

17:30 - 18:50 — Victor Veselago Room

Session 2A30

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

17:30 : Keynote talk

The Challenge of META is (Aperiodic) Inverse Electromagnetic Design

Eli Yablonovitch

UC Berkeley (USA)

Meta-Materials were inspired by the idea of periodic sub-wavelength geometrical units that produce an effective permittivity and permeability. Inevitably, in human affairs, there is a goal, and it may be that the goal is best achieved by an aperiodic rather than a periodic design. This is a universal problem in electromagnetics: What is the best design to achieve a specific goal? I will introduce the "Adjoint Method", a combination of calculus and linear algebra that generates optimal electromagnetic designs.

18:00 : Keynote talk

Wavelength conversion through plasmonic photoconductive nanostructures

Mona Jarrah

University of California Los Angeles (USA)

In this presentation, I will briefly overview the physics and possible macroporous, inch-scale nanostructures with submicron lattice parameter and discuss strategies for potential extension of more complex and functionally optimized hierarchical nanostructures.

18:30 : Invited talk

Nonlinear and electro-optic nanostructured metal-oxides

Felix Richter, Viola Valentina Vogler-Neuling, Helena Weigand, Flavia Timpu, Artemios Karvounis, Rachel Grange

ETH Zurich (Switzerland)

Non-centrosymmetric metal-oxide media possess a plethora of functional optoelectronic properties like second harmonic generation, electro-optic and electro-mechanical response. We present a variety of approaches to exploit these mechanisms at the nanoscale with the use of all-dielectric metasurfaces and photonic crystals based on barium titanate as well as plasmonic metasurfaces based on gold-barium titanate nanostructures.

17:30 - 18:50 — Allan Boardman Room

Session 2A31

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

17:30 : Invited talk

Enhancing second-harmonic generation using dipolar-parity modes in non-planar plasmonic nanocavities

Feng Wang, Hayk Harutyunyan

Emory University (USA)

Efficient optical nonlinear sources at the nanoscale are important for the development of photonic circuitry, quantum optics, and biosensing. In this work, we use metal-insulator-metal nanocavities to demonstrate strong second harmonic generation in nanoscale plasmonic cavities. We show that in contrary to the common belief, double-resonant cavities can be designed with dipolar-type modes for both fundamental and emission resonances.

17:50 : Invited talk

Theory of hot-carrier generation and relaxation in plasmonic nanoparticles

Yu Zhang

Los Alamos National Laboratory (USA)

A quantum-mechanical model for hot-carrier generation, relaxation, and extraction is developed. The plasmon excitation and hot-carrier generation from the plasmon decay are derived from the linear-response time-dependent density functional theory. And its connection to the semiclassical model is discussed. Relaxation due to electron-electron and electron-phonon scatterings are treated on equal footing. With this development,

the initial distribution of hot-carriers and lifetimes of hot-carriers induced by different excitation are investigated.

18:10 : Invited talk

Manipulating Plasmonic Excitonic Nanomaterials with Coherent Phonons

Matthew S. Kirschner, Wendu Ding, Craig T. Chapman, Xiao-Min Lin, Lin X. Chen, George C. Schatz, Richard D. Schaller

Northwestern University (USA)

Coherent acoustic phonons can modulate electronic interactions with proximal excitonic molecular species. Gold bipyramids with systematically altered aspect ratios and corresponding localized surface plasmon resonance energies, functionalized with an excitonic, J-aggregated dye molecule, produce two hybridized states that exhibit anti-crossing behavior with a Rabi splitting energy of 120 meV. Such oscillatory plasmonic-excitonic nanomaterials offer a route to manipulate and dynamically-tune the interactions of plasmonic/excitonic systems and unlock a range of potential applications.

18:30 : Invited talk

Plasmon-Induced Resonance Energy Transfer for Photocatalysis, Biosensing and Photodynamic Therapy

Nianqiang Wu

University of Massachusetts Amherst (USA)

This talk presents a new energy transfer mechanism from metal to semiconductor, that is, plasmon-induced resonance energy transfer (PIRET). This talk will demonstrate the use of PIRET mechanism to design new materials and devices for solar energy conversion, photocatalysis, biosensing and photodynamic therapy.

17:30 - 19:05 — Tatsuo Itho Room

Session 2A32

Advanced modeling techniques for the design of metasurface devices

Organized by: Patrice Genevet and Stéphane Lanteri

Chaired by: Patrice Genevet and Stéphane Lanteri

17:30 : Invited talk

T-operator limits on electromagnetic power transfer: nanophotonic upper bounds on far-field cross sections and Purcell enhancement

Sean Molesky, Prashanth S. Venkataram, Weiliang Jin, Pengning Chao, Alejandro Rodriguez

Princeton University (USA)

We present a scheme based on the scattering T operator to set physical bounds on any single electromagnetic design problem that can be framed as a net power process. The technique is found to predictively quantify and differentiate the relative performance of dielectric and metallic devices for both far and near-field sources (e.g. Purcell enhancement). The resulting framework, which relies on Lagrange duality, has immediate applications for guiding, understanding, and improving inverse design.

17:50 : Invited talk

Optimization Paradigms for Metasurface Inverse-design

Eric B. Whiting, Sawyer D. Campbell, Ronald P. Jenkins, Pingjuan L. Werner, Douglas H. Werner

The Pennsylvania State University (USA)

Metasurfaces with high efficiency and broadband performance hold the potential to revolutionize optical system design. However, advanced optimization methods and design techniques must be exploited to achieve performances that surpass conventional optical systems. In this paper, two metasurface design strategies are presented and their advantages and challenges discussed. The first strategy is based on topology optimization while the second is based on deep learning. finally, we discuss how hybridizing these techniques could surpass the state-of-the-art for metasurface optimization.

18:10 : Invited talk**Metasurface optimization based on coupled mode theory****Zongfu Yu***University of Wisconsin Madison (USA)*

We develop a coupled mode theory to model metasurfaces. It is much faster than full-wave solvers. Combined with the adjoint method, we show that the CMT approach can be used to efficiently design complex metasurfaces for a variety of functions. As an example, we also demonstrate thermal holograms based on metasurfaces.

18:30 : Invited talk**Bounds on achievable performance via Lagrange duality****Guillermo Angeris, Jelena Vuckovic, Stephen Boyd***Stanford University (USA)*

While the problem of designing physical devices is computationally hard, heuristics such as inverse design are often successful in finding designs with very good performance. A natural question to ask is: how does the performance of these designs compare to the best possible performance? We show how to construct a convex (and therefore easily solvable) problem whose optimal value bounds the best achievable performance, for a given objective. We also provide a proof of the NP-hardness of physical design.

18:50 : An efficient global optimization technique based on statistical learning for highly efficient metasurface designs at visible regime**Mahmoud M. R. Elsayw, Patrice Genevet, Régis Duvigneau, Stéphane Lanteri***Université Côte d'Azur (France)*

The object of this contribution is twofold. first, we present a brief review of the common inverse design techniques in the field of metasurfaces. Second, we present a global optimization technique based on statistical learning for optimizing highly efficient metasurface devices. Using our method, a light deflection metasurface with efficiency reaching 85 % for both TM and TE polarizations has been realized at the visible regime.

17:30 - 19:05 — Christian Huygens Room**Session 2A33****Non-Hermitian Photonics: From Plasmonics, and Metamaterials, to Topological and Quantum systems**

Organized by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

Chaired by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

17:30 : Invited talk**Encircling exceptional points as a non-Hermitian extension of rapid adiabatic passage****Alexander Schumer¹, Juraj Feilhauer², Jorg Doppler¹, Alexei A. Mailybaev³, Julian Bohm⁴, Ulrich Kuhl⁴, Nimrod Moiseyev⁵, Stefan Rotter¹**

¹Vienna University of Technology (Austria), ²Slovak Academy of Sciences (Slovak Republic), ³Instituto Nacional de Matematica Pura e Aplicada - IMPA (Brazil), ⁴Université côte d'Azur (France), ⁵Technion-Israel Institute of Technology (Israel)

Coherent transfer of population between different energy levels is a key task in quantum mechanical applications with rapid adiabatic passage (RAP) being one of the most prominent protocols. By reframing the underlying process in the domain of non-Hermitian physics through adding a variable amount of loss we theoretically and experimentally disclose a fundamental connection between the symmetric state flip of RAP and the asymmetric state transfer when encircling an exceptional point (EP).

17:50 : Correlations of indistinguishable photons in quantum walks under broken PT-symmetry**Friederike U. J. Klauck¹, Markus Grafe², Matthias Heinrich¹, Alexander Szameit¹**

¹University of Rostock (Germany), ²Fraunhofer-Institut für Angewandte Optik und Feinmechanik (Germany)

We experimentally study the correlations of indistinguishable photons in a one-dimensional Parity-Time (PT) symmetric quantum walk. Introducing a symmetric loss distribution in a homogeneous waveguide system gives rise to PT-symmetry, which drastically alters the correlations of indistinguishable photons, giving rise to e.g. correlations similar to distinguishable classical particles in a random walk.

18:05 : Invited talk

Optical thermodynamics in topological Haldane lattices

Pawel Jung¹, Midya Parto¹, Fan Wu¹, Yuzhou Liu², Mercedeh Khajavikhan², Demetrios Christodoulides¹

¹University of Central Florida (USA), ²University of Southern California (USA)

In this work we show that a nonlinear topological Haldane lattice exhibits a number of interesting optical thermodynamic properties such a heat capacity function with the Schottky anomaly effect or a metastable responses leading to different final temperatures and chemical potentials at equilibrium condition.

18:25 : Invited talk

Topological band theory for dissipative quantum and classical systems

Simon Lieu

University of Maryland (USA)

I will present a topological classification applicable to dissipative systems governed by a general class of Lindblad master equations. The dynamical matrix governing the equations of motion for "quadratic Lindbladians" must belong to one of ten non-Hermitian Bernard-LeClair symmetry classes which reduce to the Altland-Zirnbauer classes in the closed limit. These symmetries protect edge modes in a dissipative environment. I will discuss current efforts to extend the formalism to bosonic systems, with relevant implications for topological photonics.

18:45 : Invited talk

Non-Hermitian Constant Intensity Waves

Ziad H. Musslimani

Florida State University (USA)

In this talk, we will discuss recent progress related to linear and nonlinear wave propagation in non Hermitian media. We shall show that under certain circumstances, a constant intensity waves can propagate in non-homogeneous media with imposed parity-time symmetry. Such waves can even exist in a complex non-hermitian random media.

17:30 - 19:00 — Augustin Fresnel Room

Session 2A34

Light-matter interactions in new materials and meta-architectures

Organized by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

Chaired by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

17:30 : Invited talk

Gyrotropic responses mediated by Jahn-Teller and spin-orbit interactions

Blai Casals, Gervasi Herranz

Institute for Materials Science of Barcelona ICMA-B-CSIC (Spain)

We report on gyrotropic responses mediated by Jahn-Teller polarons and spin-orbit coupling, whereby a spin-reversal is induced by photons and can be selectively detected by circularly polarized light. We show that spin-reversal requires the concurrent action of orthorhombic and tetragonal Jahn-Teller modes, which opens up a pathway to explore quantum effects that involve spin and orbital degrees of freedom. Additionally, our results provide an original, general pathway towards the generation of magnetic-responsive gyrotropic responses.

17:50 : Invited talk

Conformable metasurfaces for linear and nonlinear applications**Jialing Xiao, Usenobong Akpan, Sebastian A. Schulz, Andrea Di Falco***University of St Andrews (United Kingdom)*

Conformable metasurfaces retain all the properties of their rigid counterparts, while presenting unique opportunities both in terms of optical properties and applications. Here we discuss our most recent results in imaging, antenna design and nonlinear optics based on them.

18:10 : Invited talk**Interplay of Absorption and Scattering in Metal-Dielectric Cylinders: Extreme Refractive Index Limits and Response Sensitivities****Nasim Mohammadi Estakhri¹, Nooshin M. Estakhri²**¹ *Chapman University (USA)*, ² *University of Michigan, Ann Arbor (USA)*

Structural singularities such as sharp corners are linked to extreme field concentration and presence of hotspots in small particles, typically resulting in enhanced absorption efficiency. In this work we analytically investigate the interplay of absorption and scattering in a class of singular metal-dielectric particles in the shape of joined half-cylinders. We will discuss the role of refractive indices and material loss in the response of these geometries and pinpoint several phenomena including resonant scattering/absorption, sharp absorption response, and invisibility.

18:30 : Keynote talk**Tunable Light-Matter Coupling in Low-Dimensional Excitonic Semiconductors****Deep Jariwala***University of Pennsylvania (USA)*

The focus of this talk will be light-matter interactions in excitonic semiconductors such as 2D transition metal dichalcogenides, carbon nanotubes as well as hybrid perovskites. Results on exciton-plasmon coupling and exciton-polariton formation under reflectance and emission spectroscopy will be presented.

17:30 - 18:45 — Ibn Al-Haytham Room**Session 2A35****Exotic Meta-media - Time-dependent, Nonlocal and Other Novel Responses**

Organized by: Martin McCall, Jonathan Gratus and Paul Kinsler

Chaired by: Martin McCall, Jonathan Gratus and Paul Kinsler

17:30 : Invited talk**Boundary Conditions in Time Dependent Materials****Rebecca Seviour¹, Johnathan Gratus², Paul Kinsler²**¹ *University of Huddersfield (United Kingdom)*, ² *Lancaster University (United Kingdom)*

Currently there is growing interest in time dependent media, materials whose constitutive relations change with time. Most models assume the permittivity and permeability are constant. This approach works if both permittivity and permeability are real. However to model lossy materials it is natural to consider complex constitutive relations. In this paper we demonstrate with a simple example that such a model is unphysical in a time dependent media. Furthermore we derive the correct boundary conditions necessary dependent media.

17:50 : Invited talk**Twisted light in metamaterials with spatial dispersion****Alexey Tishchenko***National Research Nuclear University "MEPhI"(Russia)*

We report on the theoretical studies of generation of non-classical light carrying orbital angular momentum and its propagation in metamaterials with spatial dispersion. The distinctive feature of such a light is a longitudinal component of the field, the effect of which increases the degree of its twisting. As a mechanism of

generation of such waves we consider Cherenkov radiation. We also discuss the role of relativistic effects and causality principle in terms of the phenomena considered.

18:10 : Invited talk

Refraction and Impedance Patterns in Moving Media

Zoe-Lise Deck-Leger¹, Christophe Caloz²

¹Polytechnique Montréal (Canada), ²Katholieke Universiteit Leuven (Canada)

To study wave propagation in moving media, a common graphical tool is the isofrequency diagram, or refractive index pattern, which provides the direction and the velocity of a wave. Here, we complement this tool with the impedance pattern, which provides information on the wave amplitude. We argue that the two tools should be used on an equal footing when solving problems involving moving media.

18:30 : Nonlocal and Soft Plasmonics in Ion Particle Chains

Christin David

Friedrich-Schiller-University Jena (Germany)

Plasmonic properties of charged fluids are discussed via a two-fluid model. Additional degrees of freedom (ion mass, ion charge, concentration) allow a wide range of bulk resonance frequencies. In analogy to metal nanoparticles, I include commonly neglected charge interactions thus deriving analytic expressions for non-local Mie coefficients for coupled charge carriers. Plasmonic ionic response is studied in microspheres and chains and compared to metal systems.

17:30 - 18:25 — Gaston Floquet Room

Session 2A36

Parity-Time and quasi-normal modes in Photonics, Plasmonics, Acoustics

Organized by: Anatole Lupu and Henri Benisty

Chaired by: Anatole Lupu and Henri Benisty

17:30 : Invited talk

Non-Hermitian Properties of Photonic and Plasmonic Nanocavities

Christophe Sauvan

Institut d'Optique - CNRS (France)

Photonic and plasmonic nanocavities confine light at a deep subwavelength scale but generally suffer from large radiative leakage and absorption losses. As a consequence, nanocavities are inherently non-Hermitian systems. The non-Hermitian character provides them peculiar properties, especially regarding the control of spontaneous emission. Quasinormal modes, i.e., natural eigenmodes of a non-Hermitian system, are the adequate tools for describing light-matter interactions in photonic and plasmonic nanocavities.

17:50 : Invited talk

Supersymmetric signatures of topological states

Henning Schomerus

Lancaster University (United Kingdom)

By contrasting settings of photonic graphene, topological CROWs, quantum walks, and mesoscopic devices, I clarify the physical consequences of supersymmetry, which is hidden behind the mathematical description of topological effects in chiral systems but leaves concrete signatures in experimentally observable states.

18:10 : Observation of exceptional arcs and cusp singularities in acoustics

Kun Ding¹, Weiyuan Tang¹, Xue Jiang¹, Yi-Xin Xiao², Zhao-Qing Zhang², C. T. Chan², Guancong Ma¹

¹Imperial College London (United Kingdom), ²Hong Kong University of Science and Technology (China)

The known fact that two singularities can be connected by geometric arcs reveals fundamental excitations of materials, such as Fermi arc and the associated interface modes. However, it is still an open question that which geometric structure connects two higher-order exceptional points (EPs). In this work, by using the

ternary acoustic cavity system we demonstrate both theoretically and experimentally that it is the exceptional arc that bridges two order-3 EPs and produces the cusp singularity at the order-3 EP.

17:30 - 18:15 — Lawrence Bragg Room

Session 2A37

Machine learning for metamaterials and metasurfaces

Organized by: Mohamed Bakr and Willie Padilla

Chaired by: Mohamed Bakr and Willie Padilla

17:30 : Nanostructured Materials for artificial neural computing

Zongfu Yu

University of Wisconsin Madison (USA)

We show that optical waves passing through a nanophotonic medium can perform artificial neural computing. Complex information, such as an image, is encoded in the wave front of in-put light. The medium continuously transforms the wave front to realize highly sophisticated computing tasks such as image recognition.

17:45 : Unidirectional non-Hermitian structures on demand by genetic optimization

Waqas W. Ahmed¹, Ramon Herrero², Muriel Botey², Ying Wu¹, Kestutis Staliunas²

¹King Abdullah University of Science and Technology (Saudi Arabia), ²Universitat Politècnica de Catalunya (UPC) (Spain)

We propose a general approach based on genetic optimization to achieve 'on demand' asymmetric light transport in non-Hermitian structures. The procedure allows designing the imaginary part of permittivity distribution from a given (arbitrary) real part of permittivity distribution for asymmetric reflection in a broad range of frequencies. We demonstrate a selective spectral unidirectional light reflection control, in such a way that it switches from left to right (or vice versa) with varying operating frequency.

18:00 : Inverse design of one-dimensional multilayer structure by artificial neural network

Kirill Safronov, Vladimir Bessonov, Andrey Fedyanin

Lomonosov Moscow State University (Russia)

We propose new method of artificial neural network (ANN) training for inverse design of multilayer structures. The ANN directly learns to predict parameters of structures by target reflection spectrum. The transfer matrix method is employed during ANN training to overcome the issue of non-unique inverse problem solution. We compare the target spectrum with spectrum of structure predicted by ANN during the training process. Our ANN is able to design structures with a target spectrum with high fidelity.

18:15 - 18:55 — Lawrence Bragg Room

Session 2A38

Opto-mechanical metasurfaces and metamaterials

Organized by: Dibakar Roy Chowdhury, Jayasri Dontabhaktuni and Harshavardhan Kalathur

Chaired by: Dibakar Roy Chowdhury, Jayasri Dontabhaktuni and Harshavardhan Kalathur

18:15 : Invited talk

Extreme Nonreciprocity in Spatio-Temporally Modulated Metasurfaces

Andrew E. Cardin, Sinhara R. Silva, Shai. R. Vardeny, Wilton J. M. Kort-Kamp, Hou-Tong Chen, Diego A. R. Dalvit, Abul K. Azad

Los Alamos National Laboratory (USA)

In this work, we experimentally demonstrate a spatio-temporal modulated reflectarray metasurface that is capable of dynamical wave-front shaping and nonreciprocal propagation of free space electromagnetic radiation. Arbitrary space-time phase distributions in reflection are achieved by embedding electronically modulated varactors into the resonators of our metasurface. Our experimental measurements reveal on-demand wave-front control of frequency conversion processes. We also demonstrate maximum violation of Lorentz reciprocity in both beam steering due to nonreciprocal excitation of surface waves along the metasurface.

18:35 : Invited talk

Applications of metasurface optics in atom-based sensors

Jennifer T. Choy

University of Wisconsin - Madison (USA)

The discrete energies of atomic electrons and the ability to probe and control them with light have enabled a host of precision measurements. In this talk, I will describe the benefits and challenges of state-of-the-art atomic sensors via examples of atom-based inertial sensors and magnetometers. I will discuss critical developments in nanophotonic engineering that are needed to improve the utility and performance of these sensors and our progress towards developing a photonic-integrated atomic magnetometer using metasurface polarization optics.

17:30 - 18:30 — Rene Descartes Room

Session 2A39

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

17:30 : Invited talk

Strong light-matter coupling: new trends in plexitronics

Yury P. Rakovich

CSIC-UPV/EHU (Spain)

We report our recent results of research of plasmon-exciton interactions in Au and Au@Ag nanorods integrated with J-aggregates of cyanine dyes is presented. In all developed hybrid nanostructures, the anti-crossing behavior of the hybridized modes can be tracked using a number of spectroscopic techniques such as absorption, photoluminescence (PL) and magnetic circular dichroism (MCD).

17:50 : Invited talk

In situ electron energy-loss spectroscopy for nanoscale optical devices

Soren Raza

Technical University of Denmark (Denmark)

We demonstrate dynamic electromechanical control over the coupling of a gold nanodisk dimer all the way to sub-nanometer-sized gaps. By combining EELS with in situ electrical actuation, we can follow the evolution of the gap size and optical properties with unprecedented spatial and spectral resolution. We show that our electromechanical device can be used as light modulator with low power consumption and high speed.

18:10 : Invited talk

Plasmonic dynamic color modulation

Eric Hopmann, Abdulhakem Y. Elezabi

University of Alberta (Canada)

Plasmonic-electrochromic ("plasmochromic") devices have recently gained significant interest in the research community, due to the dynamic optical properties of the electrochromic material and the high sensitivity of the plasmon to its dielectric environment. Here, plasmochromic resonance modulation is used to create a dynamic reflective display with a wavelength modulation of over 64 nm in the visible range. The results are verified via FDTD analysis, which projects a maximum wavelength shift of over 100 nm.

18:00 - 19:00 — Tutorial Room**Session 2A40****Conference Tutorials II**

Organized by: Ishwar Aggarwal

18:00 : Tutorial**Design of Active and Reconfigurable Metasurfaces****Harry A. Atwater***California Institute of Technology (USA)*

A grand challenge for nanophotonics is the realization of comprehensively tunable metasurface nanoantenna arrays enabling dynamic, active control of the key constitutive properties of light – amplitude, phase, wave-vector and polarization. Achieving this will open new photonics applications in phased-array optical beam steering, visible light modulation for communications and thermal radiation management. This tutorial will discuss design approaches for active and reconfigurable metasurfaces including selection of active materials, electromagnetic design and time-modulation. We will also survey status and outlook for electronically tunable and reconfigurable plasmonic and all-dielectric metasurfaces, whose elements are arbitrarily reprogrammable, enabling a wide array of functions, including steering, focusing, and frequency multiplexing of scattered radiation.

Thursday 22nd July, 2021

09:00 - 11:00 — Victor Veselago Room

Session 3A1

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

09:00 : **Invited talk**

Enhancing the second harmonic generation from nonlinear crystals by plasmonic nanostructures

Emre Gurdal, Anke Horneber, Nadim Shaqura, Alfred J. Meixner, Dieter P. Kern, Dai Zhang, Monika Fleischer

Eberhard Karls Universitat Tübingen (Germany)

Frequency conversion plays an important role in non-linear optics, and nonlinear crystals are frequently employed as optical elements. However, processes such as frequency doubling by second harmonic generation (SHG) still suffer from limited conversion efficiency. Here we combine commercial LiNbO₃ crystals with plasmonic nanoparticles in order to boost SHG from the crystal surface. The interaction is fundamentally investigated for a single nanodisc, and maximized by a dense array of particles.

09:20 : **Invited talk**

Hybrid metal/organics in strong coupling for switchable fluorescence

Joel Bellessa, Jean-Michel Benoit, Clementine Symonds, Kevin Chevrier

University Lyon 1 (France)

In this paper we evidence bifunctionality properties, photochromaticity and luminescence, in strongly coupled optical system. Strong coupling has been achieved between an electromagnetic mode, the surface plasmon, and two organic emitters: a J-aggregate, known for its high absorption properties and a photochromic material which absorption can be optically switched on and off. We show that the emission of the aggregated dye can be reversibly shifted between the activated and deactivated form of the photochromic material.

09:40 : **Invited talk**

Sensing Spatial Coherence of Light with Planar Metallic Metamaterials

Thomas Frank, Oleksandr Buchnev, Tamsin Cookson, Malgosia Kaczmarek, Pavlos Lagoudakis, Vassili A. Fedotov

University of Southampton (United Kingdom)

We report on a discovery that homogeneous metallic non-diffracting metamaterials of a certain type respond differently to spatially coherent and incoherent light, enabling robust speckle-free discrimination between different degrees of coherence. The effect has no direct analogue in natural optical materials and may find applications in compact metadevices enhancing imaging, vision, detection, communication and metrology.

10:00 : **Invited talk**

Measurement of the quantum geometric tensor and of the anomalous Hall drift in a continuous medium

Antonio Gianfrate¹, Olivier Bleu², Lorenzo Dominici¹, Vincenzo Ardizzone¹, Milena De Giorgi¹, Dario Ballarini¹, Giovanni Lerario¹, Kenneth West³, Loren N. Pfeiffer³, Dmitry Solnyshkov², Daniele Sanvitto¹, Guillaume Malpuech²

¹CNR NANOTEC (Italy), ²Université Clermont Auvergne (France), ³Princeton University (USA)

We report a direct measurement of the Berry curvature and of the quantum metric in a 2D continuous photonic medium. The measured components of the quantum geometric tensor are used for a quantitative prediction of the anomalous Hall effect, which is then confirmed by experiment.

10:20 : **Invited talk**

Chiroptically Active Metal Nanoparticles Inherently composed of Atomic Chirality

Zhifeng Huang*Hong Kong Baptist University (Hong Kong)*

Chiral nanoparticles (CNPs), fabricated by glancing angle deposition with fast substrate rotation, are composed of chiral lattices at the atomic scale and bear optical activity (OA). The chiral lattices at their surfaces give rise to enantiospecific adsorption of molecules, resulting in enantioselective amplification of molecular OA and asymmetric photocatalysis. These results provide an insight into developing potential applications of functional CNPs in the practically vital fields of enantiodifferentiation, bio-detection, bio-imaging and asymmetric catalysis.

10:40 : Invited talk**Measurement of saturable absorption behavior of CNT/PDMS coated high-Q microcavity towards mode-locking of Er-doped laser****Riku Imamura, Keigo Nagashima, Takasumi Tanabe***Keio University (Japan)*

We developed a method for coating CNT/PDMS on a high-Q silica microcavity that allows us to obtain saturable absorption that is needed for building a microresonator based Er-doped mode-locked laser.

09:00 - 11:00 — Allan Boardman Room**Session 3A2****Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy**

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

09:00 : Invited talk**Photo-thermoelectric Conversion of Plasmonic Nanohole Array****Wakana Kubo***Tokyo University of Agriculture and Technology (Japan)*

We report a plasmonic photodetector consisting of an ultra-thin silver film with nanohole array, whose photo-detection mechanism is based on thermoelectric conversion triggered by plasmonic local heating. The contribution of plasmonic local heating to thermoelectric conversion is verified experimentally and numerically, in order to discuss the mechanisms governing light detection. This plasmonic photo-thermoelectric conversion will be available for improving the power conversion of thermoelectric device by creating thermal gradients across the device.

09:20 : Invited talk**Accelerated Foerster-Type Nonradiative Energy Transfer of Semiconductor Nanocrystals****Pedro Ludwig Hernandez Martinez¹, Hilmi Volkan Demir²**¹*Nanyang Technological University Singapore (Singapore)*, ²*Nanyang Technological University (Singapore)*

We present a review of accelerated nonradiative energy transfer employing semiconductor nanocrystals by using either the plasmonic effect or the dimensionality effect. We start with enhanced Foerster-type resonance energy transfer (FRET) accelerated via selective plasmon mediation between NCs. Next, we present FRET accelerated via dimensionality and show the ultrafast exciton transfer from a thin film of CdSe/CdS nanoplatelets to a MoS2 monolayer. Our findings indicate that such accelerated FRET of NCs holds great promise for optical sensing and photodetection

09:40 : Keynote talk**Chiral Near-field Properties of Plasmonic Nanomaterials: Imaging and Functions****Hiromi Okamoto***Institute for Molecular Science (Japan)*

Based on near-field and far-field optical activity microscopic methods, we showed gold nanostructures give

highly enhanced optical activity in the local sites near the nanostructures (~ 102 times enhancement as compared with macroscopic optical activity), even in achiral nanostructures, suggesting strong chiral near-field interaction. The strong chiral near-field interaction gives highly circularly polarized luminescence from achiral dye molecules adsorbed on chiral gold nanostructures. We also demonstrate dissymmetry of optical trapping behavior of chiral gold nanoparticles.

10:10 : Invited talk

Novel nonlinear chiroptical effects

Ventsislav K. Valev

University of Bath (United Kingdom)

Following our recent discovery of the Hyper Rayleigh Scattering Optical Activity (HRS OA), further developments and related, new nonlinear chiroptical effects are reported. The new effects enable chiroptical characterization in tiny volumes of illumination.

10:30 : Up- and Down-Conversions Coupled to Surface Lattice Resonances in Aluminum Periodic Arrays

Shunsuke Murai, Yuan Gao, Katsuhisa Tanaka

Kyoto University (Japan)

Aluminum is known to plasmonic up to ultraviolet frequencies, while its plasmonic performance in near infrared is less explored. In this study, for the purpose of verifying that aluminum is working effectively from ultraviolet to near infrared regions, we fabricate arrays of aluminum nanocylinders with the lattice periodicity ranging from 150 to 980 nm. Amplified up- and down-conversion are exhibited by coupling emitters on top of the lattice that resonates at their excitation wavelengths.

10:45 : Nanoscale ZnO growth via localized photothermal energy conversion in plasmonic nanoantennas

Christophe Pin¹, Hideki Fujiwara², Tatsuro Suzuki¹, Keiji Sasaki¹

¹*Hokkaido University (Japan)*, ²*Hokkai-Gakuen University (Japan)*

In this work, a new bottom-up nanofabrication method based on plasmon-assisted hydrothermal synthesis (PAHS) is demonstrated. By engineering the polarization-dependent optical and thermal properties of a gold nanogap antenna, we achieve localized growth of a few-nm-thick zinc oxide (ZnO) layer at the targeted central position of the antenna. It is numerically shown that the back-action of the material synthesis on the plasmonic resonance can be used to achieve self-limited material growth.

09:00 - 11:00 — Tatsuo Itho Room

Session 3A3

Thermal Plasmonics and Metamaterials for Environment and Energy Application

Organized by: Junichi Takahara and Kotaro Kajikawa

Chaired by: Junichi Takahara

09:00 : Invited talk

Thermophotovoltaic system using a photonic crystal emitter operating above 1300K

Masahiro Suemitsu, Takashi Asano, Susumu Noda

Kyoto University (Japan)

We demonstrate a thermophotovoltaic system comprising a selective thermal emitters based on Si rod-type photonic crystal and InGaAs photovoltaic cells. A high heat to electricity conversion efficiency of 11.2% is obtained, which is 1.65 times higher than the previously reported record value(6.8%).

09:20 : Invited talk

Development of Surface Plasmon Resonance Enhanced Photo-electric Conversion Devices

Akira Baba, Chutiparn Lertvachirapaiboon, Kazunari Shinbo, Keizo Kato

Niigata University (Japan)

Surface plasmon resonance (SPR) phenomena have attracted considerable attention since many years because of the extremely strong enhancement and confinement of electric fields near metal surfaces. Recently, we have studied grating-coupled surface plasmon-based multiple plasmonic structures for electric-field enhanced organic device applications such as biosensors and photo-electric conversion devices. In this presentation, we will introduce our recent progress on surface plasmon enhanced photo-electric conversion systems.

09:40 : Invited talk

Thermal Emissions by Tamm Plasmon Polaritons

Kuo-Ping Chen¹, Zih-Ying Yang², Satoshi Ishii², Takahiro Yokoyama², Thang Duy Dao², Mao-Guo Sun¹, Pavel S. Pankin³, Ivan V. Timofeev³, Tadaaki Nagao²

¹National Chiao Tung University (Taiwan), ²National Institute for Materials Science (NIMS) (Japan), ³Siberian Federal University (Russia)

Infrared wavelength selective thermal emission based on Tamm plasmon is experimentally demonstrated with ultra-sharp resonance. The excitation of Tamm plasmon polaritons (TPPs) occurs between a distributed Bragg reflector (DBR) and a metal film. The proposed structure could achieve twice higher Q-factor for the measured emissivity comparing to typical plasmonic thermal emitters.

10:00 : Invited talk

Titanium nitride for light-to-heat and heat-to-light conversions

Satoshi Ishii¹, Manpreet Kaur¹, Shunsuke Murali², Shinya Goya², Makoto Higashino², Katsuhisa Tanaka², Zih-Ying Yang³, Kuo-Ping Chen³, Tadaaki Nagao¹

¹National Institute for Materials Science (NIMS) (Japan), ²Kyoto University (Japan), ³National Chiao Tung University (Taiwan)

Metallic properties of titanium nitride (TiN) can be used to enhance non-radiative decay processes in the nanoscale. In the current paper, our recent works using TiN nanostructures for photothermal heating and wavelength-selective thermal emissions are reviewed.

10:20 : Invited talk

Waveguiding of Radiation from Silica Plates

Kotaro Kajikawa

School of Engineering, Tokyo Institute of Technology (Japan)

Passive radiative cooling (RC) refrigerates objects, such as solar panels, without using electricity. In this paper, improvement of RC ability of a silica plate with silica particles is demonstrated. This is due to the decrease in the reflectivity at the flat surface of a silica plate at the mid-infrared wavelength range. It is also shown that handling of RC is possible using aluminum waveguide. This expands the possibilities of the method for various applications.

10:40 : Invited talk

Metasurfaces for Heat-Shield Windows and Their Near-Infrared Reflectance Control using Movable Thin-films

Yoshiaki Kanamori

Tohoku University (Japan)

Metasurfaces for heat-shield windows which transmit visible wavelengths and reflect near-infrared (NIR) wavelengths for sun light were designed and fabricated. Moreover, by combining the metasurfaces with movable thin-films, reflectance switchable filters that could change reflectance at NIR wavelengths whereas there was little change in visible wavelengths were designed. It can be expected to be used as smart glasses that control transmittance of the thermal wavelengths in summer and winter.

09:00 - 10:50 — Christian Huygens Room

Session 3A4

Non-Hermitian Photonics: From Plasmonics, and Metamaterials, to Topological and Quantum systems

Organized by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

Chaired by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

09:00 : Invited talk

Topological field Theory far from Equilibrium

Federico Tonielli¹, Jan C. Budich², Alex Altland¹, Sebastian Diehl¹

¹University of Cologne (Germany), ²Technische Universität Dresden (Germany)

Observable properties of topological quantum matter are often described by topological field theories. We extend this principle beyond thermal equilibrium by studying the response to an external gauge field of a driven-dissipative system with a Chern insulator stationary state. We show that, if the particle number is conserved, such response is described at late times by the same abelian Chern-Simons theory valid at thermal equilibrium. As a corollary, we predict chiral edge modes stabilized by a dissipative bulk.

09:20 : Instability of nonreciprocal non-Hermitian media

Henning Schomerus

Lancaster University (United Kingdom)

I develop a general response theory for non-conserving optical, mechanical or acoustic media in which Hermiticity is broken due to nonreciprocal effects. I show that these media undergo an instability phase transition when they display the non-Hermitian skin effect. This makes these media ideally suited as sensors and amplifiers.

09:35 : Hamiltonian and Liouvillian exceptional points in noisy non-Hermitian systems

Jan Wiersig

Otto-von-Guericke-Universität (Germany)

We discuss the relation between Hamiltonian and Liouvillian exceptional points (EPs) in non-Hermitian systems with parametric noise. Conclusions for the performance of EP-based sensors are drawn.

09:50 : Topological bulk lasing mode in non-Hermitian kagome lattices

Stephan Wong, Sang Soon Oh

Cardiff University (United Kingdom)

Due its robustness against disorders, topological edge modes have been used to enhance the performances of lasers. The quality of topological lasers can be further improved using novel topological phases in one-dimensional non-Hermitian photonic topological insulators. Here, we demonstrate topologically protected mode extended over the bulk of a two-dimensional kagome lattice with rhombus geometry by introducing an imaginary gauge field. This shows the possibility to achieve a phase-locked broad-area topological lasers in two-dimensional lattices.

10:05 : Quantum interference with bianisotropic metasurface

Hong Liang¹, Kai Ming Lau¹, Wai Chun Wong¹, Shengwang Du², Wing Yim Tam¹, Jensen Li¹

¹The Hong Kong University of Science and Technology (Hong Kong), ²The University of Texas at Dallas (USA)

We investigate how quantum interference can be controlled by using bianisotropic metasurfaces. By considering a bianisotropic metasurface with material loss from metal, we numerically demonstrate an asymmetric control of quantum interference with an origin from an exceptional point of the metasurface in the classical regime.

10:20 : Keynote talk

Rigorous modal analysis of micro and nanoresonators

Philippe Lalanne

Université de Bordeaux (France)

Microcavities and nanoresonators are characterized by their modes, called quasinormal modes because they are the eigensolutions of non-Hermitian operators. In contrast to waveguide and free space modes,

quasinormal modes are not well documented in the literature, although nanoresonances play an essential role in current developments in nanophotonics. The reason is due to mathematical difficulties, see details in the recent review article [LPR 12, 1700113 (2018)], and especially to the fact that quasinormal modes cannot be normalized by their energy.

09:00 - 11:00 — Augustin Fresnel Room

Session 3A5

Light-matter interactions in new materials and meta-architectures

Organized by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

Chaired by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

09:00 : Invited talk

Near-field spectroscopy of Photonic Crystal Cavities characterized by Fano Local Density of States
 Francesca Intonti¹, Nicoletta Granchi¹, Matteo Ciardi², Dario Balestri¹, Daniele Pellegrino³, Francesco Pagliano⁴, Andrei Silov⁴, Frank van Otten⁴, Tong Wu⁵, Kevin Vynck⁵, Philippe Lalanne⁵, Andrea Fiore⁴, Massimo Gurioli¹

¹University of Florence - LENS (Italy), ²University of Florence (Italy), ³Eindhoven University of Technology (The Netherlands), ⁴Eindhoven University of Technology and nanoPHAB (The Netherlands), ⁵LP2N-Institut d'Optique (France)

Microcavities and nanoresonators are characterized by their quality factors and mode volumes. While Q is unambiguously defined, there are questions on V and, in particular, on its complex-valued character, whose imaginary part is linked to the non-Hermitian nature of open systems. The complex modal volume implies a deep revision of the Purcell factor expression, with counterintuitive effects, such as non-Lorentzian local density of states. We experimentally demonstrate these predictions in coupled photonic crystal cavities with low optical losses.

09:20 : Invited talk

Enhanced light coupling into nanostructured arrays as an enabler for advanced Raman-based metrology

Thomas Nuytten¹, Janusz Bogdanowicz¹, Andrzej Gawlik¹, Yusuke Oniki¹, Karine Kenis¹, Yusuke Muraki², Anne L. Charley¹, Claudia Fleischmann¹, Ingrid De Wolf¹, Paul van der Heide¹

¹IMEC (Belgium), ²Tokyo Electron Technology Solutions (Japan)

Recently it was found that when coherent polarized light impinges on nanoscale grating-like structures, the coupling of light is exceptionally sensitive to the parameters that define the periodicity, i.e. the pitch and CD. This effect re-opens the possibility for Raman spectroscopy to excel as a non-contact probe for phase, stress but also CD of nanostructured devices far beyond the diffraction limit. We will show how Raman spectroscopy is able to probe stress, composition, critical dimension, etc. at the nanoscale.

09:40 : Invited talk

Adaptive Photonic Metamaterials by Self-Assembly of Liquid Crystals in Nanoporous Solids

Kathrin Sentker¹, Mark Busch¹, Andriy V. Kityk², Patrick Huber¹

¹Hamburg University of Technology (Germany), ²Czestochowa University of Technology (Poland)

Self-organized multiscale porosity in terms of precise pore size, shape, and orientation has been achieved in many base materials. Here we exemplify that in combination with self-assembly of liquid crystals in pore space this provides particularly versatile pathways for the engineering of photonic metamaterials. We present temperature-dependent structural characterizations of the thermodynamic and structural self-assembly behavior of the liquid crystals confined in nanoporous solids in combination with experiments on the resulting effective optics of the hybrid materials.

10:00 : Invited talk

Exciton Resonance Tuning in Atomically-Thin Optical Elements

Jorik van de Groep¹, Jung-Hwan Song², Qitong Li², Umberto Celano³, Pieter G. Kik⁴, Mark L. Brongersma²
¹University of Amsterdam (The Netherlands), ²Stanford University (USA), ³IMEC Leuven (Belgium), ⁴University of Central Florida (USA)

Next-generation flat optics require dynamic control over optical functionalities. We demonstrate accurate control over light scattering by exciton resonances in monolayer WS₂ by controlling the dielectric environment of the monolayer. Next, we demonstrate actively-tunable and atomically-thin optical lenses by carving them directly out of monolayer WS₂. Using ion-liquid gating to dynamically manipulate the material's exciton resonance we show active modulation of the focal intensity.

10:20 : Invited talk

Frequency tripling via sum-frequency generation by single AlGaAs nanocylinders

Attilio Zilli¹, Davide Rocco², Marco Finazzi¹, Lamberto Duò¹, Carlo Gigli³, Giuseppe Marino³, Giuseppe Leo³, Costantino De Angelis², Michele Celebrano¹

¹Politecnico di Milano (Italy), ²University of Brescia (Italy), ³Université de Paris (France)

Dielectric nano-antennas of III-V semiconductor materials offer a strong nonlinear response governed by the electro-magnetic resonances of the system. We investigate individual AlGaAs cylinders and observe efficient sum-frequency generation by mixing an input beam of telecom frequency with its second harmonic. Our work highlights a convenient frequency-tripling pathway at the nanoscale based on the second-order nonlinearity of the material.

10:40 : Invited talk

Linear and Nonlinear Interactions with Epsilon-Near-Zero Metamaterials

Humeyra Caglayan

Tampere University (Finland)

We have investigated both the linear and nonlinear effects of a multilayer epsilon-near-zero metamaterial.

09:00 - 10:45 — Ibn Al-Haytham Room

Session 3A6

Metasurfaces, flat optics, FSS and HIS

09:00 : Double caustic beams generation based on metasurface

Rui Chen¹, Yi Zhou¹, Wenjie Chen¹, Ruipin Chen², Naeem Iqbal¹, Yungui Ma¹

¹Zhejiang University (China), ²Zhejiang Sci-Tech University (China)

Metasurfaces provide a powerful tool to manipulate electromagnetic waves in multidimensions and can easily integrate multiple functions into a single device. Here, we propose a compact method to generate double caustic beams by polarization multiplexing. The output can be dynamically tuned by changing the incident polarization. The research result has potential applications in dynamic beam shaping, biosensing and micro-particle manipulation.

09:15 : Ultra-Broadband Blackbody-Like Perfect Absorber

Ze-An Chen¹, Yu-Jung Lu², Ta-Jen Yen¹

¹National Tsing Hua University (Taiwan), ²Academia Sinica (Taiwan)

We propose a multilayer perfect absorber with omnidirectional independence, polarization independence and broadband response simultaneously. The proposed perfect absorber possesses higher than 80 % absorption within 900-1900 nm wavelength range and incidence angle independent in both s- and p-polarization which are consistent in both simulation and experiment. Most importantly, we can easily fabricate large area sample due to the lithography-free fabrication process which is more practical in real application.

09:30 : Broadband Metasurfaces through first Order Approximation of Surface Impedances

Ashif Aminulloh Fathnan, Andreas E. Oik, David A. Powell

University of New South Wales (Australia)

We introduce a procedure to realize a broadband metasurface based on inductance-capacitance (LC) reso-

nators in series and parallel configurations. To broaden the operational bandwidth, the L and C values must be obtained by a first order approximation method considering the equivalent surface impedance and its derivative in the center frequency. We verified our method using experiment in millimeter-wave frequencies and show that the broadband metasurface achieves a significant increase of bandwidth (more than 80 % increase) compared to a single-frequency design.

09:45 : Transparent Metamaterial Absorber with Broadband RCS Reduction for Solar Arrays

Shunliu Jiang, Xiangkun Kong, Lingqi Kong, Qi Wang

Nanjing University of Aeronautics and Astronautics (China)

In this paper, a metamaterial absorber with simultaneous high optical transparency and broadband microwave absorption is presented. By tailoring the reflection response of meta-atoms, 80 % absorption performance from 6.8GHz to 18GHz. In the meantime, by employing transparent substrates, including indium tin oxide and anti-reflection glass, good optical transmittance are obtained. The absorptivity of the proposed metamaterial absorber is simulated and measured experimentally. Both results demonstrate highly consistence which verify the reliability of our design for use in solar arrays.

10:00 : High-efficiency anomalous reflection with acoustic metasurfaces

Xing-Feng Zhu, Jie Yao, Qi Wei, Da-Jian Wu, Xiao-Jun Liu

Nanjing Normal University (China)

We present the theoretical, numerical and experimental investigations on the high-efficiency anomalous reflection (AR) by acoustic metasurfaces (AMs). Closed-end tubes are designed to realize the passive-lossless AM. A considerable improvement of 31 % in the overall efficiency is confirmed in the AM compared to that based on generalized reflection law when the incident and reflected angles are 0° and 75° respectively. Experimental results are in accord with finite element simulations and theoretical analyses.

10:15 : Influence on wide-angle metasurface doublet due to different types of all-dielectric metasurface

Hidemitsu Toba, Hidetsugu Takagi, Michio Ohashi, Katsura Otaki

Nikon Corporation (Japan)

We compare the incidence angle dependence among three types of subwavelength structures in all-dielectric polarization-insensitive metasurfaces through electromagnetic simulations. As a result, the waveguide-type metasurface was found to be most suitable for wide-angle metalenses. Therefore, we performed full-wave electromagnetic simulations of cylindrical doublet metalenses to compare the influence on lens performance between microposttype metasurfaces and waveguide-type metasurfaces. These results indicate that the waveguide-type metasurface improves the lens performance of the doublet lens previously introduced.

10:30 : Polarization-reconfigurable Water-based Frequency Selective Resorber With Four Operating Modes

Ling Qi Kong, Xiang Kun Kong, Shun Liu Jiang, Xiang Xi Yan, Lei Xing

Nanjing University of Aeronautics and Astronautics (China)

A multifunctional polarization-reconfigurable water-based frequency-selective resorber (FSR) was presented in this communication. The proposed structure is composed of a cross-gap FSS and two layers of parallel water channels fixed on the opposite sides of a dielectric substrate, where the layers are arranged in orthogonal polarization with independent control of the top and bottom channels. The novelty of the design lies in its reconfigurability between four different operating modes by injecting water or not in the top and bottom channels.

09:00 - 11:00 — Gaston Floquet Room

Session 3A7

Topology in photonic crystals, metamaterials, and metasurfaces: physics and design

Organized by: Yang Li and Benfeng Bai

Chaired by: Yang Li and Benfeng Bai

09:00 : Invited talk**Broadband Optical Modulation via Dual Epsilon-Near-Zero Modes****Qin Chen, Long Wen***Jinan University (China)*

Epsilon-near-zero (ENZ) modes have attracted extensive interests due to its ultrasmall mode volume resulting in extremely strong light-matter interaction. However, the operation bandwidth is usually limited by the ENZ wavelength range. In this paper, broadband optical modulation is demonstrated by dual ENZ modes in a TCOs/dielectric/silicon nanotrench configuration. Different types of carrier accumulations in both silicon and TCOs give rise to the ENZ states at two wavelengths determined by the carrier densities.

09:20 : Invited talk**Discovering Topological Surface States of Dirac Points In An Acoustic Crystal****Hengbin Cheng¹, Yixin Sha², Rongjuan Liu¹, Chen Fang¹, Ling Lu¹**¹Chinese Academy of Sciences (China), ²Peking University (China)

Dirac materials, unlike the Weyl materials, have not been found in experiments to support intrinsic topological surface states, as the surface arcs in existing systems are unstable against symmetry-preserving perturbations. Utilizing the proposed glide and time-reversal symmetries, we theoretically design and experimentally verify an acoustic crystal of two frequency-isolated three-dimensional Dirac points with four branched gapless helicoid surface states.

09:40 : Invited talk**Active optical metasurfaces based on VO₂ phase change materials****Tongtong Kang, Zheng Peng, Min Xiao, Jun Qin, Longjiang Deng, Lei Bi***University of Electronic Science and Technology (China)*

We report several active metasurface devices based on VO₂ hybrid metaatoms. We show several prototype devices for ultrafast optical modulation, optical switch and flexible active photonic device applications.

10:00 : Invited talk**Observation of a Topological Edge State in the X-ray Band****Zhiwei Guo, Haitao Jiang, Yong Sun, Yunhui Li, Hong Chen***Tongji University (China)*

In this work, the topological edge state is theoretically proposed and experimentally demonstrated at the interface of two kinds of photonic crystals having different bandgap topological characteristics in the X-ray regime. Remarkably, this topologically protected edge state is immune to the thickness disorder as long as the zero-average-effective-mass condition is satisfied. The results extend the topological photonics to the X-ray regime and paves the way for the development of novel X-ray optics such as high-resolution X-ray filters/monochromators.

10:20 : Invited talk**Realization of Photonic Charge-2 Dirac Point by Engineering Super-modes in Topological Superlattices****Mengying Hu¹, Kun Ding², Tong Qiao¹, Xi Jiang¹, Qiang Wang¹, Shining Zhu¹, Hui Liu¹**¹Nanjing University (China), ²Imperial College London (United Kingdom)

We report on the first experimental realization of the charge-2 Dirac point in the visible region by deliberately engineering hybrid topological states called super-modes in a 1-D optical superlattice system with two additional synthetic dimensions. Utilizing direct reflection and transmission measurements, we also show the approach to manipulating two spawned Weyl points that are identically charged in synthetic space. Moreover, topological end modes uniquely resulting from the charge-2 Dirac point can be delicately controlled within

truncated superlattice samples.

10:40 : Invited talk

Generation of optical phase vortex beams by momentum-space polarization vortices around BIC
Lei Shi

Fudan University (China)

An optical phase vortex (OV) is a beam with spiral wave front and screw phase dislocation. Here we theoretically and experimentally realized a novel but easy approach to generate OVs. This new class of OV generators operates in the momentum space, meaning that there is no real-space center of structure. Any even order of OV, which is actually a quasi-non-diffractive high-order quasi-Bessel beam, at any desired working wavelength could be achieved in principle.

09:00 - 09:40 — Lawrence Bragg Room

Session 3A8

Opto-mechanical metasurfaces and metamaterials

Organized by: Dibakar Roy Chowdhury, Jayasri Dontabhaktuni and Harshavardhan Kalathur

Chaired by: Dibakar Roy Chowdhury, Jayasri Dontabhaktuni and Harshavardhan Kalathur

09:00 : Invited talk

Nanophotonic platforms for light emitting metasurfaces

Naresh Kumar Emani, Saurabh Kishen, Jinal Tapar

Indian Institute of Technology Hyderabad (India)

In this talk we focus on two major research themes (a) Two-dimensional PT (parity-time-reversal) symmetric metamaterials and (b) Improvement of light emission efficiency in electrically driven MIM tunnel junctions. We will briefly discuss our key findings given below.

09:20 : Invited talk

Concept of mechanically tunable terahertz circular polarizer

Xiaolong You, Christophe Fumeaux, Withawat Withayachumnankul

The University of Adelaide (Australia)

A terahertz circular polarizer is developed based on an extended analytical method incorporating network analysis and genetic algorithm. Simulations reveal that the design enables a 15-dB extinction ratio relative bandwidth of 14.3%. The center operation frequency can be tuned by design from 220 to 330 GHz through varying the spacer thicknesses. The performance suggests the potential of the structure to function as a mechanically tunable terahertz circular polarizer, provided that air gaps are included in between the dielectric spacers.

09:40 - 11:00 — Lawrence Bragg Room

Session 3A9

Integrated lithium-niobate photonics: structures, devices, systems and applications

Organized by: Xianfeng Chen and Yang Li

Chaired by: Xianfeng Chen and Yang Li

09:40 : Invited talk

Spectral engineering of LNOI waveguides: from ultranarrow to broadband**Katia Gallo, Alessandro Prencipe, Halvor Fergestad***KTH - Royal Institute of Technology (Sweden)*

The development of advanced nano-structuring capabilities on LNOI is paving the way towards low-footprint photonic circuits leveraging appealing functionalities of LiNbO₃ for ultrafast signal processing and wavelength conversion. In the talk we shall present ultra-narrow bandpass and multi-resonance filters, implemented with phase-shifted Bragg gratings in LNOI photonic wires. We shall also discuss designs of dispersion engineered waveguides for broadband second harmonic generation, appealing for wavelength multicasting, ultrashort pulse frequency doubling and enhanced quadratic cascading in the telecom band.

10:00 : Invited talk**Periodically poled lithium niobate microdisk resonators on insulator****Zhenzhong Hao, Li Zhang, Qiang Luo, Xiaomei Gao, Wenbo Mao, Feng Gao, Fang Bo, Guoquan Zhang, Jingjun Xu***Nankai University (China)*

We developed a fabrication process of periodically poled lithium niobate (PPLN) photonic devices on insulator with the assistance of piezo response force microscopy (PFM), providing us the abilities to fabricate micro-domain structures down to the scale of several tens of nm. The efficient harmonic generation with d₃₃, the largest nonlinear coefficient of LN, utilized in the whole microcavity and high-order harmonic generations were demonstrated in PPLN microcavities with single and double spacial periods, respectively. The work paves the way to achieve efficient frequency conversion in on-chip LN photonic devices.

10:20 : Invited talk**Lithium niobate thin films for integrated optics****Hui Hu***Shandong University (China)*

Lithium niobate (LN) thin film on a low refractive index silica (SiO₂) cladding layer (LN on insulator, LNOI) has been fabricated by using ion implantation and wafer bonding technologies. The LN thin film has the excellent physical properties. The high refractive index contrast can enable ultra-small waveguides, facilitating the development of ultra-compact photonic integrated devices and circuits. The progress of LNOI material is reviewed.

10:40 : Invited talk**Fabrication of lithium niobate on insulator device for nonlinear frequency conversion****Shijie Liu, Yuanlin Zheng, Xianfeng Chen***Shanghai Jiaotong University (China)*

We have done a lot of work on the lithium niobate microdisk, including fabrication, second harmonic generation, cascading quadratic processes and high-Q exterior WGM. We will also present an on-chip Soltype device based on transverse electro-optic effect at the telecommunication wavelengths in a periodically poled lithium niobate on insulator ridge waveguide. Cascaded sum-frequency generation and electro-optic polarization coupling is also observed in this device.

09:00 - 09:55 — Rene Descartes Room**Session 3A10****Topology in photonic crystals, metamaterials, and metasurfaces: physics and design**

Organized by: Yang Li and Benfeng Bai

Chaired by: Yang Li and Benfeng Bai

09:00 : Invited talk**Unidirectional Radiation Enabled by Topological Charge Evolution**

Xuefan Yin¹, Jicheng Jin², Marin Soljacic³, Chao Peng¹, Bo Zhen²

¹*Peking University (China)*, ²*University of Pennsylvania (USA)*, ³*MIT (USA)*

In this work, it is demonstrated that the unidirectional emission can be realized even without mirrors on the bottom, by manipulating the evolution of topological charges in the polarization field. The observed up-down asymmetry ratio is up to 27.7 dB. Our work provides an effective approach to achieve unidirectional radiation without the need for mirrors and represents a vivid example of applying topological principles to design and optimize the photonic devices.

09:20 : Invited talk

Recent experimental and theoretical developments in synthetic dimensions including the frequency axis of light

Luqi Yuan¹, Avik Dutt², Qian Lin², Momchil Minkov², Meng Xiao³, Mingpu Qin¹, Xianfeng Chen¹, Shan-hui Fan²

¹*Shanghai Jiao Tong University (China)*, ²*Stanford University (USA)*, ³*Wuhan University (China)*

The field of synthetic dimensions in photonics is under rapid development and is interesting to the broad community. Recent efforts on the synthetic space including the frequency axis of light have been largely explored with their linear properties. Moreover, we show the possibility of creating an effective nonlinear Bose-Hubbard Hamiltonian with local interactions along the frequency dimension. Our works trigger further interests in synthetic dimensions and point towards potential possibilities for manipulating internal degrees of freedom of light.

09:40 : Discovery of topological metamaterials by symmetry relaxation and smooth topological indicators

Cyrill Bösch¹, Tena Dubcek¹, Frank Schindler², Andreas Fichtner¹, Marc Serra-Garcia¹

¹*ETH Zurich (Switzerland)*, ²*Princeton University (USA)*

Topological metamaterials have emerged, both, to realize and study topological physics, as well as in the search of practical applications. However, systematically designing topological metamaterials has remained a challenge due to the very nature of topology, its robustness. Here we present a novel approach, based on symmetry relaxation and gradient methods that, for the first time, tunes the topological index directly. We have successfully applied the method to design both conventional and higher-order topological systems.

09:55 - 11:10 — Rene Descartes Room

Session 3A11

Plasmonics and nano-optics

09:55 : Self-assembly of Cubic Plasmonic Nanoparticles for Unnaturally High Refractive Index at Optical Frequency

Ji-Hyeok Huh, Jaewon Lee, Seungwoo Lee

Korea University (Korea)

In this work, we demonstrate self-assembly of the cubic plasmonic nanoparticles superlattice to achieve unnaturally high refractive index (n) at optical frequencies (i.e., near infrared (NIR)). First, we (i) synthesize highly uniform gold nanocubes (Au NCs) and then, (ii) self-assemble them into 2D superlattice. Through this self-assembly approach, Au NC superlattices are successfully obtained over the large area, exhibiting n of 6.4 at the resonant (NIR), which were not reached thus far.

10:10 : Conformal Symmetry and its Application to Plasmon Localization

Lizhen Lu¹, Emanuele Galiffi², Kun Ding², Tianyu Dong¹, Xikui Ma¹, John Pendry²

¹*Xi'an Jiaotong University (China)*, ²*Imperial College London (United Kingdom)*

Transformation optics provides us an elegant and insightful way to harness symmetry. By leveraging the conformal symmetry, we propose a general strategy for the design of light-stopping plasmonic metasurfaces, which is able to localize surface plasmon polaritons in space longer than their plasmon lifetime. Furthermore, by investigating the properties of conformal symmetry, we stress its significance in tuning the band struc-

tures of subwavelength plasmonic systems, which can be extended to a variety of applications via different conformal transformations.

10:25 : High-Performance Transparent Conducting Electrodes (TCEs) Using Plasmonic Wires

Chin-Chien Chung¹, Rober J. Visser², B.Leo Kwak², Hyunsung Bang², Chung-Chia Chen², Wan-Yu Lin², Ta-Jen Yen¹

¹National Tsing-Hua University (Taiwan), ²Corporate CTO Office (USA)

We design the structure of metal to control the plasma frequency at near infrared region. We obtain highly transparent metal within visible regime. Accompanied by the intrinsically low resistivity, we achieve high performance transparent conducting electrode based on the structure-designed metal. We integrate the TCE with OLED. Our device demonstrates high current density and luminance.

10:40 : In-situ imaging the Hygroscopic Growth of Individual Aerosol Nanoparticles using a Surface-Sensitive Photonic Microscope

Yan Kuai, D. G. Zhang

University of Science and Technology of China (China)

The water uptake of ultrafine atmospheric aerosols affect public health, atmospheric chemistry and the Earth's climate. In this paper we present a surface-sensitive photonic microscope that was proposed to measure the hygroscopic growth of single aerosol nanoparticle, and for the first time recorded the phase change process with the good time resolution. These time-resolved measurements will be critical for understanding of the interactions between water vapor and aerosols that significantly affect chemical reactivity, aging, and atmospheric composition.

10:55 : Anapole-assisted Dark Plasmon Excitation in Hollow Silicon Disk

Kang Du¹, Pei Li¹, Heng Wang², Kun Gao¹, Ting Mei¹

¹Northwestern Polytechnical University (China), ²School of Physical Science and Technology, Northwestern Polytechnical University, Xi'an, China (China)

The anapole mode excited by silicon nanodisk provides a mildly enhanced and low loss field while exhibiting non-radiating property. Due to the special nature of anapole mode, much research has focused on the light-matter interaction engineered by the anapole-assisted enhanced field. Here, based on the anapole mode in hollow silicon disk, we proposed a new method for exciting dark plasmon mode.

Coffee Break and Exhibit Inspection

Session 3P1

Poster session V

11:00 - 11:40

P1: Temperature effects on the surface plasmon resonance in Bi₂O₃-Ag eutectic composite

Kamil Szlachetko¹, Monika Tomczyk¹, Marcin Raczkiwicz², Dorota Pawlak²

¹University of Warsaw (Poland), ²Lukasiewicz Research Network (Poland)

The temperature dependences of spectral position, width and intensity of the surface plasmons of metallic Ag nanoparticles in Bi₂O₃-Ag were studied in the temperature range from ambient conditions to 500 °C. The increase of temperature leads to nonlinear and appreciable redshift and broadening of the localized surface plasmon resonance band in Bi₂O₃-Ag. The observed dependences were compared with temperature dependences and simple Drude-Lorentz approximation.

P2: Optical studies of the Bi₂ZnOB₂O₆ glass doped with silver nanoparticles and Er³⁺ ions

Katarzyna Sadecka¹, Andrzej Klos¹, Barbara Surma¹, Ryszard Piramidowicz², Anna Jusza², Krzysztof Anders², Dorota A. Pawlak¹

¹Institute of Electronic Materials Technology (Poland), ²Warsaw University of Technology (Poland)

This work reports the optical studies of the Bi₂ZnOB₂O₆ (BZB) glass doped with silver nanoparticles (nAg) and Er³⁺ ions. The material was obtained by modified micropulling down method. LSPR of nAg resonance in BZB glass occurs at 620 nm. Annealing of BZB:nAg:Er³⁺ material at 450 °C caused two fold increase of the

luminescence at 1532 nm.

P3: Linear and nonlinear effects in metamaterials based on magnonic crystals and semiconductors

Oleg Matveev, Dmitrii Romanenko, Maria Morozova

Saratov State University (Russia)

We present results of theoretical and experimental investigations of propagation of spin waves in heterostructure magnonic crystal - semiconductor wafer. Direct current in semiconductor wafer allows to control of band gap characteristics in such structure. In addition, we observed propagating of soliton-like pulses and electrical control of number and velocity of such pulses.

P4: Quadriwave lateral shearing interferometry for characterization of metasurfaces

Samira Khadir¹, Daniel Andr n², Ruggero Verre³, Qinghua Song⁴, Serge Monneret³, Patrice Genevet⁴, Mickael K ll², Guillaume Baffou³

¹Centre de recherche sur l'h t ro pitaxie et ses applications (CRHEA) (France), ²Chalmers University of Technology (Sweden), ³Institut Fresnel (France), ⁴Universit  Cote d'Azur (France)

We report in this work a new investigation method for characterization, inspection and diagnosis of meta-surface based devices. It is an optical method based on quadriwave lateral shearing interferometry (QLSI), a high-resolution, quantitative phase microscopy technique capable of mapping both the intensity and phase profiles of a light beam. In this study, we focus on conventional metalenses and linear phase-gradient metasurfaces, including resonant and geometrical Pantcharatnam-Berry phase metasurfaces.

P5: Radiative rate enhancement in spherical lead halide perovskite particle for optical cooling

Pavel Tonkaev, George Zograf, Sergey Makarov

ITMO University (Russia)

Recent studies demonstrated high interest in halide perovskite materials for different applications including solar cells, light emitting diodes and many others. Moreover, it was revealed that perovskite film can be cooled by laser irradiation due to effective up-conversion. In the other hand, particle supporting Mie-resonance can enhance spontaneous emission rate, which has crucial importance for optical cooling. In this work we theoretically reveal enhancement of radiative rate emission in single spherical lead halide perovskite particle.

P6: Boosting Faraday rotation in a one-dimensional coupled resonator magnetoplasmonic structure made by silica matrix doped with magnetic nanoparticles

Mounir Bouras, Nassim Dermeche, Ahmed Kahlouche, Abdesselam Hocini

University of Mohamed Boudiaf of M'sila (Algeria)

The present study aimed to evaluate the magneto-optic Faraday rotation of one-dimensional coupled resonator magnetoplasmonic structure by metallic cover layer in each resonator. To this purpose, transfer matrix method was used where crystals made by SiO₂/ZrO₂ or SiO₂/TiO₂ doped with magnetic nanoparticles using sol-gel process in different configurations and use and 10-nm thick gold or silver layer for the excitation of the surface plasmon polaritons (SPPs).

P7: Doping an Eutectic 3D Material to get Plasmonics Effects

Miguel Cuerva¹, Marcin Raczewicz², Dorota A. Pawlak²

¹University of Warsaw (Poland), ²Institute of Electronic Materials Technology (Poland)

Due to the development of new manufacturing technologies and the increasing availability of nanomaterials, plasmonic has become an emerging field of photonic research. Although the fabrication of metal elements has already been widely demonstrated, the development of three-dimensional plasmonic materials progresses slowly. Hence, we report the development of a eutectic compound that incorporates nanometric silver to provide surface plasmonic resonance around 600 nm

P8: Competing Interactions in Doped Rare-earth Manganites nanostructural system

Wiqar Hussain Shah

International Islamic University (Pakistan)

The Structural, magnetic and transport behavior of La_{1-x}CaxMnO₃+(x=0.48, 0.50, 0.52 and 0.55 and =0.015) compositions close to charge ordering, was studied through XRD, resistivity, DC magnetization and AC susceptibility measurements. With time and thermal cycling (T<300 K) there is an irreversible transformation of the low-temperature phase from a partially ferromagnetic and metallic to one that is less ferromagnetic and highly resistive.

P9: Underground Wireless Communication based on Metamaterial enhanced Magnetic Induction for Smart Irrigation**B. Raagavi, Sugumar Swathi, S. Sakthivel M.***Anna university (India)*

An appropriate communication strategy for transmitting underground information to achieve agricultural automation is based on magnetic induction techniques. To increase the transmission distance range, metamaterial is introduced in-between the magnetic induction coils. The proposed Permeability Negative Metamaterial is placed between the magnetic induction coils to increase the received power strength. In future, metamaterial enhanced magnetic induction based underground wireless communication system can be used effectively to transmit the sensor data to the farmer's mobile app using cloud.

P10: Quantum Dot Enhanced Performance of Plasmonic Refractive Index sensor for Cancer Detection**Simitha S.¹, Jesly Jacob², Vibin Ipe Thomas¹***¹CMS college, Kottayam (India), ²Assumption College, Changanacherry (India)*

This paper proposes a novel biosensor based on quantum dots for detection of diseased cells. We aim to report a simple, label-free and enhanced figure of merit (FOM) plasmonic cavity model which has potential to exhibit sensitivity and selectivity for detecting cancer cells. Proposed structure is studied using finite Element Method. The structure with and without quantum dot is optimized to make biosensor with better performance. The proposed sensor with dot shows improved sensitivity and FOM.

P11: Modelling Nanostructures for Application in Plasmonically Enhanced Hot-Electron Devices**Noushin Dolati, Kylie Catchpole, Thomas White, Fiona J. Beck***Australian National University (ANU) (Australia)*

Here we design nanostructures to have strong, tunable absorption in Au nanoparticles (NPs) for application in hot electron devices. Using 2D optical modelling, we show the tunability in small NPs absorption in the infrared spectral region by mounting NPs on grating structures. We demonstrate that introducing a thin dielectric layer between the NPs and the gratings can increase the absorption in the NPs up to 3.8-fold.

P12: Spectroscopic properties of freshwater diatoms frustules: a type of self-assembled nanoporous silica structures**Ankur Gogoi¹, Jayur Tisso¹, Partha P. Nath², Gazi A. Ahmed²***¹Jagannath Barooah College (India), ²Tezpur University (India)*

Fourier Transform Infrared (FTIR) and photoluminescence (PL) spectroscopy of nanoporous silica frustules of various fresh water diatoms were conducted to investigate their potential as unique silica structures for photonic and luminescent device applications. The internal chemical structure of the frustules was revealed by the FTIR spectra which showed characteristic peaks for diatom biosilica, such as bending of siloxane (Si-O-Si) and stretching of silanol (Si-OH) groups.

P13: Spatial filtering Enabled Spectral Characterization of a Photonic Crystal Cavity**Shilpi Gupta, Naresh Sharma, Govind Kumar, R. Vijaya***IIT Kanpur (India)*

We experimentally demonstrate a technique to estimate resonance wavelength of a photonic crystal cavity by spatially filtering a laser beam incident at different angles and imaging the transmitted beam profile. The transmitted beam exhibits two features: an annular beam and a central spot. Under the resonance condition, the two features overlap spatially. We develop calibration curves for the spectral characterization using transfer matrix method. The estimates from our cost-efficient technique match well with measurements obtained using a spectrophotometer.

P14: A Double layer SRR Meta- material Absorber for C band applications**Shubhangi Mahendra Palekar, Neeraj Rao***VNIT Nagpur (India)*

Metamaterials have great applications in microwave circuits. They are used to improve antenna parameters. Metamaterial absorber absorb maximum of incident radiation in certain frequency band. Here, a circular and a square split ring resonator are individually designed. Then a circular split ring is placed over a square ring resonator and the combined structure is simulated. Simulation results shows that the obtained structure is a double layer meta material absorber for C band applications.

P15: Acousto-Optic Coupling in a fishbone Phoxonic Nanobeam CavityJin-Chen Hsu¹, Chieh-Chun Chang², Tzy-Rong Lin²¹National Yunlin University of Science and Technology (Taiwan), ²National Taiwan Ocean University (Taiwan)

Acousto-optic (AO) coupling in a fishbone phoxonic nanobeam cavity is studied. Because of the dual photonic and phononic bandgaps in the structure, optical and acoustic modes can be confined in the gradient phoxonic cavity. With the spatial overlap between the modes, AO coupling can be enhanced. Boosting of the interface effect mainly contributes to the AO coupling enhancement. Here we calculate the phoxonic band structures, cavity modes, and coupling rates to discuss the AO coupling in the proposed system.

P16: Acoustic Imaging by an Three-dimensional Acoustic Meta-lens

Sang-Hoon Kim, Jung-Woo Kim

Mokpo National Maritime University (Korea)

A 3D acoustic Luneburg lens or an acoustic Luneburg ball of the radius of 60 mm was manufactured by a 3D printer. The focusing ability was simulated by the commercial FEM analysis program, COMSOL Multiphysics simulator and measured in experiment, too. Acoustic plane wave of 5–15 kHz were applied to the lens and the acoustic gain was obtained. The 3D imaging by the Luneburg ball is presented.

P17: The power law behaviors of Q factor for impedance matched zero index media

Liangsheng Li, Yong Zhu, Jie Gao, Hongcheng Yin, Yansong Wang

Science and Technology on Electromagnetic Scattering Laboratory (China)

The reflectivity, resonance width and eigen-frequency have been investigated for a single slab with the impedance matched zero index models. Moreover, we found that the resonance widths exhibit the power law behaviors in the vicinity of BICs. Typically, for the IMZI model, the resonance widths are proportional to θ . Finally, the power law behaviors of real frequency are the sub-leading term, and the exponent values are dependent on the model.

P18: Active photo-tuning of resonances in symmetric THz Fano MetasurfacesSubhajit Karmakar¹, Ravendra Kumar Varshney¹, Dibakar Roy Chowdhury²¹Indian Institute of Technology (IIT) (India), ²Mahindra Ecole Centrale (India)

We theoretically propose active photo tuning of different resonances in symmetric terahertz (THz) Fano metasurfaces. With the increase of substrate conductivity, dipole dip and transparency peak get depleted keeping Fano dip intact. The relative effect of photo-tuning also depends on lattice periodicity. Such results can be extremely beneficial to several futuristic metaphotonics applications in THz regime.

P19: A transparent fluoropolymer for daytime radiative coolingHoang Thi Thanh Tam¹, Mana Toma¹, Takayuki Okamoto¹, Mio Hidaka², Kensuke Fujii³, Yasuhiro Kuwana³, Kotaro Kajikawa¹¹Tokyo Institute of Technology (Japan), ²AGC Inc. (Japan), ³AGC Inc., Marunouchi (Japan)

We introduce the fluoropolymer film that is transparent 98% of incident sunlight and emitting 97% in the atmospheric transparency window. By the indoor experiment, the 50 μm -thick fluoropolymer film cools to 7.6 °C below ambient air temperature. Moreover, the paint-ability and the weather-resistant allow to apply the fluoropolymer film in the wide range of human life.

P20: Towards Spiral-Like Cold Atom States

Vasileios E. Lembessis, Anwar Al Rsheed, Andreas Lyras, Omar M. Aldossary

King Saud University (Saudi Arabia)

We solve the Schrodinger equation and calculate the wave functions and corresponding energies for a Rb atom trapped by the optical dipole potential created by the interaction of the atom and a Helical Optical Tube (HOT) light field. The wave functions have a twisted spiral-like spatial structure. Our work paves the way for the generation of twisted atom laser beams.

P21: Polarization-insensitive broadband achromatic metalens from ultraviolet to near-infrared regions

Yue Li, Zheng Peng, Lei Bi, Longjiang Deng, Jianliang Xie, Bo Peng

University of Electronic Science and Technology of China (China)

Metalenses have presented extraordinary abilities in compact optical devices. However, there still remains a challenge of achromatic aberration, which limits the usefulness for broadband applications. Here, we propose a polarization-insensitive metalens made of multi-layer metasurfaces to achieve a broadband achromatic

focusing from ultraviolet to near-infrared regions. A wavelength-independent focal length is achieved at 355, 450 and 785 nm with focusing efficiencies of 12 %, 30 % and 58 %. Our work provides a general approach in applications of various flat achromatic devices.

11:40 - 12:40 — Victor Veselago Room

Session 3A12

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

11:40 : **Invited talk**

High-performance Ultraviolet Metasurfaces using Wide-bandgap Dielectrics

Cheng Zhang¹, Wenqi Zhu², Junyeob Song², Qingbin Fan³, Lu Chen², Jinghui Yang², Ting Xu³, Amit Agrawal², Henri J. Lezec²

¹Huazhong University of Science and Technology (China), ²National Institute of Standards and Technology (USA), ³Nanjing University (China)

I will present our recent efforts of achieving low-loss ultraviolet metasurfaces using wide-bandgap dielectric materials. The demonstrated devices perform an array of representative wave-front shaping functionalities and operate down to the deep-UV regime, with efficiencies up to 72 %.

12:00 : **Invited talk**

Suppressing Exciton Interaction in 2D Semiconductors

Jeongyong Kim

Sungkyunkwan University (Korea)

We investigated the effect of metal screening of exciton interaction in 1L-WS2 by using Au film substrate spaced by hBN layer. We found that PL lifetime and QY can be greatly enhanced at high exciton density, which we attributed to the suppressed exciton-exciton annihilation due to screened exciton interaction.

12:20 : **Invited talk**

Flat Nonlinear Optics with Intersubband Polaritonic Metasurfaces

Daeik Kim¹, Jaeyeon Yu¹, Inyong Hwang¹, Seongjin Park¹, Hyeongju Chung¹, Frederic Demmerle², Gerhard Boehm², Mikhail A. Belkin², Jongwon Lee¹

¹Ulsan National Institute of Science and Technology (Korea), ²Technische Universitat Munchen (Germany)

In this work we report electrically tunable nonlinear response and giant nonlinear circular dichroisms based on intersubband polaritonic nonlinear metasurfaces. Experimentally we achieved 0.75 μm of the second harmonic generation spectral peak tuning and over 86 % of nonlinear circular dichroisms around 10 μm wavelength.

11:40 - 12:45 — Allan Boardman Room

Session 3A13

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

11:40 : **Keynote talk**

Photophysical Effects behind the Efficiency of Hot Electron Injection in Plasmon-Assisted Catalysis

Miguel A. Correa-Duarte

Universidade de Vigo (Spain)

The present study examines the joint role played by the morphology and composition of plasmonic nanoparticles in their photosensitizing capabilities. The critical influence of these parameters is evidenced by the effect of Au and core-shell Au@Ag nanorods on a TiO₂-driven photochemical probe reaction. In this case, the use of the bimetallic composites leads to a remarkable increase in the photocatalytic activity of the semiconductor compared to that found for the monometallic Au sensitizers.

12:10 : Invited talk

Excitons in nanophotonic landscapes: fluctuating, diffusing, annihilating

Alberto G. Curto

Eindhoven University of Technology (The Netherlands)

Excitons in atomically thin semiconductors are sensitive to their electronic and photonic environments. Here we present our experimental and theoretical results on the fluctuation, diffusion, and annihilation of excitons near nanostructures. Our results have implications for exciton-based sensors, single-photon sources based on 2D materials, and efficient and high-power light-emitting devices.

12:30 : New Synthesis Approach of Aluminum Nanoparticles for UV- Plasmonics

Marion Castilla¹, Silvère Schuermans¹, Florian Lamaze¹, Thomas Maurer¹, Gil Markovich², Uri Hananel², Davy Gérard¹, Jérôme Martin¹, Jérôme Plain¹, Julien Proust¹

¹University of Technology of Troyes (UTT) (France), ²Tel-Aviv University (Israel)

Aluminum nanostructures appear to be a good alternative to gold or silver because of the broad range of their plasmonic resonances (UV to NIR) and their reduced cost. In this paper, we present a new way of synthesis of Al nanoparticles based on sonochemistry and solvothermal reaction. By tuning the solvothermal reaction time, we are able to control the size of the nanoparticles between 10 to 100nm. finally, we will present some applications of such nanoparticles.

11:40 - 12:50 — Tatsuo Itho Room

Session 3A14

Thermal Plasmonics and Metamaterials for Environment and Energy Application

Organized by: Junichi Takahara and Kotaro Kajikawa

Chaired by: Kotaro Kajikawa

11:40 : Invited talk

A Suspended TiN film for Thermal Plasmonics Platform

Kentaro Iwami, Yuta Otome, Azusa Tahara, Satoshi Ikezawa

Tokyo University of Agriculture and Technology (Japan)

In this study, A suspended TiN film on a silicon substrate is studied and fabricated through sputter deposition and sacrificial layer etching. The balance of the internal stresses between TiN and sacrificial layers are found to be important to release suspended film structures. Modification of spectral reflectance was demonstrated by introducing hole array structure to the suspended film.

12:00 : Omnidirectional Narrow-band Thermal Radiation by Mie Resonator on Refractory Metal

Junichi Takahara, Akihiro Kawano

Osaka University (Japan)

We propose a narrow-band thermal radiation emitter based on a refractory metasurface composed of Germanium Mie resonators on Tungsten substrate. The metasurface radiates omnidirectional thermal radiation with a quality-value Q of 132 and an emissivity of 0.96 at mid-IR range in simulation. We fabricated the device and demonstrated Q~34 at 7.8micron in experiment.

12:15 : Invited talk

New polaritonic materials for radiative cooling

Juliana Jaramillo¹, G. L. Whitworth², A. Francone², N. Kehagias², P. D. Garcia², C. M. Sotomayor-Torres²

¹*Universitat Autònoma de Barcelona (Spain)*, ²*Catalan Institute of Nanoscience and Nanotechnology (Spain)*

Implementing sustainable methods for cooling is urgent, due to global warming and the high energy consumption associated to modern cooling technologies. In this work, we propose novel polaritonic materials as an energy-efficient solution for above-ambient radiative cooling.

12:35 : Heat transfer regulation for textiles using tailorable metallic wires

Muluneh G. Abebe¹, Eric Khouzakoun², Jeremy Odent¹, Jean-Marie Raquez¹, Sylvain Desprez², Bjorn Maes¹

¹*University of Mons (Belgium)*, ²*Materia Nova (Belgium)*

We numerically explore the concept of dynamic, switchable infrared transmittance using electromagnetic and thermal calculations, for the use of smart, temperature regulating textiles. We discuss the photonic effects of metallic and shape-memory-polymer coated mono-filaments on the temperature dependent transmittance of the textile fabric.

11:40 - 12:25 — Christian Huygens Room

Session 3A15

Non-Hermitian Photonics: From Plasmonics, and Metamaterials, to Topological and Quantum systems

Organized by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

Chaired by: Konstantinos Makris, Manas Kulkarni, Ramy El-Ganainy and Sahin Ozdemir

11:40 : Nonlinear spectral singularities and tunable laser with 2D material coating

Hamed Ghaemidzicheh

Lancaster University (United Kingdom)

We investigate the application of nonlinear spectral singularity in a nonlinear non-Hermitian optical system consisting of an infinite planar slab that is coated with a two-dimensional (2D) material in arbitrary transverse electric TE and transverse magnetic TM modes. Here, we explore the effects of placing the slab between Graphene and 2D Weyl semimetal sheets. We show that the 2D material introduces additional physical parameters for tuning the output intensity of the laser.

11:55 : Quantum correlations in PT-symmetric systems

Federico Roccati¹, Salvatore Lorenzo¹, Gioacchino Massimo Palma¹, Gabriel Landi², Matteo Brunelli³, Francesco Ciccarello¹

¹*Università degli Studi di Palermo (Italy)*, ²*Universidade de Sao Paulo (Brazil)*, ³*(United Kingdom)*

We study the dynamics of correlations in a paradigmatic setup to observe PT-symmetric physics: a pair of coupled oscillators, one subject to a gain one to a loss. Quantum correlations (QCs) are created, despite the system being driven only incoherently, and can survive indefinitely. We link PT-symmetry breaking to the long-time behavior of QCs, which display different scalings in the PT-broken/unbroken phase and at the exceptional point (EP). The EP in particular stands out as the most classical configuration.

12:10 : Exotic atom-photon interactions in a non-Hermitian photonic lattice

Federico Roccati¹, Salvatore Lorenzo¹, Giuseppe Calajò², Gioacchino Massimo Palma¹, Angelo Carollo¹, Francesco Ciccarello¹

¹*Università degli Studi di Palermo (Italy)*, ²*CFO-Institut de Ciències Fotoniques (Spain)*

We study emission properties and dipole-dipole interactions for a set of quantum emitters (atoms) coupled to a photonic lattice with engineered losses which exhibits the non-Hermitian skin effect. A number of exotic quantum optics effects occur such as loss-induced chiral emission, exactly localized metastable dressed states, chiral photon-mediated atom-atom interactions. At a lattice exceptional point, the effective couplings between the emitters are exactly non-reciprocal and short-range.

11:40 - 12:40 — Augustin Fresnel Room

Session 3A16

Light-matter interactions in new materials and meta-architectures

Organized by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

Chaired by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

11:40 : **Invited talk****Switchable nanooptics with conducting polymer nanoantennas****Magnus Jonsson***Linköping University (Sweden)*

I will present our recent research on switchable nanooptical resonances in nanostructures of organic conducting polymers, and the use of such materials in display applications.

12:00 : **Invited talk****Excitonic Behavior and Photo-Carriers Transport in 2D Quantum Confined Metal Organic Chalcogenides****Lorenzo Maserati, Sivan Refaely-Abramson, Christoph Kastl, Mirko Prato, Stefano Pecorario, Bianca Passarella, Andrea Perinot, Anna Anupa Thomas, filippo Melloni, Mario Caironi***Istituto Italiano di Tecnologia (Italy)*

We unveil anisotropic 2D excitons in a self-assembled, layered bulk silver benzeneselenolate coordination polymer, $[\text{AgSePh}]_{\infty}$. This in-plane covalently bonded hybrid multiple quantum well nanostructure, resemble the 2D hybrid metal halides perovskites, but it is non-polar and air stable. We therefore investigate the charge carriers' transport across a $[\text{AgSePh}]_{\infty}$. The photo-response of this material suggests possible use of this materials as air-stable UV photodetector UV even on flexible substrates.

12:20 : **Invited talk****Spin dynamics in thin films and nanostructures based on Fe60Al40****Anna Semisalova¹, Tanja Strusch¹, Ralf Meckenstock¹, Rantej Bali², Jonathan Ehrler², Kay Potzger², Kilian Lenz², Jürgen Lindner², Michael Farle¹**¹*University of Duisburg-Essen (Germany)*, ²*Institute of Ion Beam Physics and Materials Research (Germany)*

We report on ferromagnetic resonance detected spin pumping in Fe60Al40/Pd and Fe60Al40/permalloy bilayers, and laterally patterned Fe60Al40 nanostructures with periodical ferromagnetic/paramagnetic interfaces. Due to its magnetostructural phase transition Fe60Al40 can be utilized as a paramagnetic or ferromagnetic material at the same temperature depending on its structural order parameter. In this study we investigate the dual role of this alloy as a spin source and a spin sink.

11:40 - 12:40 — Ibn Al-Haytham Room

Session 3A17

Light-matter interactions in new materials and meta-architectures

Organized by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

Chaired by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

11:40 : **Tunable and dynamic structural colors in nano-optics: toward future applications****Junsuk Rho, Jaehyuck Jang***Pohang University of Science and Technology (POSTECH) (Korea)*

In this abstract, I will discuss our efforts in realizing multifunctional a-Si:H metaholograms that can encode multiple pieces of information in a monolayer device for anticounterfeiting applications.

11:55 : Redox-tunable structural colouration by UV-patterned conducting polymer nanofilms on metal surfaces**Shangzhi Chen, Magnus P. Jonsson***Linköping University (Sweden)*

In this presentation, I will introduce our latest studies on the redox state tunable structural colouration based on conducting polymers and the use of UV-patterning technique for fabricating multi-colour images within single steps.

12:10 : Plasmonic Electronic Paper**Marika Gugole, Oliver Olsson, Jolie Blake, Kunli Xiong, Andreas Dahlin***Chalmers University of Technology (Sweden)*

We work on developing reflective displays (electronic paper) in color by combining plasmonic nanostructures and electrochromic materials. The main motivation is to save energy in comparison with emissive displays. (Further details in submitted file.)

12:25 : Narrow Linewidth Plasmonic Color filters for Transmission with Enhanced Out-Of-Band Suppression**Anabel De Proft¹, Kristof Lodewijks², Pol Van Dorpe², Xavier Rottenberg²***¹Katholieke Universiteit Leuven (Belgium), ²IMEC (Belgium)*

We report on a novel plasmonic color filter design leveraging the interference of a dielectric post array and a metallic hole array, complemented with a metal cover on the dielectric posts to suppress out-of-band transmission. Simulations predict a combination of linewidth (down to 25 nm), transmission (45 %) and out-of-band transmission surpassing previously reported plasmonic color filters for transmission.

11:40 - 12:40 — Gaston Floquet Room**Session 3A18****Topology in photonic crystals, metamaterials, and metasurfaces: physics and design**

Organized by: Yang Li and Benfeng Bai

Chaired by: Yang Li and Benfeng Bai

11:40 : Invited talk**Topological bulk laser****Renmin Ma***Peking University (China)*

Here, we propose and experimentally demonstrate a topological bulk laser in a judiciously architected nanocavity array, where a bulk state is harvested for lasing in a topological way. The topological bulk laser is based on a novel band-inversion induced reflection, which provides a new lasing-mode selection mechanism and renders emission directionality.

12:00 : Invited talk**Prediction of topological transitions in photonic crystals using machine learning****Bei Wu¹, Kun Ding², Che Ting Chan³, Yuntian Chen¹***¹Huazhong University of Science and Technology (China), ²Imperial College London (United Kingdom), ³Hong Kong University of Science and Technology (China)*

We train artificial neural networks to distinguish the geometric phases of a set of bands in photonic crystals by feeding the network with the information of photonic crystals encoded with Maxwell's equations. Interestingly, we find that the trained network yields remarkably accurate predictions of the topological phases for photonic crystals, even for those geometric and material parameters that are beyond the range of the parameter space in the training dataset.

12:20 : **Invited talk**

Six-dimensional light-matter interaction

Ouyang Xu, Yi Xu, Yaoyu Cao, Xiangping Li

Jinan University (China)

Pursuing new physical dimensions for tailoring and manipulating light-matter interaction is valuable from fundamental science and application points of view. In general, the intrinsic disorder effect during nanofabrication is generally considered to be useless. In sharp contrast to topological photonics that is proposed to mitigate the involuntary disorder effect, we utilize the combination of intentional disorder effect in plasmonic crystal and the polychromatic and polarized vortex beam to synthesize a record high dimension for light-matter interaction.

11:40 - 12:40 — Lawrence Bragg Room

Session 3A19

Plasmonics and nano-optics

11:40 : Mimicking a Wormhole and Giant Tidal Force with Curved Optical Spaces

Runqiu He, G. H. Liang, S. N. Zhu, H. Liu

Nanjing University (China)

We apply curved optical waveguides to simulate the evolution of a beam around a wormhole and we found the huge tidal forces acting on a beam around a wormhole, which cause the wave packets deformed or even destroyed. The experimental system provides an effective method for simulating gravity and will be helpful for Astronomical observation and prediction in the future.

11:55 : Fabrication of GaN nanopillars by metal-assisted chemical etching combined with ICP etching

Xiaomeng Zhang, Hailiang Li, Changqing Xie

Institute of Microelectronics, Chinese Academy of Sciences (China)

We present a facile and effective approach for fabricating GaN nanopillar arrays, using a combination of inductively coupled plasmas(ICP) etching and Pt/Au metal assisted chemical etching (MacEtch). By introducing defects in the metal covered area before wet etching we can achieve a faster process and greater depth. Further, the effects and significances of illumination conditions on the etching process are demonstrated.

12:10 : Electron kinetics in epsilon-near-zero optical nonlinearity

Heng Wang, Kang Du, Xinhai Dai, Wending Zhang, Soo Jin Chua, Ting Mei

Northwestern Polytechnical University (China)

The physical mechanisms of epsilon-near-zero (ENZ) optical nonlinearity are modeled and the Drude model is extended to intraband transition induced optical nonlinearity by adopting the statistical parameters. The electron overall effective mass and overall scattering time are proposed, which takes into account the distribution of energy dependent electrons in the nonparabolic band.

12:25 : Enhanced circular polarization discriminative photoresponse in the metamaterial integrated anisotropic active materials

Jing Zhou, Zeshi Chu, Xu Dai, Xiaoshuang Chen, Wei Lu

Chinese Academy of Sciences (China)

Circular polarization discrimination is desired for use in many optoelectronic applications. Nevertheless, the circular polarization extinction ratio (CPER) for absorption of the active materials, typically below 2.5. Based on double polarization selection mechanism, the integration of an asymmetric metamaterial with an anisotropic material can drastically enhance CPER by 6 to 10 times and enhance photoresponse of active materials.

11:40 - 12:40 — Rene Descartes Room

Session 3A20

Structured and topological photonic fields

Organized by: Jamal Berakdar

Chaired by: Jamal Berakdar

11:40 : Invited talk

Compact Meta-Spectrometer for Mobile Applications

jaesoong lee

Samsung Advanced Institute of Technology (Korea)

We demonstrated an extremely compact and efficient meta-structure-based spectrometer for use in the near-infrared range. The spectrometer consists of a mobile-phone CMOS imager and silicon-nanoposts-embedded dielectric multilayers fabricated directly on top of the imager. The spectrometer shows good transmission and excellent spectral resolutions. In addition, the presence of the metaposts embedded in the dielectric layers greatly simplifies the fabrication process to generate individual spectral channels. It shows promise of integrating compact spectrometers in smartphones for diverse applications.

12:00 : Invited talk

Photonic nodal rings associated with electromagnetic anti-duality symmetry

Ruo-Yang Zhang, Qinghua Guo, Biao Yang, Che Ting Chan

Hong Kong University of Science and Technology (Hong Kong)

We investigate a kind of dispersive chiral media, and show that there exists a novel kind of nodal rings existing out of any high-symmetry planes in the momentum space. These nodal rings arise from the accidental electromagnetic anti-duality symmetry at a specific frequency. More interestingly, when two such materials with opposite chiralities are adjoined at an interface, the anti-duality symmetry also ensures that the bands of interface states can intersect at the anti-duality frequency, forming two-fold degenerate Fermi arcs.

12:20 : Invited talk

High harmonic generation with topological light fields

Jonas Wätzel, Jamal Berakdar

Martin-Luther-University Halle-Wittenberg (Germany)

Results of quantum mechanical calculations of an atomic gas irradiated by three-dimensional topological light fields are presented. The investigation of the high harmonic generation reveals that the generation of topologically protected and polarization-structured light fields in the X(UV) frequency regime is possible with only one driving field.

Lunch

12:40 - 14:00

14:00 - 15:15 — Victor Veselago Room

Session 3A21

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

14:00 : Invited talk

Controlling the propagation of light and sound in silicon waveguides by means of subwavelength nanostructuring

Jianhao Zhang¹, Dorian Oser¹, Thi-Thuy-Duong Dinh¹, Xavier Le Roux¹, Miguel Montesinos¹, Christian Lafforgue¹, Diego Perez-Galacho², Daniel Benedikovic¹, Elena Duran Valdeiglesias¹, Vladyslav

Vakarin¹, Omar Ortiz¹, A. Rodriguez³, Olivier Alibert³, Pavel Cheben⁴, Sebastien Tanzilli³, Laurent Labonté³, Delphine Marris-Morini¹, Eric Cassan¹, Daniel Lanzillotti-Kimura¹, Laurent Vivien¹, Carlos Alonso-Ramos¹

¹Université Paris-Saclay (France), ²Universitat Politècnica de Valencia (Spain), ³Université Côte d'Azur (France), ⁴University of Ottawa (Canada)

Periodically patterning silicon with a pitch sufficiently small to suppress diffraction effects opens new degrees of freedom to control light propagation in silicon photonic circuits with unprecedented flexibility. Concurrently, near-infrared photons and GHz phonons in nanoscale Si waveguides have comparable wavelengths. Thus, subwavelength nanostructuring has a great potential to shape light-sound interactions in silicon waveguides. Here, we present our most recent results on the subwavelength engineering of silicon for the implementation of high-performance photonic and optomechanical components.

14:20 : Invited talk

On-chip guiding of spoof terahertz surface plasmon polaritons on metasurface pathways

Sven Becker, Tassilo Fip, Marco Rahm

TU Kaiserslautern (Germany)

We investigate the guiding and routing of spoof terahertz surface plasmon polaritons (terahertz SSPPs) along metasurface pathways of subwavelength width by means of numerical calculations and experimental measurement of the spatio-temporal and spectro-temporal dynamics of the electric field of the SSPPs. The SSPPs are routed along straight and curved pathways of subwavelength path width. We show that terahertz SSPPs can be tightly guided within subwavelength space on metasurfaces without loss of the out-of-plane confinement.

14:40 : Invited talk

Tailoring Nanoscale Order in Metal and Semimetal - Based Metamaterials for Enhanced Light Harvesting, Sensing, and Switching

Johann Toudert, Rosalia Serna

Instituto de Optica-CSIC (Spain)

Self-assembled metamaterials based on metal and semi-metal nanostructures are reported. Their nanoscale order is tailored to achieve enhanced optical properties, such as a broadband near-perfect absorption of light or a broad optical phase tuning. They enable an optimal sunlight harvesting, 3D-multiplexed remote sensing and synaptic optical switching.

15:00 : Experimental Probes of Thermal Transport and Lattice Stability of Important Optoelectronic Semiconductor Nanomaterials

Richard D. Schaller

Northwestern University (USA)

We performed transient X-ray diffraction experiments on semiconductor nanocrystals and lead halide perovskites. Bragg peak shifts relate heating and peak amplitude reduction confers lattice disordering. II-VI nanoparticles melt upon absorption of ~ 0.89 excitations/nm³. Certain perovskites are found to undergo solid-solid phase transitions prior to disordering. Diffraction intensity recovery kinetics occur over hundreds of picoseconds with slower recoveries for larger particles. These findings highlight questions of physical stability of advanced materials and related electronic impacts in high intensity excitation applications.

14:00 - 15:15 — Allan Boardman Room

Session 3A22

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

14:00 : Invited talk

Light management at the nanoscale bioinspired by photosynthesis photonics

William P. Wardley, Johannes. W. Goessling, Miguel Castillo, Martin Lopez Garcia

International Iberian Nanotechnology Laboratory (Portugal)

Cost-effective nanofabrication methods can be used to reproduce the phenomena of enhanced absorption and structural colour found in the leaves of some plants. Such nanophotonic biomimetic approaches can work as a playground to better understand the natural system and as inspiration for new strategies on light harvesting. We will also show that the silica exoskeleton of diatom microalgae contains slab photonic crystals that can be used as a naturally produced nanophotonic material, opening the door to bioengineered nanophotonic devices.

14:20 : Invited talk

On the mechanisms of plasmon-enhanced chiroptical response

Thomas Weiss¹, Steffen Both¹, Egor A. Muljarov²

¹University of Stuttgart (Germany), ²Cardiff University (United Kingdom)

We present a rigorous theory based on the resonant state expansion that allows us to analyze all electromagnetic contributions to the plasmonic enhancement of the chiroptical response of chiral media. Potential applications are optimized optical sensors for molecular handedness detection.

14:40 : Invited talk

Optical injection of plasmonic Janus-nanoparticles into living cells

Theobal Lohmüller

LMU München (Germany)

15:00 : Doping zinc oxide nanohybrids for application in white light emitting diodes

Alexandra Apostoluk¹, Yu Zhang¹, Christophe Theron¹, Thibaut Cornier¹, Beata Derkowska², Malgorzata Sypniewska², Stéphane Danièle¹, Bruno Masenelli¹

¹Université de Lyon (France), ²Nicholas Copernicus University (Poland)

We present a study concerning the effect of doping in inorganic/organic ZnO/PAAH (polyacrylic acid) nanohybrids. The doping atoms vary in their ionic size, electronic valence and concentration. Some of them have been known to provide ZnO with magnetic properties. The effect on the optical properties of the ZnO nanoparticles of three different concentrations (0.1 %, 1 % and 5 %) of dopants is compared. The luminescent properties of the undoped and doped nanohybrids are reported.

14:00 - 15:20 — Tatsuo Itho Room

Session 3A23

Thermal Plasmonics and Metamaterials for Environment and Energy Application

Organized by: Junichi Takahara and Kotaro Kajikawa

Chaired by: Junichi Takahara

14:00 : Invited talk

Complex metamaterials for energy applications

Andrea Fratalocchi

Kaust University (Italy)

In this invited talk I will summarize recent research of my group on complex metamaterials for various energy and environmental applications ranging from world record water splitting, photo-catalysis, solar desalination and structural coloration.

14:20 : Invited talk

Smart Meta-optical Solar Reflector based on Vanadium Dioxide

Kai Sun¹, Wei Xiao¹, Ioannis Zeimpekis¹, Mirko Simeoni², Alessandro Urbani², Matteo Gaspari², Sandro Mengali², Ivano Indiveri³, Behcet Alpat³, Lars Kildebro⁴, Javier Aizpurua⁵, Dan Hawak¹, C.H. (Kees)

de Groot¹, Otto L. Muskens¹

¹University of Southampton (United Kingdom), ²Consorzio C.R.E.O. (Italy), ³Maprad S.r.l. (Italy), ⁴NIL Technology (Denmark), ⁵CSIC-UPV/EHU (Spain)

Optical solar reflectors (OSRs) play a crucial role in the thermal control of a spacecraft. We present novel 'smart' metasurface-based OSRs using vanadium dioxide (VO₂), which has its infrared emittance modulated by temperature. We have investigated the infrared property of the ALD VO₂ films under different anneal conditions and found the anneal condition can optimize VO₂ infrared response. We have demonstrated that a superior emittance tunability and lower solar absorption for meta-OSRs than the planar film device.

14:40 : Invited talk

Titanium nitride and silicon nanostructures for photothermal applications

Osamu Takayama¹, Evgeniy Shkondin¹, Shinya Goya², Makoto Higashino², Shunsuke Murai², Katsuhisa Tanaka², Satoshi Ishii³, Tadaaki Nagao³, Andrei V. Lavrinenko¹

¹Technical University of Denmark (Denmark), ²Kyoto University (Japan), ³National Institute for Materials Science (Japan)

We report photothermal conversion of high aspect ratio TiN and Si nanostructures. We fabricate two types of TiN nanostructures: trenches and tubes, and two types of Si nanostructures: trench and air hole arrays in Si. We study the photothermal properties of these nanostructures by measuring Raman shift of the materials that corresponds to the temperature in terms of laser power. These high aspect ratio nanostructures exhibit significant temperature increase, showing the potential for highly efficient photothermal converters.

15:00 : Invited talk

Metasurface thermal emitter

Ming Zhou, Zongfu Yu

Wisconsin Madison (USA)

We show metasurface emitter can exploit collective wave effects of thermal photons to realize focused thermal emission and thermal holography.

14:00 - 15:00 — Christian Huygens Room

Session 3A24

Local enhancement and control of light-matter interaction

Organized by: Antonio Ambrosio

Chaired by: Antonio Ambrosio

14:00 : Invited talk

Nanooptics in Hyperbolic Van der Waals Materials

Javier Martin-Sanchez

University of Oviedo (Spain)

Polaritons - hybrid light-matter excitations - play a crucial role in fundamental and applied sciences, as they enable nanoscale control of light. Here, we present propagation of hyperbolic polariton along the surface of the Van der Waals (vdW) material α -MoO₃ with amplitude record lifetimes. Additionally, we will discuss possible applications for the in-plane manipulation of light based on the in-plane and ray-like directional propagation of polaritons. Our findings open new avenues for fully planar photonic technologies at the nanoscale.

14:20 : Invited talk

P-plates: Pure vortex generation with a single metasurface

Marco Piccardo, Antonio Ambrosio

Fondazione Istituto Italiano di Tecnologia (Italy)

Since the introduction of the concept of optical vortex in 1989 a number of methods to generate such beams have emerged. After presenting a timeline tracing the appearance of vortex generation tools in the last 30 years, we will introduce p-plates - metasurfaces enabling the creation of vortices with high purity.

14:40 : Invited talk

Flat Optics and High Quality Factor Nanoresonators with Phonon Polaritons in Van Der Waals Materials

Michele Tamagnone

Fondazione Istituto Italiano di Tecnologia (Italy)

New types of nanophotonic devices based on phonon polaritons in hexagonal boron nitride and molybdenum trioxide are demonstrated. Different substrates (metal, phase change materials or other van der Waals materials) can be used to engineer the propagation of these polaritons. High quality factor resonators using isotopically pure boron nitride are also discussed. Polaritons in van der Waals materials are an excellent mid infrared counterpart to plasmon polaritons in mobile metals.

14:00 - 15:20 — Augustin Fresnel Room

Session 3A25

Light-matter interactions in new materials and meta-architectures

Organized by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

Chaired by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

14:00 : Invited talk

Light-absorption in nano-antennas: from self-heating to reconfigurable metasurfaces

Giulia Tagliabue

EPFL (Switzerland)

By engineering light absorption in dielectric and metallic nanoantennas, we explore new opportunities for the manipulation of temperatures at the nanoscale as well as the design of reconfigurable metasurfaces.

14:20 : Invited talk

Plasmon-driven chemical reactions: photothermal effects, near-fields, and hot charge carriers

Andrea Baldi

Vrije Universiteit Amsterdam (The Netherlands)

Metal nanoparticles scatter and absorb light thanks to surface plasmon resonances. The decay of these resonances is used to drive chemical reactions via photothermal heating, electromagnetic field enhancement, and ejection of hot charge carriers. All these mechanisms typically occur on ultrafast timescales and are therefore very difficult to disentangle experimentally. In our group, we develop strategies to disentangle and quantify the relative contribution of these activation mechanisms in plasmon-driven chemical reactions, both in ensemble and at the single-particle level.

14:40 : Invited talk

Time-Resolved Cathodoluminescence in a Transmission Electron Microscope

Sophie Meuret¹, Nikolay Cherkashin¹, Luiz Tizei², Yves Auad², Robin Cours¹, Sebastien Weber¹, Florent Houdellier¹, Mathieu Kociak², Arnaud Arbouet¹

¹CEMES/CNRS (France), ²Laboratoire de Physique des Solides (France)

Time resolved Cathodoluminescence (TR-CL) is a unique technique that allows to measure the lifetime of radiative transition down to the picosecond at the nanometer scale. In this presentation, we will discuss the first experimental demonstration of TR-CL in a transmission electron microscope. We will show its potential, and the opportunities offer by its complementarity with other electron based spectroscopies, to correlate the optical and structural properties of materials.

15:00 : Invited talk

Unlocking the far-IR potential in plasmonics with metal oxide perovskites

Nikolaos Kalfagiannis

Nottingham Trent University (United Kingdom)

Metal-oxide perovskites are important technological materials due to their excellent properties. Here we ex-

Explore the optical properties of SrTiO₃ and BaTiO₃ in the mid- and far-infrared regime. In this spectral range, both materials demonstrate an exceptional behavior due to their combined phonon-polariton modes allowing their real permittivity to remain negative for an extended region, in contrast to other polar dielectrics. We thus present the exceptional flexibility of both materials as active nano-phonic components.

14:00 - 14:40 — Ibn Al-Haytham Room

Session 3A26

Structural Color for Displays and Imaging

Organized by: Debashis Chanda

Chaired by: Debashis Chanda

14:00 : **Invited talk**

Structural Blue: Journey from fundamental research to real-life application as first omnidirectional structural color pigment

Debasish Banerjee

Toyota Research Institute of North America (USA)

In this talk we will discuss optical design strategies based on multilayer stack of thin-films to maintain brilliancy, omnidirectional characteristics of structural colors as well as feasible production methods for commercialization. This work paves the way to access of durable naturally brilliant and high chroma colors to automotive market and beyond.

14:20 : **Invited talk**

High-purity hybrid structural colors by enhancing optical absorption of organic dyes in resonant cavity

L. Jay Guo, Zhengmei Yang, Chengang Ji

University of Michigan (USA)

We present a simple and new method of incorporating an ultrathin dye film into a planar thin-film structure to realize high-purity reflective colors with drastically increased lifetime.

14:40 - 15:25 — Ibn Al-Haytham Room

Session 3A27

Photonic bandgap structures

14:40 : **Free space topological surface states at the surface of square photonic crystals**

Anna Tasolamprou¹, Maria Kafesaki¹, Costas Soukoulis², Eleftherios Economou¹, Thomas Koschny²

¹IESL - FORTH (Greece), ²Iowa State University (USA)

We present a square photonic crystal that sustains topological surface states at the free space interface. Band structure and direct scattering simulations demonstrate the topological surface mode unidirectionality and immunity to defects and back-scattering.

14:55 : **Characteristics of Transmission GAPS in Photonic Hypercrystals**

Munazza Zulfiqar Ali

Punjab University (Pakistan)

Here we theoretically investigate the wave propagation in two different types of photonic hypercrystals. The emergence of different types of transmission gaps in frequency as well as in momentum space and some of the characteristics of these gaps such as their dependence on filling ratio of HMM and angle of incidence is

studied by curve plotting and simulations.

15:10 : Self-action of Bloch surface waves in a one-dimensional photonic crystal

Daniil A. Shilkin, Evgeny V. Lyubin, Andrey A. Fedyanin

Lomonosov Moscow State University (Russia)

Bloch surface waves (BSWs) in one-dimensional photonic crystals are considered as a promising platform for two-dimensional integrated optics and optical manipulation of particles. In this contribution, we present an experimental study of nonlinear optical effects that occur when BSWs are excited at the interface with a water suspension of dielectric nanoparticles. The results are in agreement with our theoretical predictions and highlight the potential of BSWs for use in two-dimensional devices operating in a nonlinear regime.

14:00 - 15:15 — Gaston Floquet Room

Session 3A28

Micro/Nano fabrication and characterization techniques

14:00 : Laser-Induced Metal/Polymer Composites For Flexible Electronics

Anna Lipovka¹, Raul D. Rodriguez¹, Fedor Gubarev¹, Sergey Schadenko¹, Gennadiy Murastov¹, Alexey Zinoviev¹, Ilia Petrov¹, Ammar Al-Hamry², Olfa Kanoun², Jin-Ju Chen³, Evgeniya Sheremet¹

¹Tomsk Polytechnic University (Russia), ²Chemnitz University of Technology (Germany), ³University of Electronic Science and Technology of China (China)

In this work, we present a fast, inexpensive, single-step, and scalable route for metal nanoparticles (MNPs) integration to polymer structures using laser beams. We propose the formation of highly conductive and mechanically robust aluminum/polyethylene terephthalate (Al/PET) composites for flexible electronic devices fabrication. These results are in accordance with our previous research connected to the incorporation of graphene-like materials into PET which indicates the versatility and generality of the laser processing method.

14:15 : What Influences Surface Plasmon Resonance Linewidth in MIM Structures Obtained by Colloidal Self-Assembly

Ye Yu¹, Daniel Schletz², Johanna Reif¹, Felix Winkler¹, Matthias Albert¹, Andreas Fery², Robert Kirchner¹

¹Technische Universität Dresden (Germany), ²Leibniz-Institut für Polymerforschung Dresden e.V. (Germany)

LSPRs have been extensively studied in the past decades. However, they often suffer from spectral broadening, a crucial origin of which is fabrication inaccuracies. Unfortunately, there is yet an efficient way to evaluate this impact, rendering the researchers having to rely on rather intensive, time-consuming experimentations. We present an approach to evaluate various fabrication contributors to the broadening, providing a possibility of efficient optimization in lieu of experiments, shedding light to automated fabrication, optimization of integrated design.

14:30 : Development of a high throughput metalens fabrication process relying on Bosch Deep Reactive Ion Etching and UV Nano Imprint Lithography

Christopher A. Dirdal, Geir Uri Jensen, Hallvard Angelskar, Jo Gjessing, Paul C. Vaagen Thrane

SINTEF Microsystems and Nanotechnology (Norway)

Current optical components are often bulky, heavy and expensive - thereby inhibiting the application areas of optical technology. Metasurfaces are highly promising in this respect due to their potential of making optical components small, lightweight and cheap. However, virtually all diffraction limited dielectric metasurface lenses to date rely on slow and expensive direct writing methods. At SINTEF Microsystems and Nanotechnology we are developing industrially relevant Bosch Deep Reactive Ion Etching and UV Nano Imprint Lithography processes for metalens fabrication.

14:45 : Comparative study of monocrystalline and polycrystalline gold plasmonic nanorods

Lukas Kejik, Michal Horak, Tomas J. Sikola, Vlastimil Krapek

Brno University of Technology (Czech Republic)

Plasmonic antennas are often fabricated by lithographic patterning of a thin metallic film and its properties

are then intimately related to the quality of the resulting structures. Here we compare two kinds of thin gold films: polycrystalline thin film deposited by magnetron sputtering, and chemically synthesized monocrystalline gold platelet. Both metallic substrates were used to fabricate plasmonic nanorods using focused ion beam lithography. The resulting nanorods were characterized by scanning transmission electron microscopy and electron energy loss spectroscopy.

15:00 : **Microstructural Properties of ZnO Powder Nanostructures and Ab Initio Study**

Salah Oudjertli, Abdelkader Safou

Research Center In Industrial Technologies (Algeria)

We used X-ray diffraction (XRD). Powder X-ray diffraction data confirm the formation of a hexagonal crystal structure with space group $P63mc$ of ZnO. The ZnO replicas was clearly observed under an optical microscope(OM). Band structure and density of states of the phase of crystal ZnO computed using Ab Initio methods, confirmed that pure ZnO is a indirect band gap semiconductor for B1 phase, whose phase B1 is of NaCl type(Rocksalt).

14:00 - 15:15 — Lawrence Bragg Room

Session 3A29

Plasmonics and nano-optics

14:00 : **Electromagnetism and Plasmon-Enhanced Light-Matter Interactions at the Nanoscale**

P. A. D. Goncalves¹, T. Christensen², N. Rivera¹, A.-P. Jauho³, N. A. Mortensen¹, M. Soljagic¹

¹University of Southern Denmark (Denmark), ²MIT (USA), ³Technical University of Denmark (Denmark)

We present a theory for quantum nanoplasmonics that incorporates nonlocality and quantum effects, such as electronic spill-out and surface-enabled Landau damping. Focusing on the planar and sphere geometries, we derive analytical expressions for the nonclassical scattering coefficients, from which the optical response of the systems can be unambiguously determined. We compute the systems' plasmonic excitations and investigate the role of quantum surface corrections to plasmon-emitter interactions, e.g., Purcell enhancement, dipole-forbidden transitions rates, and plasmon-mediated energy transfer.

14:15 : **Influence of primary beam energy on localized surface plasmon resonances mapping by STEM-EELS**

Michal Horak, Tomas Sikola

Brno University of Technology (Czech Republic)

We present an experimental study of the influence of primary beam energy on localized surface plasmon resonances mapping by scanning transmission electron microscopy combined with electron energy loss spectroscopy. The best results are obtained using a medium primary beam energy (120 keV) as the primary beam energy should be high enough to suppress the scattering in the sample and at the same time should be low enough to avoid the appearance of relativistic effects.

14:30 : **Towards implantable SERS substrates: testing the fabrication approaches**

Raul Rodriguez¹, Ilia Petrov², Elizaveta Sviridova², Elena Dorozhko², Ali Issa³, Evgenia Sheremet², Pavel Postnikov², Safi Jradi³

¹Tomsk Polytechnic University (Russia), ²Tomsk Polytechnic University (Russia), ³University of Technology of Troyes (France)

The use of Surface Enhanced Raman Spectroscopy (SERS) in vivo has attracted increasing interest in biomedical applications. In this work, we demonstrate different approaches for in vivo SERS substrates fabrication and outlook on their perspectives and future applications.

14:45 : **Highly Anisotropic Tungsten Oxide Nanocrystals for Tunable NIR-Plasmonic Absorption and Scattering**

Yannis Cheref, Capucine Cleret de Langavant, Eric Larquet, Axel Laborieux, Thierry Gacoin, Jongwook Kim

LPMC - Ecole Polytechnique (France)

We report on the plasmonic optical properties in the near-infrared range of Hexagonal Tungsten Bronze (HTB) nanocrystals. We present full LSPR bands dynamics where particles grow from strongly colored HTB platelets to long HTB rods invisible to the eye. Furthermore, control of the particle size allowed us to measure LSPR light scattering for the first time in doped semiconductor nanoparticles, despite carrier densities in these materials being lower than in the well-studied noble metal nanoparticles.

15:00 : **Embedded Annular-Hole Arrays Enabling Multiband Near-Zero-Index Transmission at Near-Infrared**

Andriy E Serebryannikov¹, Hodjat Hajian², Ekmel Ozbay²

¹Adam Mickiewicz University (Poland), ²Bilkent University (Turkey)

Annular-hole periodic arrays enabling high-efficiency transmission in multiple bands are studied at near-infrared. Up to four subwavelength transmission bands can be obtained by placing several annular holes within each unit cell. High efficiency in transmission is achieved due to near-zero-index behavior at the cutoffs of the plasmonic modes propagating along the coaxial wave guide channel.

14:00 - 15:20 — **Rene Descartes Room**

Session 3A30

Structured and topological photonic fields

Organized by: Jamal Berakdar

Chaired by: Jamal Berakdar

14:00 : **Invited talk**

Optical super-resolution sensing of a trapped ion's wave packet size

Martin Drechsler¹, Sebastian Wolf¹, Christian Tomas Schmiegelow², Ferdinand Schmidt-Kaler¹

¹Universitaet Mainz (Germany), ²IFIBA, UBA CONICET (Argentina)

In this work we demonstrate a super-resolution optical scheme to determine the size of the wave packet of a single trapped ion. Our method is inspired by the well known ground state depletion (GSD) technique. We use a hollow beam to strongly saturate a dipole-forbidden transition around a sub-diffraction limited area at its center and observe state dependent fluorescence. By spatially scanning this laser beam over a single trapped 40Ca^+ ion, we are able to distinguish the wave packet sizes of ions cooled to different temperatures. Using a depletion beam waist of $4:2(1) \mu\text{m}$ we reach a spatial resolution which allows us to determine a wave packet size of $39(9) \text{ nm}$ for a near ground state cooled ion.

14:20 : **Invited talk**

The physics of the magnetoelectric near fields

Eugene Kamenetskii

Ben Gurion University of the Negev (Israel)

A continuous variation of field structure across the interfaces defines the near field. In a case of near fields of dielectric materials, one observes the evanescent wave character of the fields with continuous variation of field amplitudes and energies across the interfaces. In a case of a material with magnetoelectric (ME) properties, along with the question of variation of field amplitudes, questions also arise of variation of field phases and angular-momentum conservation across the interfaces.

14:40 : **Invited talk**

field Distributions and Atom Trapping in Focused Axially-Shifted Counter-Propagating light Beams

Koray Koksal¹, Vassilis E. Lembessis², Jun Yuan³, Mohamed Babiker³

¹Bitlis Eren University (Turkey), ²King Saud University (Saudi Arabia), ³University of York (United Kingdom)

This talk deals with a special kind of structured light when focused optical beams interfere with their focal planes shifted axially by a finite distance d . For doughnut beams we show how these lead to finite ring lattices, Ferris-wheels and conveyor belts. Furthermore, a new all-optical atom trapping environment arises solely due to the scattering force on atoms when the beam waists are of sub-wavelength dimensions. Our

findings are discussed with reference to sodium atoms.

15:00 : Invited talk

Metasurface spin-to-orbital angular momentum converters

Antonio Ambrosio

Istituto Italiano di Tecnologia (Italy)

We demonstrate beams with independent OAM coupled to user-defined linear or circular polarisation states, all from the same laser. Our nanostructured metasurfaces are compact and power scalable, for the creation of arbitrary angular momentum states of structured light.

Coffee Break and Exhibit Inspection

Session 3P2

Poster session VI

15:20 - 16:00

P1: Tuning the decay coefficient of sound in a 2D viscous metamaterial

Jesus Arriaga¹, José Doporto¹, Martin Ibarias¹, Arkadii Krokhin²

¹Universidad Autonoma de Puebla (Mexico), ²University of North Texas (USA)

The homogenization theory developed previously for a phononic crystal of cylinders embedded in a viscous fluid is used to calculate the decay coefficient of sound due to viscosity. We consider different Bravais lattices and different cross sections of the cylinders in order to tune the decay coefficient of sound. We observe that, in the low-frequency limit a phononic crystal with asymmetric unit cell behaves like a dissipative homogeneous metafluid with anisotropic viscosity

P2: Optical Rectification in Meta-Gratings with Broken Inversion Symmetry

Petr Moroshkin, Jimmy Xu

Brown University (USA)

We report on optical rectification effect generated by infrared absorption and diffraction in a periodic 1D meta-grating with a broken inversion symmetry in its unit cell. The photon-drag enabled effect in this case is substantially enhanced by surface plasmon polaritons, resulting in infrared driven ratchet transport of electrons.

P3: Plasmonic study of FeS₂/Au and FeS₂/Ag nanoparticles

R. Eyi¹, S. Lee¹, O. Manasreh¹, R. Rumpf²

¹University of Arkansas (USA), ²University of Texas at El Paso (USA)

Iron pyrite (FeS₂), FeS₂/Au and FeS₂/Ag nanoparticles were synthesized. The absorption and PL were measured before and after the coupling. No significant increase due to the plasmonic effects of the metallic particles were observed in the optical properties after coupling. finite difference frequency domain (FDFD) simulations were performed to further investigate the effects of the coupling. An increase in the optical properties was observed. Plasmonic peaks attributed to the semiconductor were also observed in the near infrared region

P4: Plasmons, Excitons, and Polaritons in Aligned Carbon Nanotubes

Weilu Gao, Junichiro Kono

Rice University (USA)

We have recently developed a controlled vacuum filtration technique to fabricate wafer-scale films of highly aligned and densely packed SWCNTs. Here, we summarize our recent accomplishments using these unique samples.

P5: Temporal Dynamics of Strongly-Coupled Epsilon Near-Zero Metasurfaces

Mehdi Haji Ebrahim¹, Andrea Marini², Vincenzo Bruno¹, Daniele Faccio¹, Matteo Clerici¹

¹University of Glasgow (United Kingdom), ²University of L'Aquila (Italy)

We demonstrate a significant slow-light effect in a deeply subwavelength epsilon near-zero plasmonic system

with decorated gold nanoantennae. We find that the longitudinal and transverse field components dominate the temporal response at the higher and lower resonant frequencies, respectively. We show that the slow-light effect is particularly pronounced within the strong coupling region and thus such ENZ plasmonic systems can provide an alternative platform for control and manipulation of light.

P6: Novel Photonic Glasses: Potassium Fluoride with Tellurium Oxide Doped With Rare Earth

Hamed Algarni¹, Norah A. Al Saffar¹, Manuela Reben², Kamal Dammak³, Mohammed Alqahtani¹, Elsayed Yousef¹

¹King Khalid University (Saudi Arabia), ²AGH - University of Science and Technology (Poland), ³Sfax University (Tunisia)

The glasses within composition in mol % 64TeO₂- 10WO₃- 10Nb₂O₅- 15KF- 1.0 La₂O₃-x Tm₂O₃ (where x= 0.0 (sample A), 1'104 (sample B), 2'104 (sample C), 3'104 (sample D) and 5'104ppm (sample E) were prepared. Spectroscopic properties of these glasses were estimated. The linear refractive index and third-order non-linear susceptibility increases with increase Tm³⁺ ions. The emission and gain cross-section increase with increasing Tm³⁺ ions in the host glass. These glasses can be fabricated as a promising laser material and optical devices.

P7: Luminescence and Radiation Shielding Parameters of Optical Halohalide Phosphate Glass with doping Rare Earth

Abdulrahman Hussain¹, Kamal Dammak², Mohammed Alqahtani¹, Hamed Algarni¹, Manuela Reben³, Elsayed Yousef¹

¹King Khalid University (Saudi Arabia), ²Sfax University (Tunisia), ³AGH - University of Science and Technology (Poland)

A homogeneity Pr³⁺ doped glass with the composition (in mol %) 40P₂O₅- 30ZnO- 20LiCl-10BaF₂-3.5Pr₂O₃ referred as PZLBP_r glass, was prepared by the melt quenching technique (see fig. 1). Herein the luminescence and shielding parameters such as radiative life time, branching ratio, magnetic dipole, mass attenuation coefficients (MAC), half-value layer (HVL), and mean free path, (MFP), were computed in range of ionizing radiation energies. These results indicate that the optical glass can be used in various diagnostic and therapy applications.

P8: Surface plasmon-assisted spin precession in Au/YIG heterostructures

Artsiom Kazlou¹, Alexander Chekhov², Alexander Stognij³, Ilya Razdolski¹, Andrzej Stupakiewicz¹

¹University of Bialystok (Russia), ²Free University Berlin (Germany), ³Scientific-Practical Materials Research Centre of the NASB (Belarus)

We report amplification of laser-induced spin precession in Co-doped YIG employing a surface plasmon excitation in a metal-dielectric magneto-plasmonic crystal. Our results are important for non-thermal control of all-optical magnetization reversal in dielectrics and its nanoscale localization.

P9: Terahertz near-field microscopy for quantitative measurements of the conductivity and charge carrier density on the nanoscale

Matthias M. Wiecha, Rohit Kapoor, Hartmut G. Roskos

Goethe University Frankfurt (Germany)

The suitability of a terahertz near-field microscope to quantitatively determine the conductivity and the charge carrier density of semiconductors is explored. For doped and optically excited silicon, the charge carrier density is successfully extracted from the relative phase of the terahertz nearfield signals. This technology is promising for non-contact and nanoscale-resolved characterization of electronic devices and materials.

P10: Curved structured beams: towards practical manipulation of light at the nanoscale from free space to in-plane plasmonic

Igor V Minin¹, Oleg V Minin²

¹Tomsk Polytechnic Univ. (Russia), ²Tomsk Polytechnic Univ. (Russia)

Among the ways of manipulation of light propagation, the interest is directed at the implementation of curving the trajectory of localized beam at the nanoscale in near field. It was discovered that an electromagnetic field can be made to curve after propagation through a simple dielectric material with broken symmetry. Photonic hooks features the radius of curvature, which is subwavelength - this is the smallest curvature radius of electromagnetic, acoustic and plasmonic waves ever reported.

P11: Influence of the Cell Number for finite Size Artificial Magnetic Conductor**Céline Ha¹, Jean-François Pintos¹, Priscillia Daquin², Serge Bories¹**¹Université Grenoble Alpes (France), ²CNES (France)

This paper examines the influence of the cell number for finite size AMC under normal plane wave incidence. A comparison is carried out on the operational frequency, the fractional bandwidth, and the reflection coefficient magnitude between the finite size AMC and an infinite structure, for different cell numbers and for three different finite size screens.

P12: Fabrication of SERS Substrates via Laser Induced Surface Nanostructuring of Silicon**Alp Akbryik, Nardin Avishan, Emre Yüce, Alpan Bek***Middle East Technical University (Turkey)*

In this work, we fabricate SERS substrates using laser assisted chemical etching as a surface nano structuring technique for silicon surface. On top of the etched surface, silver is thermally deposited to form hot spots with the silicon. Silver thickness on top of the silicon is shown to influence significantly to the SERS signal obtained from the substrates along with the various parameters that affect surface roughness of silicon.

P13: Purcell enhancement and photoluminescence spectra shift of the APbI₃ (A=Cs, FA) Perovskite nanocrystals coupled to Hyperbolic Metamaterials**Hamid Pashaei Adl¹, Setatira Gorji¹, Mojtaba Karimi Habil², Isaac Suarez¹, Vladimir S. Chirvony¹, Andrés F. Gualdrón-Reyes³, Ivan Mora-Sero³, Carlos J. Zapata-Rodríguez⁴, Juan P. Martínez-Pastor¹**¹Universidad de Valencia (Spain), ²University of Tabriz (Iran), ³Universitat Jaume I (Spain), ⁴Universitat de València (Spain)

Hyperbolic metamaterials are properly designed, simulated and fabricated as an outstanding photonic structure able to control the emission rate of lead halide perovskite nanocrystals (PNCs) deposited on the top. Geometrical parameters are optimized to enhance coupling between the structure and the exciton confined in the PNCs. The device is tested for CsPbI₃ and FAPbI₃ PNCs, and demonstrates an increase of the exciton radiative recombination rate around a factor of 2-3 together with the red shift of the emission spectra.

P14: Properties of Ni-doped ZnO thin films prepared by Sol-gel Spin Coating and Study of ZnO Powder Nanostructures.**Salah Oudjertli¹, Abdelkader Mohammedi², Miloud Ibrir²**¹Research Center In Industrial Technologies (Algeria), ²University of M'sila (Algeria)

Thin films of Nickel-doped ZnO are prepared by sol-gel spin coating process. The percentage concentration of dopant: Nickel, by weight, was at 0 %, 2 % and 5 %. Band structure and density of states of the phase of crystal ZnO computed using Ab Initio methods, confirmed that pure ZnO is a direct band gap semiconductor for B3 phase, whose phase B3 is of ZnS type Blende.

P15: Simulation of a Smart filter Based on Non-linear Metamaterials and Micro-plasma**Motahareh Rahmani, Saeed Hasanpour Tadi, Babak Shokri***Shahid Beheshti University (Iran)*

A smart photonic device is simulated to work as a limiter. A 2D non-linear metasurface is located inside a waveguide and its interaction with microwave by different input powers is studied. It is shown that when the input power is low, there is no micro-plasma and the system is transparent and in high input power mode, the situation is vice versa.

P16: Analysis of Overcoming Independent Core Light Propagation in Multicore Photonic Crystal fibers with Non-identical Cores Coupling**Miami Mohammed Mohammed***University of Muenster (Germany)*

Multicore photonic crystal fibers with non-identical cores are analyzed numerically using Comsol Multiphysics software. Anisotropy in all cores diameters of multicores photonic crystal fibers leads to different coupling behavior. Such anisotropy causes suppressed the coupling between the core modes at some wavelengths.

P17: Tunable polaritons enhanced by the spiral nanowire metamaterials**Thanos Ioannidis¹, Tatjana Gric¹, Edik Rafailov²**¹Vilnius Gediminas Technical University (Lithuania), ²Aston University (United Kingdom)

The tunable spiral nanowire metamaterial design at optical frequency is presented, and the surface polaritons

are theoretically studied. It was found that the dispersions of the polaritons could be tuned by varying physical dimensions of the spiral nanowire metamaterial. This geometry is unique. Doing so, one may dynamically control the properties of surface polaritons.

P18: All dielectric Si nanoresonator based color filters

Vishal Vashistha¹, Andriy Serebryannikov², Maciej Krawczyk²

¹Aalto University (Finland), ²Adam Mickiewicz University (Poland)

In this study, we experimentally demonstrate high-quality colors obtained by using arrays of cross-shaped Si nanoantennas. Functionality and characteristics achieved with the aid of symmetric and asymmetric nanoantennas are compared. Si nanoantennas typically show two strong resonances: electric and magnetic dipole ones. They can be tuned by adjusting the size and shape of the nanoantennas throughout the whole visible spectrum.

P19: Spoof Localized Surface Plasmons Examination Based on Effective Medium Theory in Textured Metallic Cylinders

Amirmasood Bagheri, Mohammad Reza Tavakol, Amin Khavasi

Sharif University of Technology (Iran)

Textured closed surfaces made out of perfect electric conductors (PECs) can mimic highly localized surface plasmons (LSPs). Here, we propose and scrutinize an effective medium being able to model LSP resonances in a two-dimensional periodically decorated PEC cylinder. Thanks to the incorporation of an effective surface conductivity created at the interface of the cylinder and the homogeneous medium surrounding the structure, the proposed model is accurate. Full-wave simulations verify the validity of the proposed model.

P20: Potential of pure VO₂ building blocks for tunable metasurfaces in the visible

Peter Kepič, Filip Ligmajer, Katarina Rovenska, Martin Hrtón, Jiri Liska, Tomas Sikola

Brno University of Technology (Czech Repub)

Metasurfaces represent a new class of optical components, which can provide optical functions far beyond the current applications. Phase-change materials can upgrade them into tunable metasurfaces. Vanadium dioxide (VO₂) represents a phase-change material, which can provide such tunability and which nanostructures were barely explored for metasurfaces in the visible. Here, we focus on the investigation of VO₂ nanocylinders in the visible. For that, we use FDTD simulations based on the refractive index obtained from our optimized VO₂ thin film.

P21: Numerical study of the optical properties of disordered metallic grooves by a one-mode analytical model

Denis Langevin, Julien Jaeck, Eslam El Shamy, Riad Haïdar, Patrick Bouchon

DOTA, ONERA, Paris-Saclay University (France)

Metallic grooves of sub-wavelength dimensions behave like Fabry-Perot nano-cavities able to resonantly enhance the electromagnetic field. In this summary, the consequences of positional disorder on the optical behavior of groove arrays are presented. We show, with a specifically developed simulation tool, that disorder leads to a redistribution of energy compared to the periodic case. By studying arrays of variously shaped grooves, we also show that their optical response can be described by the individual sub-array responses.

16:00 - 17:10 — Victor Veselago Room

Session 3A31

Plenary Session III

16:00 : Plenary talk

Applications of metasurfaces: From multispectral imaging to optical communications and biosensing

Maiken H. Mikkelsen

Duke University (USA)

In this talk, I will present advances in plasmonic metasurfaces with examples of potential future applications

in a wide-variety of areas from multispectral imaging to optical communications and biosensing.

16:35 : Plenary talk
4D Structured Waves

Nader Engheta

University of Pennsylvania (USA)

Varying materials parameters in time, in addition to (or instead of) spatial inhomogeneities in material platforms, can provide additional degrees of freedom in structuring and sculpting waves that lead to interesting functionalities in wave-matter interaction. While the spatial and/or temporal variations of material parameters have certain analogies and similarities, they exhibit important differences. In this talk, I will present an overview of some of the ongoing research programs on this topic in my group, will discuss the salient features, and will forecast possible future directions.

Break

17:10 - 17:30

17:30 - 19:00 — Victor Veselago Room

Session 3A32

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

17:30 : Keynote talk

Design and Knowledge Discovery in Metastructures Using Manifold Learning

Ali Adibi

Georgia Institute of Technology (USA)

New techniques for design and optimization of electromagnetic nanostructures using manifold-learning techniques are discussed. Using the strong correlation among features of an electromagnetic problem, deep-learning techniques are employed to considerably reduce the dimensionality of the problem and thus, the computation complexity, without imposing considerable error. Deep-learning algorithms can be trained to relate the reduced-dimensionality design and response spaces and facilitate the solution of the inverse design problems that are complicated to solve with conventional techniques. In addition, by training manifold-learning algorithms in the reduced-dimensionality spaces, valuable insights about the feasibility of the response and the roles of design parameters can be obtained. This talk explains the importance of these approaches and their applications to high-impact photonic nanostructures.

18:00 : Keynote talk

Volumetric microscale gradient refractive index lenses and waveguides for ultra-dense 3D optics

Paul V. Braun

University of Illinois Urbana-Champaign (USA)

Here, we present Subsurface Controllable Refractive Index via Beam Exposure (SCRIBE), a lithographic approach that enables the fabrication of volumetric microscale gradient refractive index lenses and waveguides. The basis of SCRIBE is multiphoton polymerization inside monomer-filled nanoporous silicon and silica scaffolds. Adjusting the laser exposure during printing enables 3D submicron control of the polymer infilling and thus the refractive index over a range of greater than 0.3 and chromatic dispersion tuning. Combining SCRIBE's unprecedented index range and 3D writing accuracy has realized the world's smallest (15 μm diameter) spherical Luneburg lens operating at visible wavelengths.

18:30 : Keynote talk

Active Epsilon-Near-Zero Photonics

Howard Lee¹, Aleksei Anopchenko¹, Sudip Gurung¹, Khant Minn², Jingyi Yang¹

¹University of California, Irvine (USA), ²Baylor University (USA)

This talk will review our recent development on conducting oxide and metallic nitride epsilon-near-zero optics. I will present our recent advances on the study of enhanced ultrafast nonlinearity and broadband and field-effect tunable absorption in ultrathin transparent conducting oxide ENZ materials meta-film fabricated by atomic layer deposition technique. In addition, I will discuss the photoluminescence enhancement of 2D materials on epitaxial titanium nitride thin films grown by molecular-beam-epitaxy. These studies enrich the fundamental understanding of emission and nonlinear properties on ENZ thin films that could be important for the development of advanced nanoscale lasers/light sources, optical/bio-sensors, and nano-optoelectronic devices.

17:30 - 19:00 — Allan Boardman Room

Session 3A33

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

17:30 : Invited talk

Photoelectrochemical Methanol Oxidation under Visible and UV Excitation of TiO₂-supported TiN and ZrN Plasmonic Nanoparticles

Olga Baturina¹, Albert Epshteyn¹, Jonathan A Boltersdorf², Gregory T Forcherio³, Asher C Leff⁴, Andrew Purdy¹, Blake Simpkins¹, Eva Y Santiago⁵, Alexander Govorov⁵

¹US Naval Research Laboratory (USA), ²US Army Research Laboratory (USA), ³Naval Warfare Center (USA), ⁴General Technical Services (USA), ⁵Ohio University (USA)

Combined experimental and computational approaches are used to optimize the performance of TiO₂-supported TiN and ZrN nanoparticles towards photoelectrochemical methanol oxidation. Issues leading to performance losses are analyzed and mitigation strategies are proposed.

17:50 : Invited talk

Non-Hermitian Approach for Modelling Hybrid Quantum Dot/Plasmon Systems

Cristian L. Cortes, Matthew Otten, Stephen K. Gray

Argonne National Laboratory (USA)

A non-Hermitian model for modelling quantum dot/plasmon interactions including dissipation and dephasing is analysed. Optical spectra in the linear regime are adequately described and the model also describes time-dependent coherences qualitatively when both dissipation and dephasing are present, and quantitatively with neglect of dephasing. Results for a large number of quantum dots allow assessing the role of coupling disorder. Interestingly, disorder can help stabilize the ensemble average towards a dark steady-state, a result of potential relevance to sensing applications.

18:10 : Invited talk

3D-Printed Terahertz Resonant Nanocones

Andrea Rovere¹, Riccardo Piccoli¹, Andrea Bertoni², Young-Gyun Jeong¹, Stéphane Payeur¹, François Vidal¹, O-Pil Kwon³, Seung-Heon Lee³, Roberto Morandotti¹, Carlo Liberale², Luca Razzari⁴

¹INRS - Énergie, Matériaux et Télécommunications (Canada), ²KAUST (Saudi Arabia), ³Ajou University (Korea), ⁴INRS - Énergie, Matériaux et Télécommunications (Canada)

Arrays of gold-coated 3D-printed nanocones resonating in the terahertz frequency range are investigated.

18:30 : DNA Based Optical Nano-sensor for Hazardous Molecules Detection

Sarabjeet Kaur, Safi Jradi, Julien Proust

Universite de Technologie de Troyes (France)

The prolonged use of toxic pesticides in agriculture lead to their introduction into the food chain which results in the interaction of these molecules with DNA and could create some mutations and induce translocation

leading to many diseases such as cancer. The aim of this research is to develop a unique Nano-biosensor which can detect chemical agents that interact and induce changes in structure of DNA. The detection is based on field enhancement from coupled gold nanoparticles (AuNPs) in solution.

18:45 : Light induced adaptation of structural colour and light absorption enhancement in photosynthetic photonic organelles

Miguel A. Palhinha Castillo, William P. Wardley, Martin Lopez-Garcia
International Iberian Nanotechnology Laboratory (Portugal)

In this work, we communicate a theoretical investigation on a naturally occurring photonic crystal: the iridoplast, an adapted photosynthetic organelle found in plants living under low light conditions. Our numerical study suggests that these structures could be controlling the absorption and the reflectance of light in order to enhance photosynthetic activity. We model purely light dependent structural changes based on experimental reports. This could therefore have potential interest in other technological areas such as coloured solar cells.

17:30 - 19:00 — Tatsuo Itho Room

Session 3A34

Plasmonics and nano-optics

17:30 : A Near-Perfect Optical Metamaterial Absorber and Its Applications

Mahdi Safari, Joel Y. Y. Loh, George V. Eleftheriades, Nazir P. Kherani
University of Toronto (Canada)

Here, we propose a metamaterial structure composed of Cu nano-cubes distributed on a Au-Cu bilayer thin film stack for near-perfect broadband optical absorption. We demonstrate *textgreater*95 % absorption in this structure over wavelengths ranging from 200 to 600 nm.

17:45 : Integrated Photodetection Leveraging Plasmonic Radiation Pressure

Jared H. Strait¹, Christian Haffner¹, Junyeob Song¹, Glenn Holland¹, Wei Zhou², Amit Agrawal³, Henri J. Lezec¹

¹National Institute of Standards and Technology (USA), ²Virginia Tech (USA), ³University of Maryland (USA)

We present novel integrated photodetectors based on the radiation pressure of a plasmonic mode. Light absorbed in a plasmonic or hybrid plasmonic/optical waveguide builds a voltage along the length of the waveguide via the photon-drag effect. We implement this device concept for the first time and investigate its potential for fast, broadband, and inexpensive optical detection in an integrated platform.

18:00 : Photoinduced Forces on Dielectric Nanoantennas

Martin Poblet¹, Yi Li², Gustavo Grinblat¹, Emiliano Cortés², Stefan Maier², Andrea Bragas¹

¹Universidad de Buenos Aires (Argentina), ²Ludwig-Maximilians-Universität München (Germany)

In this work we study optical forces produced by an all-dielectric germanium nanoantenna illuminated at resonance, through Photo-Induced Force Microscopy (PIFM). We investigate the electric and magnetic components of the optical forces and analyze the results through numerical simulations.

18:15 : Ultrafast all-optical modulation in dielectric nanoantennas at anapole resonances

Gustavo Grinblat¹, Michael Nielsen², Rodrigo Berté³, Haizhong Zhang⁴, Yi Li³, Leonid Krivitsky⁴, Benjamin Tilmann³, Emiliano Cortés³, Arseniy Kuznetsov⁴, Rupert Oulton⁵, Stefan Maier³

¹Universidad de Buenos Aires (Argentina), ²University of New South Wales (Australia), ³Ludwig-Maximilians-Universität München (Germany), ⁴A*STAR (Singapore), ⁵Imperial College London (United Kingdom)

In this investigation we study the ultrafast optical modulation performance of individual silicon and gallium phosphide (GaP) nanoantennas through pump-probe spectroscopy using sub-10 fs pulses. We find that a GaP nanoantenna excited at the anapole resonance outperforms its analog silicon resonator by two orders of magnitude, showing modulation depths close to 30 % at sub-30 fs time scales.

18:30 : Epitaxial Growth of Single Crystal Noble Metals for Plasmonic and Nanophotonic Applications

Sasan V. Grayli¹, Xin Zhang², finlay MacNab², Dmitry Star², Gary Leach², Michael Reimer¹

¹University of Waterloo (Canada), ²Simon Fraser University (Canada)

Plasmonic structures made from monocrystalline metals display lower absorption losses and much higher stability than polycrystalline devices which are subject to many losses due to the presence of grain boundaries. With the help of an epitaxial electroless chemistry, ultrasmooth gold films are grown on monocrystalline silver. Our approach enables the growth of gold with improved pattern transfer yield, reduced optical and resistive losses, and tailored local fields to yield greater optical response as compared to those of polycrystalline films.

18:45 : Width-graded plasmonic resonator arrays for broadband infrared spectroscopy

Katelyn Dixon¹, Arthur Montazeri², Liang Chen², Moein Shayegannia¹, Stefano Cabrini², Nazir Kherani¹, Hoi-Ying Holman²

¹University of Toronto (Canada), ²Lawrence Berkeley National Laboratory (USA)

Plasmonic field enhancement is used to amplify signals in various optical sensing techniques. However, the design of resonant devices in the infrared remains a challenge due to the high aspect ratios required and lack of repeatable fabrication methods. We present a facile technique for the fabrication of metal-insulator-metal (MIM) thin film resonator arrays suitable for field enhancement across the infrared range. Plasmonic absorption in these arrays is demonstrated through infrared spectroscopy and corroborated via simulation.

17:30 - 18:30 — Christian Huygens Room

Session 3A35

Metamaterials and metasurfaces

17:30 : A perfect lens for static magnetic fields

Rosa Mach-Battle¹, Mark G Bason², Nuria Del Valle¹, Jordi Prat-Camps²

¹Universitat Autònoma de Barcelona (Spain), ²University of Sussex (United Kingdom)

Perfect lenses are one of the most intriguing devices brought by negative refraction. In this work, we show that the recent introduction of negative permeability in magnetostatics enables the creation of a device analogous to a perfect lens in the magnetostatic limit. Experimental results demonstrate that this lens achieves a strong focusing of magnetic fields and makes it possible to create and cancel magnetic sources remotely, something unachievable with conventional devices with positive permeability.

17:45 : Microplotter printing of ultra-thin metasurfaces for the sub-terahertz domain

Cyprien Brulon, Arthur Salmon, Baptiste fix, Patrick Bouchon

ONERA (France)

Inkjet printing is a promising low-cost alternative to conventional optical lithography for the fabrication of terahertz metasurfaces. In this work, we discuss a new printing process and the fabrication of sub-terahertz metallic metasurfaces composed of periodical arrays of split ring resonators on ultra-thin substrate. The optical response in transmission of the metasurfaces shows resonances around 70 GHz, which is consistent with numerical simulations and demonstrates the use of microplotter system for rapid prototyping of low-loss terahertz passive components.

18:00 : A dielectric metasurface-polarimeter for single-shot detection of arbitrary polarization states

Yash Diptesh Shah, James P. Grant, Charles Altuzarra, Ashley Lyons, Daniele Faccio

University of Glasgow (United Kingdom)

We present a dielectric metasurface using an asymmetric design that exploits exotic multipolar interactions forming polarisation-dependent resonant eigenmodes. We experimentally demonstrate single-shot full-state polarimetry.

18:15 : Reconfigurable Terahertz Holographic CMOS Metasurface using Electromagnetic-Circuit Co-design Approach

Suresh Venkatesh, Xuyang Lu, Hooman Saeidi, Kaushik Sengupta

Princeton University (USA)

In this article, we demonstrate a digitally reconfigurable metasurface at 0.3THz exploiting the electromagnetic-circuit co-design approach. The metasurface is fabricated using industry-standard 65-nm CMOS process and the reconfiguration happens through digitally controlled CMOS switches. We also show scalability by chip tiling approach and also demonstrate multi-functionality (beamforming, holographic projections, spatial light modulation).

17:30 - 18:50 — Augustin Fresnel Room

Session 3A36

Light-matter interactions in new materials and meta-architectures

Organized by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

Chaired by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

17:30 : **Invited talk**

Ultra-Compact Photonic Modulator based on Accumulation-Layer Surface Plasmons

Viktoriia Babicheva

University of New Mexico (USA)

Most commonly studied electro-optic modulators employ several electro-optical mechanisms such as Pockels, Kerr, or carrier concentration change effects to modulate the phase and amplitude of light propagating through the device. In this work, we consider a transparent conducting oxide (TCO) based electro-optic modulator and demonstrate the propagation of surface plasmons due to the accumulation of carriers in the thin TCO layers under external electrical bias.

17:50 : **Invited talk**

Photonics with Metallic Alloys

Marina S. Leite

University of California (USA)

We propose metallic alloys and intermetallic materials as a platform to control the electromagnetic spectrum in the UV-NIR range. Opportunities for developing optical components using metallic materials beyond noble metals for applications including superabsorbers, NIR photodetectors, and catalysis, among others will be discussed.

18:10 : **Invited talk**

Polar Semiconductors as Long-Wavelength Epsilon-Near-Zero Materials

Milan Palei, Irfan Khan, Owen Dominguez, Junchi Lu, Ryan Roeder, Anthony Hoffman

University of Notre Dame (USA)

Polar semiconductors exhibit a region of negative permittivity between the longitudinal optical (LO) and transverse optical (TO) phonons. At energies close to the LO phonon energy, the real part of the optical permittivity approaches zero and the imaginary part is also very small, the so-called epsilon-near-zero (ENZ) spectral regime. We show how the ENZ and negative permittivity regions can be leveraged for a new generation of long-wavelength optical devices and materials.

18:30 : **Invited talk**

Diagnostic Colorimetric Metasurfaces Visualize Disease in fibrous Biological Tissue

Lisa V. Poulikakos¹, Zaid Al Haddadin¹, Trinity C. Pike¹, Jebin J. Moses¹, Aniket S. Puri¹, Mark Lawrence², David R. Barton³, Stefanie S. Jeffrey⁴, Jennifer A. Dionne⁴

¹University of California (USA), ²Washington University St. Louis (USA), ³Harvard University (USA), ⁴Stanford University (USA)

We leverage the unique properties of anisotropic, colorimetric metasurfaces to scale down the complex manipulation of light and selectively visualize disease-relevant fiber density and orientation in biological tissue. Ranging from Alzheimer's disease to heart disease, fibrosis or cancer, we discuss the potential of metasurfaces to yield rapid, precise, low-cost diagnostics.

17:30 - 19:00 — Ibn Al-Haytham Room**Session 3A37****Quantum and topological photonics****17:30 : Optical Magnetism in a Quantum System without Metamaterials****Kyle Ballantine, Janne Ruostekosk***Lancaster University (United Kingdom)*

We propose how to synthesize optically active magnetism without metamaterials using quantum-mechanical electric dipole transitions of naturally occurring atoms. We extend the method for toroidal dipoles and anapoles, and a physical realization of a quantum Huygens' surface that allows for extreme wavefront engineering even at a single photon level. Examples of focusing and steering light, and generations of entangled quantum superposition states with additional photons are provided.

17:45 : Majorana-like bound state in Kekule distorted sonic lattices**Penglin Gao¹, Daniel Torrent², Francisco Cervera³, Pablo San-Jose⁴, Jose Sanchez-Dehesa³, Johan Christensen¹***¹Universidad Carlos III de Madrid (Spain), ²Universitat Jaume I (Spain), ³Universitat Politècnica de Valencia (Spain), ⁴Instituto de Ciencia de Materiales de Madrid (ICMM-CSIC) (Spain)*

Topological phases have recently been unveiled in bosonic systems to facilitate the manipulation of waves in unprecedented ways. Here we demonstrate a new type of matter supporting non-propagating localized state that is topologically protected and robust against any parameter perturbations given that the intrinsic particle hole symmetry is preserved, i.e., an acoustic analogue of Majorana bound state. Our work might broaden exciting avenues for robust sound confinement and energy harvesting in industries.

18:00 : Integration of nitrogen-vacancy center into an one-dimensional photonic crystal cavity**Jan Olthaus¹, Philip P. J. Schrinner², Carsten Schuck², Doris E. Reiter¹***¹University of Munster (Germany), ²Center for NanoTechnology CeNTech and Center for Soft Nanoscience SoN (Germany)*

Photonic crystal cavities based on on-substrate tantalum pentoxide waveguides are optimised for coupling to nitrogen vacancy centers in nanodiamond using 3D-FDTD simulations. Coupling conditions depending on the position and size of the nanodiamonds are studied. Antibunching of the photoluminescence signal of an integrated nitrogen-vacancy center in nanodiamond coupled to a 1D photonic crystal cavity is observed experimentally.

18:15 : Spontaneous emission enhancement of handed molecules**Karolina Slowik, Miriam Kosik***Nicolaus Copernicus University (Poland)*

A theory of interactions of nanostructured quantum vacuum with atomic systems beyond the electric dipole approximation has recently been developed. Based on that input, in this work we investigate spontaneous emission enhancement of right- and left-handed molecules via the Purcell effect in proximity of plasmonic or dielectric nanocavities. Our conclusion is that the handedness of an atomic system may have major influence on its emission rate in suitably tailored nanophotonic environments.

18:30 : Dispersion engineering for an ultraviolet frequency comb**Ali E. Dorche¹, Dogan Timucin², Krishnan Thyagarajan², Noble Johnson², Thomas Wunderer², David E. Schwartz²***¹Georgia Institute of Technology (USA), ²Palo Alto Research Center (PARC) Inc. (USA)*

Anomalous dispersion engineering in the ultraviolet (UV) spectrum is used for efficient bright soliton-based optical frequency comb generation. We have overcome normal dispersion in a III-Nitride platform to create a wideband Kerr frequency comb, with a free spectral range of 400GHz, centered at 442 nm, to ensure that the Raman gain does not compete with the four-wave mixing gain. This paves the way for applications of fine UV spectral lines in quantum computing, metrology, optical clocks, inertial navigation and secure communication technologies.

18:45 : Lamb-Dicke Confinement of Cold Atoms in Ferris Wheels

V. E. Lembessis, A. Lyras, O. M. Aldossary

King Saud University (Saudi Arabia)

We investigate the case of strong confinement of a Cs cod atom that is trapped in an optical Ferris wheel light field. We show that it is possible to reach the Lamb-Dicke limit for parameter values that are typically used in cold atom physics experiments.

17:30 - 18:50 — Gaston Floquet Room

Session 3A38

Topology in photonic crystals, metamaterials, and metasurfaces: physics and design

Organized by: Yang Li and Benfeng Bai

Chaired by: Yang Li and Benfeng Bai

17:30 : Invited talk

Inverse design of metasurfaces and photonic systems for enhanced Raman scattering

Rasmus E. Christiansen¹, Jérôme Michon², Ying Pan², Ole Sigmund¹, Juejun Hu², Steven G. Johnson²

¹Technical University of Denmark (Denmark), ²Massachusetts Institute of Technology (USA)

We propose and investigate designs for nano-patterned surfaces and devices, tailored to maximize the surface enhanced Raman scattering (SERS) occurring from molecules placed near them. The designs are created using our extended version of a recently proposed approach, utilizing topology optimization as an inverse design tool. The modified approach takes additional limitations in the fabrication process into account in order to minimize the discrepancies between the design blueprint and the fabricated metasurface.

17:50 : Invited talk

Inverse Design Towards Global Performance Bounds

Owen Miller

Yale University (USA)

We demonstrate the capability to design structures approaching fundamental nanophotonic performance limits. We use inverse design in tandem with analytical and computational bound techniques to identify opportunities for improvement across various applications, and discover designs with state-of-the-art performance. We design minimal-thickness perfect absorbers, maximal-efficiency grating couplers, and minimal-mode-volume dielectric resonators (where we show the capability to incorporate fabrication and multi-frequency constraints). More broadly, we discuss general principles for identifying nanophotonic limits and designing devices to approach them.

18:10 : Invited talk

T Operator Bounds for Electromagnetic Power Transfer

Sean Molesky¹, Pengning Choa¹, Prashanth S. Venkataram¹, Weiliang Jin², Alejandro W. Rodriguez¹

¹Princeton University (USA), ²Stanford University (USA)

We present a method for utilizing power transfer constraints on the electromagnetic scattering operator to set physical bounds on any single material design problem that can be framed as a net emission, scattering or absorption process. The technique is found to predictively quantify and differentiate the relative performance of dielectric and metallic materials for both far and near-field sources. The broad applicability of scattering theory means that similar application to acoustics, quantum mechanics, and other wave physics are likely possible.

18:30 : Invited talk

Multi-Layered Meta-Optics by Inverse Design

Zin Lin, Steven G. Johnson

Massachusetts Institute of Technology (USA)

To successfully address the issues besetting the state-of-the-art LPA-based metasurface designs, we pro-

pose a novel computational approach with three key components: multiple interacting layers with deeply sub-wavelength separations (three-dimensional single-piece nanophotonics) designed by large-scale optimization techniques (inverse design and topology optimization), using more sophisticated overlapping domain-decomposition methods superior to LPA.

17:30 - 19:00 — Lawrence Bragg Room

Session 3A39

Nonlinear metamaterials

17:30 : All-Optical Switching of an Epsilon-Near-Zero Plasmon in ITO

Justus Bohn¹, Craig Tollerton¹, Ting Shan Luk², Igal Brener², Bill Barnes¹, Euan Hendry¹

¹University of Exeter (United Kingdom), ²Sandia National Laboratories (USA)

A new class of materials with a vanishing permittivity, known as epsilon-near-zero (ENZ) materials, has been reported to exhibit unprecedented ultrafast nonlinear efficiencies within subwavelength propagation lengths. We study the pump dependent near perfect absorption ENZ plasmon in a thin indium tin oxide (ITO) layer. Utilising the Kretschmann configuration combined with the ENZ plasmon resonance could pave the way towards ultrafast switching from near-perfect absorption to total internal reflection or vice versa.

17:45 : Negative refraction in time-varying, strongly-coupled plasmonic antenna-ENZ systems

Vincenzo Bruno¹, Clayton DeVault², Stefano Vezzoli³, Zhaxylyk Kudyshev², Tahiyat Huq³, Sandro Mignuzzi³, Andrea Jacassi³, Soham Saha², Yash Diptesh Shah¹, Stefan Maier⁴, David Cumming¹, Alexandra Boltasseva², Marcello Ferrera⁵, Matteo Clerici¹, Daniele Faccio¹, Riccardo Sapienza³, Vladimir Shalaev²

¹University of Glasgow (United Kingdom), ²Purdue University (USA), ³Imperial College London (United Kingdom), ⁴Ludwig-Maximilians-Universität München (Germany), ⁵Heriot-Watt University (United Kingdom)

We demonstrate high efficiency in the generation of optical induced time-reversal phase conjugate and negative refraction waves, from a temporal modulated deeply subwavelength epsilon-near-zero (ENZ) film integrated within a plasmonic metasurface. The strong coupling between the plasmonic resonance and the ENZ modes leads to a conversion efficiency that is more than 4 orders of magnitude greater than the bare ENZ film.

18:00 : Interaction based nonlinearity of a deep subwavelength plasmonic heterodimer

Avi Niv

Ben-Gurion University of The Negev (Israel)

Conventionally, optical nonlinearity emerges from the linear fields that echo the linear optical response. Contrarily, we demonstrate a pure second-harmonic resonance, without a linear counterpart, and propose a dynamic model of its origin.

18:15 : Modelling the Thermo-Optic Non-Linear Behavior of 2D Photonic Crystal Cavities

Simone Iadanza¹, Marco Clementi², Changyu Hu³, Sebastian A. Schulz⁴, Dario Gerace², Matteo Galli², Liam O'Faolain¹

¹Cork Institute of Technology (Ireland), ²Università di Pavia (Italy), ³University College Cork (Ireland), ⁴University of St. Andrews (United Kingdom)

We present a first principles model for the thermo-optic nonlinearities of PhC cavities. The match between the calculations and experiments demonstrated here renders the model a crucial predicting tool for the development of nonlinear microcavities

18:30 : Exploiting time-dependent metamaterials for frequency conversion in guided-wave structures

Victor Manuel Pacheco Peña¹, Nader Engheta²

¹Newcastle University (United Kingdom), ²University of Pennsylvania (USA)

In this work we study theoretically the effect of using time-dependent metamaterials filling the cladding of optical waveguides and ring resonators as a technique to effectively change the frequency of the wave traveling in such guided-wave structures in real time. The physics behind using such temporal metamaterial clads is

discussed and demonstrated numerically at telecommunication wavelengths.

18:45 : A quasi-planer approach to dynamic meta-surfaces via band-gap engineered, electro-optic quantum effects

Walter R Buchwald

University of Massachusetts Boston (USA)

finite element simulations motivate placement of a 2-level quantum well within the near surface depletion region of a Schottky contact producing a quantum system of three-dimensionally confined hetero-interface charge allowing both voltage and optical control of electron occupancy. This electro-optic quantum system is discussed in relation to dynamic nano-photonics metasurfaces.

17:30 - 18:40 — Rene Descartes Room

Session 3A40

Structured and topological photonic fields

Organized by: Jamal Berakdar

Chaired by: Jamal Berakdar

17:30 : Invited talk

Matter and optical vortices and their interactions

M. Babiker¹, V. E. Lembessis², Koray Kooksal³, J. Yuan¹

¹University of York (UK), ²King Saud University (Saudi Arabia), ³Bitlis Eren University (Turkey)

This talk deals with optical and matter vortices, singly and interacting. The ability to generate optical matter vortices, suggests scenarios where vortex atoms interact with vortex photons. These lead to new physics, most notably exchange of orbital angular momentum, spin-orbit coupling where the longitudinal optical field component comes into play. We highlight novel situations with chiral trapping potentials due to bi-chromatic doughnut beams. The spin-orbit interaction and the longitudinal field components are emphasised to be crucial in this context.

17:50 : Invited talk

Structured ultrafast high-harmonic pulses

Carlos Hernández-García

Universidad de Salamanca (Spain)

Coherent extreme-ultraviolet/soft x-ray pulses structured in their temporal (attosecond), spectral (line spacing) and angular momentum (polarization and topological charge) properties is nowadays possible thanks to high harmonic generation. In this talk we review our recent work in the generation and control of ultrafast structured harmonic pulses through the use of infrared driving beams with custom angular momentum properties.

18:10 : Photonic simulation of Rashba-Dresselhaus spin-orbit coupling in a tunable birefringent cavity

Katarzyna Rechcinska¹, Mateusz Krol¹, Rafal Mazur², Przemyslaw Morawiak², Rafal Mirek¹, Karolina Iempicka¹, Witold Bardyszewski¹, Michal Matuszewski³, Przemyslaw Kula², Wiktor Piecek², Pavlos G. Lagoudakis⁴, Barbara Piętko¹, Jacek Szczytko¹

¹University of Warsaw (Poland), ²Military University of Technology (Poland), ³Polish Academy of Sciences (Poland), ⁴University of Southampton (United Kingdom)

Rashba and Dresselhaus effects well known in semiconductor physics have inspired the field of spintronics by providing a means of spin manipulation. In this communication we will demonstrate how these phenomena can be simulated in a purely photonic system. Experimental spectra of a tunable liquid-crystal-based device will be presented opening the avenue for novel technical solutions which exploit the analogy between the electron's spin and photon's polarization.

18:25 : Three-dimensional vectorial holography based on machine-learning inverse design

Haoran Ren¹, Wei Shao², Yi Li³, Flora Salim², Min Gu⁴

¹Ludwig-Maximilians-Universität München (Germany), ²RMIT University (Australia), ³Southern University of Science and Technology (China), ⁴University of Shanghai for Science and Technology (China)

We demonstrate 3D vectorial holography where an arbitrary 3D vectorial field distribution on a wavefront can be precisely reconstructed using the machine-learning inverse design based on multilayer-perception artificial neural networks. Such 3D vectorial holography allows the lensless reconstruction of a 3D vectorial holographic image with near-unity 3D polarization purity. Holographic information can thus be encoded and encrypted on the wavefront of a 3D vectorial field.

18:00 - 19:00 — Tutorial Room

Session 3A41

Conference Tutorials III

Organized by: Ishwar Aggarwal

18:00 : Tutorial

New Artificial-Intelligence Techniques for Electromagnetic Metasurfaces

Ali Adibi

Georgia Institute of Technology (USA)

A survey of new artificial-intelligence-based approaches for analysis, design, optimization, and knowledge discovery in electromagnetic metasurfaces will be presented. Recent advances in using both deep learning and machine learning techniques, and their application to practical problems will be covered. These techniques will not only enable more efficient designs of the electromagnetic metasurfaces (e.g., photonic metasurfaces) but also provide valuable insight about the complex physics of light-matter interactions in such structures. Details of the training process for these algorithms as well as the challenges and limitations of these techniques for different classes of metasurfaces will be discussed. Knowledge discovery using these techniques includes the study of feasibility of a certain optical response from a given class of metasurfaces and comparing the roles of different design parameters to facilitate the inverse design process.

Friday 23rd July, 2021

09:00 - 10:55 — Victor Veselago Room

Session 4A1

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

09:00 : **Biomimetic Ultra-Broadband Perfect Absorbers Optimised with Reinforcement Learning**

Trevon Badloe, Inki Kim, Junsuk Rho

Pohang University of Science and Technology (POSTECH) (Korea)

A double deep Q-learning network (DDQN) is used to design ultra-broadband, biomimetic, perfect absorbers with various materials, based the structure of a moth's eye. By training a DDQN with chromium-based design, we transfer the learned knowledge to other, similar materials to quickly and efficiently find the optimal parameters from the ~ 1 billion possible options. The previously learned knowledge helps the network optimise new materials in fewer steps, dramatically increasing the efficiency of finding the best designs for ultra-broadband absorption.

09:15 : **Invited talk**

Near-field heat flux control with many thermal emitters: theory and applications

Philippe Ben-Abdallah

Laboratoire Charles Fabry (France)

Near-field radiative heat transfer between two bodies is an old problem in physics. In 2011, the theoretical foundations have been introduced to deal with near-field heat exchanges in many-body systems. In this presentation I will introduce the theoretical framework to describe this physics and i will discuss some of its potential applications.

09:35 : **Invited talk**

Arbitrary Order Exceptional Point Induced by Photonic Spin-Orbit Interaction

Shubo Wang¹, Bo Hou², C. T. Chan³

¹City University of Hong Kong (China), ²Soochow University (China), ³The Hong Kong University of Science and Technology (China)

Exceptional points in non-Hermitian systems have many intriguing properties and novel applications. High order exceptional points normally require delicate variations of multiple parameters and are difficulty to achieve. In this talk, we will show that photonic spin-orbit interaction can induce exceptional points of arbitrary order without tuning any parameters.

09:55 : **Invited talk**

Compact and Integrated Imaging Based on Metalenses

Tao Li, Chen Chen, Beibei Xu, Shining Zhu

Nanjing University (China)

Recent progresses in metasurface have opened a door of new type of ultra-thin optical elements for versatile applications. Here, I would like to show our recent progress in the spectral tomographic imaging based on aplanatic metalens, which was implemented by sweeping the wavelength the tuning the focal length and achieved and DOF resolved microscopic imaging. Afterwards, an integrated device has been constructed by mounting metalens directly onto a CMOS sensor to access a compact, stable, and wide-field microscopy.

10:15 : **Invited talk**

Highly efficient nanophotonic color router for sub-micron-pixel CMOS imagers

Hongkyu Park, Sookyoung Roh, Sangyun Lee, Minwoo Lim, Hyuck Choo, Seokho Yun

Samsung Advanced Institute of Technology (Korea)

We have demonstrated novel nanophotonic color-routing structures that can efficiently split and focus different

colors onto corresponding pixels of an image sensor. Direct color-separating capability without the use of absorptive filters provides each pixel with more photons of the desired wavelength, resulting in higher signal-to-noise ratios.

10:35 : Invited talk

Metasurfaces with Maxwell's demon-like nonreciprocity

Kin Hung Fung

The Hong Kong Polytechnic University (China)

09:00 - 10:10 — Allan Boardman Room

Session 4A2

Symposium I: Hybrid Nanomaterials and Metastructures for Photonics, Sensing and Energy

Organized by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

Chaired by: Jérôme Plain, Alexander Govorov, Davy Gérard and Pedro Hernandez Martinez

09:00 : Invited talk

Extracting Meaning from the Analysis of Photoluminescence Decays of Colloidal Quantum Dots: What's in a Function?

Ana Luisa Simões Gamboa¹, Evgeny N. Bodunov²

¹ITMO University (Russia), ²Emperor Alexander I Petersburg State Transport University (Russia)

We discuss functions for the description of the room-temperature photoluminescence decay of colloidal quantum dots that are practical to use and whose parameters have a straightforward physical meaning. We introduce a function that accounts for the long-time tails of the decays, highlighting processes that may be related to photoluminescence blinking, and which use can provide valuable information concerning the nature of the trap states involved in the recombination of charge carriers.

09:20 : Invited talk

High quality factor Tamm structures for the development of laser sources at room temperature

Clementine Symonds, Vincent Toanen, Jean-Michel Benoit, Alban Gassenq, Joel Bellessa

Université de Lyon (France)

Tamm structures offer very versatile approaches to develop confined lasers, polarized lasers or plasmon sources. We propose here an optimized design of these structures, enabling an increase by a factor 5 of their quality factor, which is a critical parameter for various applications. In particular, we will show that these optimized structures enable room temperature lasing operation. This first demonstration is an important step toward future applicative developments of Tamm devices.

09:40 : Surface-enhanced Absorption Principle of Thin film Sensing in the Mid-infrared

Dmytro Chubich, Roman Zvagelsky, Danila Kolymagin, Anastasia Pisarenko, Elena Zhukova

Moscow Institute of Physics and Technology (Russia)

We fabricated arrays of periodic nanoantennas with plasmonic resonances between 1300 cm⁻¹ and 2300 cm⁻¹ on a single chip and demonstrated the ability of infrared signal enhancement on specific analyte with vibrational lines close to resonances of fabricated nanoantennas.

09:55 : Single molecule studies of metal-enhanced fluorescence and resonance energy transfer interplay in graphene-metallic hybrid nanostructure

Kamil Wiwatowski¹, Karolina Sulowska², Sebastian Mackowski², Joanna Niedziolka-Jönsson¹

¹Polish Academy of Sciences (Poland), ²Nicolaus Copernicus University (Poland)

In this work, we use single molecule fluorescence spectroscopy to probe nanoscale interactions in hybrid nanostructures composed of graphene and silver nanowires. The presence of metallic nanoparticles or graphene in hybrid nanostructure can induce fluorescence enhancement, fluorescence quenching or their interplay in nearby emitter, depending on nanostructure arrangement.

10:10 - 10:55 — Allan Boardman Room**Session 4A3****Photonic bandgap structures****10:10 : Hybrid External Cavity Laser based on Silicon Nitride 1D Photonic Crystals Cavities for optical sensing in gasses and liquids****Simone Iadanza, A. Tedesco, G. Giannino, M. Grande, L. O'Faolain***Cork Institute of Technology (Ireland)*

In this paper we demonstrate a 1D silicon nitride photonic crystal cavity for optical sensing operating in the NIR wavelength range, in liquids and gasses.

10:25 : Vertical assisted directional-coupler from Silicon-on-Insulator to Silicon Nitride platforms**L. Zagaglia¹, S. Argiolas², S. Iadanza¹, L. O'Faolain¹, G. Mura², F. Floris¹, P. O'Brian¹**¹*Tyndall National Institute (Ireland)*, ²*DIEE-University of Cagliari (Italy)*

Silicon (Si) and Silicon Nitride (Si₃N₄) are two leading materials for the creation of photonic integrated circuits (PICs). The necessity of efficiently couple light between Si-PICs and Si₃N₄-PICs is becoming more relevant to face crucial applications in various technologic sectors, as high bandwidth optical interconnects and mid-infrared and visible optical sensing. However, the strong refractive index mismatch between Si and Si₃N₄ leads to challenging coupling between the elements used to carry the light throughout the PICs (waveguides). We propose a solution based on a vertical assisted directional-coupler allowing a theoretical -2.2dB insertion-loss and a 30nm bandwidth.

10:40 : Near-field optical investigation of Hyperuniform Disordered photonic structures**Nicoletta Granchi¹, Richard Spalding², Matteo Lodde³, Maurangelo Petruzzella³, Frank Wan Otten³, Andrea Fiore³, Francesca Intonti¹, Riccardo Sapienza⁴, Marian Florescu², Massimo Gurioli¹**¹*University of Florence (Italy)*, ²*University of Surrey (United Kingdom)*, ³*Eindhoven University of Technology (The Netherlands)*, ⁴*Imperial College London (United Kingdom)*

Located in-between random structures and perfectly ordered photonic crystals, there is a special class of disordered heterostructures called hyperuniform disordered (HuD) photonic structures. These materials, due to the presence of a photonic bandgap, combine the advantages of disordered systems and ordered systems: here, we underline and experimentally prove all these advantages by means of the first near-field optical characterization of HuD photonic structures in the near IR.

09:00 - 11:00 — Tatsuo Itho Room**Session 4A4****Advances in Metamaterials**

Organized by: Jeong Weon Wu, Bumki Min and Jonghwa Shin

Chaired by: Jeong Weon Wu, Bumki Min and Jonghwa Shin

09:00 : Invited talk**Strain Responsive Elastomeric Metamaterials Made from PnP****Seokwoo Jeon***KAIST (Korea)*

We propose a new type of three-dimensional (3D) nanocomposite film consisting of an ultrathin (~60 nm) alumina nanoshell inserted between the elastomers in a perfectly periodic 3D nanonetwork. Numerous light-scattering nanogaps form at the interfaces of the alumina and the elastomers under stretching. This results in the uniform modulation of transmission from ~90% to 16% at visible wavelengths and maintains its performances during repeated stretching/releasing over more than 10,000 cycles thanks to the perfect uniformity of the 3D structure.

09:20 : Invited talk

Elastic Metasurfaces tailoring reflections and refractions

Min Soo Kim, Sung Won Lee, Joo Hwan Oh

Ulsan National Institute of Science and Technology (Korea)

In this presentation, recent researches on elastic metasurfaces, thin artificial surface that can tailor elastic waves as desired, will be introduced. First, the reflection-type elastic metasurface will be focused, and a new idea of transmodal elastic metasurface will be introduced. After that, the refraction type elastic metasurface that can control the transmitted waves with a single doubly-negative unit will be explained. From these basic results, various new applications such as vibration and ultrasonic devices are expected.

09:40 : Invited talk

Metasurface for on chip nanophotonics

Soo Jin Kim

Korea University (Korea)

Metasurface has been drawn great scientific and practical interests due to its potential applications in novel optical devices. In this work, we build the designing strategy of metasurfaces for the applications in chip scale devices which feature effective spectral sorting on the nano-scaled photodetection systems. Furthermore, we show the electrical tuning of metasurface for the dynamic control of optical spin using the transparent conductive oxide combined with plasmonic metasurfaces.

10:00 : Invited talk

Soft Matters for Extremely High-Index Meta-Optics

Seungwoo Lee

Korea University (Korea)

Conventionally, meta-optics has benefitted from the rapid advances in semiconducting processing such as lithography and lift-off/etching of hard materials. Here, I'll introduce that it is time to widen our view of materials and relevant processing for meta-optics. Especially, colloidal self-assembly can extremely increase the polarization of effective medium and resultant refractive index.

10:20 : Invited talk

Cyclic group symmetric metasurface for optical spin-dependent beam separation

Yeon Ui Lee¹, Igor Ozerov², Frederic Bedu², Ji Su Kim¹, Frederic Fages², Jeong Weon Wu¹

¹*Ewha Womans University (Korea)*, ²*Aix-Marseille University (France)*

Geometric phase is introduced when a cross-polarization from nano-rod takes place. Here we introduce a cyclic symmetric metasurface composed of tapered arc nano-rods and explore how azimuthal angular dependence determines the feature of spin-dependent beam separation.

10:40 : Invited talk

On-Demand Electrical Tuning of Metasurface with Complex Modulation for Arbitrary Wavefront Shaping

Junghyun Park, Byung Gil Jeong, Sun Il Kim, Duhyun Lee, Jisan Lee, Inoh Hwang, Hyuck Choo, Kyoungho Ha

Samsung Advanced Institute of Technology (Korea)

We present an ultrafast, electrically-tunable, 50-channel metasurface array that can generate arbitrary phases and amplitudes of light in reflection and achieves on-demand beam steering. The individual metasurface of the demonstrated array consists of an index-modulating indium tin oxide layer sandwiched between the individually addressable top and bottom electrodes. Applying separate biases to the electrodes modulates the charge concentration in the ITO layer, which allows independent control of the complex reflection coefficient and consequently the phase and amplitude of reflected light.

09:00 - 10:35 — Christian Huygens Room

Session 4A5

Metasurfaces and 2D Metamaterials in microwave region

Organized by: Badreddine Ratni and Shah Nawaz Burokur

Chaired by: Badreddine Ratni and Shah Nawaz Burokur

09:00 : Invited talk**Metasurface Hologram with High Quality in Microwave Region****Xumin Ding, Tianhao Wu, Chunsheng Guan, Zhuochao Wang, Kuang Zhang, Qun Wu***Harbin Institute of Technology (China)*

Metasurfaces have shown great potential in controlling the propagation of light through the well-tailored scattering behavior of the constituent ultrathin planar elements with a high spatial resolution, making them suitable for holographic beam-shaping elements. Here, we review recent developments in the field of metasurface holography in microwave region.

09:20 : Invited talk**Independent Manipulation of Orthogonal Circular Polarizations based on Microwave Metasurface****Yueyi Yuan¹, Kuang Zhang¹, Xumin Ding¹, Badreddine Ratni², Shah Nawaz Burokur², Qun Wu¹***¹Harbin Institute of Technology (China), ²Université Paris Nanterre (France)*

In this paper, a transmissive-type metasurfaces based on multi-layer structure is proposed to independently manipulate the orthogonal circularly polarized wave in microwave region. Through combining the propagation phase and geometry phase principles, the opposite circularly polarized transmitted wave can be imposed independent spatial phase distributions to perform different wavefronts. Experimental measurement are conducted and effectively verified the feasibility of the proposed theory for artificial manipulation of circular polarization manipulation in microwave region.

09:40 : Invited talk**Angular Scattering Control with Multilayer Metasurface Stacks****Karim Achouri, Olivier J. F. Martin***EPFL (Switzerland)*

In order to implement metasurface-based optical analog processing systems, we develop an analysis technique for computing the angular scattering response of multilayer metasurface stacks. This technique is based on an improved scattering matrix method and applies to bianisotropic metasurfaces for optimal field control.

10:00 : Invited talk**Glide-Symmetric Luneburg Lens Based on Substrate-Integrated-Holes****Oskar Zetterstrom, Ramez Hamarneh, Oscar Quevedo-Teruel***KTH Royal Institute of Technology (Sweden)*

We propose a novel parallel plate waveguide (PPW) metasurface. The PPW is periodically loaded with glide-symmetrically arranged dielectric-filled cavities in both conductors. By filling the cavities with dielectric, the equivalent refractive index increases. To facilitate the manufacturing using conventional methods, the walls of the cavities are implemented with metallic vias and the proposed structure is used to design a Luneburg lens. Compared to previous works on holey metasurfaces, the designed lens is cheaper and more resilient to manufacturing errors.

10:20 : Design Concept of Dual-Polarized Microwave Superscatterers**Vitalii Shcherbinin¹, Volodymyr Fesenko¹, Vladimir Tuz²***¹National Academy of Sciences of Ukraine (Ukraine), ²Jilin University (China)*

It is shown that a subwavelength perfectly-electrically-conducting (PEC) cylinder with doubly periodic dielectric-filled slots features anisotropic surface impedance, which may ensure superscattering of both TEz- and TMz-polarized waves at a single frequency. Owing to this property, a slotted cylinder is demonstrated to operate as a dual-polarized superscatterer. A prototype of such superscatterer is designed to provide a broadband dual-polarized superscattering in the microwave range.

10:35 - 11:05 — Christian Huygens Room**Session 4A6****Metasurfaces, flat optics, FSS and HIS****10:35 : Efficient Mie Resonance of Metal-masked Titanium Dioxide Nanopillars****Xiao Shang, Lina Shi, Jiebin Niu, Changqing Xie***Chinese Academy of Sciences (China)*

Here, we propose a simple design approach based on metal-masked titanium dioxide nanopillars, which can realize strong Mie resonance in metasurfaces and enables light confinement within itself over the range of visible wavelengths. By selecting the appropriate period and diameter of individual titanium dioxide nanopillars, the coincidence of resonance peak positions derived from excited electric and magnetic dipoles can be achieved. And the optical properties in this design have been investigated with the finite-Difference Time-Domain (FDTD) solutions.

10:50 : Harnessing the Optical Performance of Metalenses with Numerical Aperture**Haowen Liang, Juntao Li***Sun Yat-sen University (China)*

We present a crystalline silicon based metalens with effectively low loss and high-NA in air. This metalens can be further front-immersed in high index liquids, presenting much higher NA and diffraction-limited focusing.

09:00 - 10:00 — Augustin Fresnel Room**Session 4A7****Organic and Perovskite Optoelectronics**

Organized by: Erik Lotfi

09:00 : Invited talk**Colloidal inorganic halide perovskite quantum dots optoelectronics****Chenghao Bi, Jifeng Yuan, Shixun Wang, Jianjun Tian***University of Science and Technology Beijing (China)*

Colloidal inorganic halide perovskite CsPbX_3 ($x=\text{Cl, Br, I}$) quantum dots (QDs) show great promising for high-performance photovoltaics and electro-optical converters owing to outstanding optoelectronic properties, such as potential multi-excitons generation, tunable band gap, high carrier mobility and narrow emission width. However, they are still suffering from low power conversion efficiency (PCE) and poor stability, which are hindering their practical application. In this talk, I will demonstrate our recent works that aimed at solving the issues above.

09:20 : Invited talk**Unique Room-Temperature Solution-Processes for High Performance Organic and Perovskite Optoelectronic Devices****Wallace Choy***The University of Hong Kong (Hong Kong)*

Room-temperature solution process will be discussed for various optoelectronic devices fabrication, while high-temperature evaporation and sputtering are commonly used. Room-temperature-processed metal-oxides-based carrier transport layers are attractive candidates for all-solution-processed optoelectronics such as organic solar cells, perovskite solar cells, dye-sensitized solar cells, and organic light-emitting diodes, which can favor the efficient transport of carriers between the photoactive layer and electrode as well as optical transparency. Additionally, new class of heavy-duty room-temperature-processed Ag nano-network is developed as efficient transparent flexible electrodes.

09:40 : Invited talk

Hole Transporting Materials with Strategy of Flexible Core and Tunable Conformation for Efficient and Stable Perovskite Solar Cells

Aung Ko Ko Kyaw

Southern University of Science and Technology (China)

I will present three HTMs based on this strategy, employing flexible saddle-shaped cyclooctatetrathiophene or bifluorenylidene as core and triphenylamine units as arms. Both cyclooctatetrathiophene and bifluorenylidene more flexible than rigid spiro unit in Spiro-OMeTAD and fit well into our strategy. The influence of the location of arms on the dihedral angles, molecular configuration, packing characteristics and the resultant film morphology will be discussed. In addition to HTMs, the passivation of perovskite with organic molecular materials will be briefly discussed.

10:00 - 11:00 — Augustin Fresnel Room

Session 4A8

Synthesis and characterization of plasmonic nanostructures

Organized by: Mohamed Boutinguiza Larosi and Juan Pou Saracho

Chaired by: Juan Pou Saracho

10:00 : Invited talk

Plasmonics in a Variable Temperature Thermodynamic Bath

Francesco Bisio¹, Michele Magnozzi², Marzia Ferrera², Maurizio Canepa²

¹CNR-SPIN (Italy), ²Università di Genova (Italy)

Electromagnetically-heated metal nanoparticles can be exploited as efficient heat sources at the nanoscale. The assessment of their temperature is, however, often performed indirectly by modelling their temperature-dependent dielectric response. Direct measurements of the optical properties of metallic nanoparticles in equilibrium with a thermodynamic bath provide a calibration of their thermo-optical response, to be exploited for refining current thermoplasmonic models or whenever direct temperature assessments are practically unfeasible.

10:20 : Invited talk

Synthesis, characterization and example applications of anisotropic plasmonic metal nanoparticles

Jan Krajczewski

University of Warsaw (Poland)

In some metallic nanoparticles, incident light of an appropriate frequency excites the collective oscillation of electron plasma, a phenomenon known as surface plasmon resonance. Electron plasma oscillations lead to a high local enhancement of the electromagnetic field near the plasmonic structure which generates increase in the effectivity of many optical processes. In this contribution various methods of synthesis of such structures and examples of their applications (in spectroscopy, catalysis, cancer detection and treatment) will be presented.

10:40 : Invited talk

Colloid- and Polymer-Based Self-Assembled Meta-atoms and Metasurfaces

Rajam Elancheliyan¹, Cian Cummins¹, Alberto Alvarez-Fernandez², Philippe Barois¹, Alexandre Baron¹, Olivier Mondain-Monval¹, Guillaume Fleury³, Virginie Ponsinet¹

¹Centre de Recherche Paul Pascal (France), ²University College London (United Kingdom), ³Univ. Bordeaux (LCPO-CNRS) (France)

We present some examples of meta-atoms and metasurfaces fabrication processes, which implement a bottom-up approach based on colloid and polymer directed self-assembly. Scalability and tunability of the processes are demonstrated, as well as the final optical properties reached by these self-assembled nanostructures.

09:00 - 10:15 — Ibn Al-Haytham Room

Session 4A9

Optical antennas and plasmonics-based devices

09:00 : Confined Orbit Angular Momentum in Deep-subwavelength Plasmonic Resonator**Xuanru Zhang, Tie Jun Cui***Southeast University (China)*

Chiral symmetry broken is introduced into the double-layer microwave plasmonic resonator, to realize confined orbit angular momentum (OAM) in deep-subwavelength scale. Electromagnetic field and chirality distributions are analyzed quantitatively. OAM mode within $1/11$ free space wavelength has been demonstrated, exhibiting superchiral fields and envisioning the potential in detecting chiral structures.

09:15 : Self-powered photoresponse enhanced by asymmetrically integrated optical patch antennas in a metal-graphene-metal structure**Jing Zhou, Shangkun Guo, Jie Ding, Yu Yu, Wei Lu, Xiaoshuang Chen***Chinese Academy of Sciences (China)*

Optical patch antennas are integrated into the metal-graphene-metal structure in an asymmetrical manner for a prominent self-driven photoresponse. A 105 times high contrast between the photoresponses at the two contact-graphene junctions is achieved, and the responsivity enhancement by this structure is one order of magnitude higher than that by a subwavelength metal grating. The resonant behavior of the antenna enables spectrum-selective photoresponse. The photoresponse time is shorter than several microseconds, and the photoresponse mechanism is mainly attributable to photothermoelectric effect.

09:30 : Independent engineering of individual plasmon modes in plasmonic dimers with conductive and capacitive coupling**Vlastimil Krapek¹, Michal Horak¹, Martin Hrton¹, Andrea Konecna², Michael Stoger-Pollach³, Filip Ligmajer¹, Tomas Sikola¹***¹Brno University of Technology (Czech Republic), ²Materials Physics Center CSIC-UPV/EHU (Spain), ³Vienna University of Technology (Austria)*

We revisit plasmon modes in nanoparticle dimers with conductive or insulating junction. In our study which combines electron energy loss spectroscopy, optical spectroscopy, and numerical simulations, we show co-existence of strongly and weakly hybridized modes. While the properties of the former ones strongly depend on the nature of the junction, the properties of the latter ones are nearly unaffected. This opens up a prospect for independent engineering of individual plasmon modes in a single plasmonic antenna.

09:45 : Angular emission of second harmonic from GaP antennas reveals multipolar modes**Pawel Wozniak, Vikas Remesh, Niek F. van Hulst***ICFO-Institut de Ciències Fotoniques (Spain)*

An SHG interaction can be enormously enhanced using resonant antennas due to their ability to confine light within sub-wavelength volume. Our investigation demonstrates a great influence of the antenna size on the SHG efficiency and, more importantly, on the SHG angular emission. Detailed modal studies reveal the excitation of an antenna mode via the nonlinear interaction, which is not allowed in the linear regime, and a nontrivial conversion of the polarization state between these two regimes.

10:00 : Subwavelength mapping of optical modes in all-dielectric nanoantennas**Aleksandr Yu. Frolov¹, Niels Verellen², Jiaqi Li², Xuezhi Zheng², Hanna Paddubrouskaya², Denitza Denkova³, Maxim R. Shcherbakov¹, Guy A. E. Vandenbosch², Vladimir I. Panov¹, Pol Van Dorpe², Andrey A. Fedyanin¹, Victor V. Moshchalkov²***¹Lomonosov Moscow State University (Russia), ²KU Leuven (Belgium), ³Macquarie University (Australia)*

We report on near-field mapping of higher order optical modes in all-dielectric nanoantennas (rod, disk, square, triangle shape) by using aperture type scanning near-field optical microscopy (SNOM). Nanoantennas have been made of amorphous Si on the glass substrate. The correspondence of the electromagnetic field components of the excited optical modes with features on SNOM maps is demonstrated by means of full 3D FDTD modeling of the scanning process.

10:15 - 11:00 — Ibn Al-Haytham Room**Session 4A10****Plasmon-enhanced photovoltaics, photocatalysis, and solar fuels****10:15 : High-Index Optical Metamaterial for Perovskite Solar Cells****Kwangjin Kim, Seungwoo Lee***Korea University (Korea)*

Recently, perovskite solar cells have attracted considerable attentions from the solar cell society due to their exotic properties such as high quantum efficiency and low recombination rate. However, the power conversion efficiency (PCE) of the perovskite solar cells, reported thus far (~25 %), is not yet comparable to their fundamental limit (referred to as Shockley-Queisser Limit). Here, we show that an increase in refractive index of perovskite via high-index optical metamaterial can further enhance the PCE of solar cell.

10:30 : Ultraviolet radiation impact on the efficiency of crystalline silicon-based photovoltaics**George Perrakis, Anna C. Tasolamprou, George Kenanakis, Eleftherios N. Economou, Stelios Tzoratzakis, Maria Kafesaki***FORTH-IESL (Greece)*

We evaluate the ultraviolet radiation impact on the temperature and efficiency of realistic photovoltaic modules. We perform this investigation for crystalline silicon-based photovoltaics that operate outdoors, by employing a thermal-electrical modeling approach, which takes into account all the major intrinsic processes affected by the temperature variation in the photovoltaic devices.

10:45 : Near-field spectroscopy of Silicon Carbide phonon polaritons resonators**Andrea Mancini¹, Christopher R. Gubbin², Rodrigo Berte¹, Alberto Politi², Yi Li³, Simone De Liberato², Stefan A. Maier¹***¹Ludwig-Maximilians-Universität München (Germany), ²University of Southampton (United Kingdom), ³Southern University of Science and Technology (China)*

Antennas supporting surface phonon polaritons are an interesting alternative to common plasmonic resonators in the infrared region, due to their reduced losses and higher field confinement. However, many applications involving SPhPs antennas require knowledge of their near-field response, which cannot be directly inferred from common far-field measurements. We study the near-field spectral response of arrays of Silicon Carbide antennas by means of scattering scanning near field microscopy and discuss the influence of the AFM tip on the experimental results.

09:00 - 10:50 — Gaston Floquet Room**Session 4A11****Metasurface for information processing**

Organized by: Jensen Li and Xianzhong Chen

Chaired by: Jensen Li and Xianzhong Chen

09:00 : Invited talk**SP3: Metasurface for information processing****Lingling Huang¹, Qunshuo Wei¹, Basudeb Sain², Yongtian Wang¹, Bernhard Reineke², Xiaowei Li¹, Thomas Zentgraf²***¹Beijing Institute of Technology (China), ²Paderborn University (Germany)*

We propose and experimentally demonstrate a novel meta-device that integrates color printing and computer-generated holograms within a single-layer dielectric metasurface by modulating spectral and spatial responses at subwavelength scale, simultaneously. In our design, such metasurface appears as a microscopic color image under white light illumination, while encrypting two different holographic images that can be projected

at the far-field when illuminated with red and green laser beams. Such a method can further extend the design freedom of metasurfaces.

09:20 : Reconfigurable Optical Switch based on Metasurface-Coated Liquid Crystal Elastomer

Jianxun Liu¹, Xiaoguo Fang¹, Ming Cheng¹, Changxiong Zheng¹, Jiawei Wang¹, Hao Zeng², Yanjun Liu¹

¹*Southern University of Science and Technology (China)*, ²*Tampere University of Technology (Finland)*

Liquid crystal elastomer, whose ordered arrangement of mesogens can be changed under light stimulation and leading reversible mechanical deformation, have emerged as an intensively explored soft material in recent years. In this work, we introduce an active optical switch with broadband working and all-optical control using a metasurface-coated liquid crystal elastomer. As a reconfigurable substrate, LCE film deforms by photothermal effect. The results demonstrate a low-cost technique to realize complex and reconfigurable light-driven optical devices of active metasurfaces.

09:35 : Measuring Both Phase and Polarization Singularities of Light with Plasmonic Spin-Hall Nano-gratings

Changjun Min

Shenzhen University (China)

Measuring the phase and polarization singularities of light is essential for modern optics, but it is still a challenge especially in integrated optical systems. Here, we propose an on-chip plasmonic spin-Hall nano-grating structure that individually or simultaneously detects the polarization and phase singularities of incident light. We experimentally demonstrated that the nano-grating structure enables unidirectional excitation of surface plasmon polariton depending on the topological charge of optical vortex, and shows chiral responses for vectorial polarization state detection.

09:50 : Anomalous Brewster Effects and Ultra-broadband Reflectionless Brewster Absorbers Protected by Reciprocity

Jie Luo¹, Jensen Li², Yun Lai³

¹*Soochow University (China)*, ²*The Hong Kong University of Science and Technology (Hong Kong)*, ³*Nanjing University (China)*

The Brewster's law predicts zero reflection of p-polarization on a dielectric surface at a particular angle. However, when loss is introduced into the permittivity of the dielectric, the Brewster condition breaks down and reflection unavoidably appears. Here, we found an exception by creating a class of nonmagnetic anisotropic metamaterials, where anomalous Brewster effects with independently tunable absorption and refraction emerge. Ultra-broadband reflectionless Brewster absorbers with bandwidth covering from dc to optical frequencies, are bestowed by the anomalous Brewster effects.

10:05 : Polarized Light Interference Imprinted Plasmonic Metasurface

Zhen Yin, Yue Shi, Jianxun Liu, Huilin He, Xiaoguo Fang, Dan Luo, Yan Jun Liu

Southern University of Science and Technology (China)

An approach for direct fabrication of silver nanoparticles structured plasmonic metasurface imprinted by two orthogonal circularly polarized light interference is presented. The correlation between the spatial distributed silver nanoparticles and the state of polarization of the interference field is investigated. The proposed approach can be further extended to create even more complex and unconventional plasmonic metasurfaces by tailoring the polarization interference.

10:20 : Anticounterfeiting visible metaholograms multiplexed with spin, direction and wavelength

Junsuk Rho, Inki Kim

Pohang University of Science and Technology (POSTECH) (Korea)

Computer-generated holography (CHG) involves iterative numerical algorithms to obtain the phase and/or amplitude profiles needed to physically realize holograms. Metasurfaces consist of arrays of subwavelength nanoresonators that can control the wavefront of light in a desired way.

10:35 : Coherent unitary transformations of light using metasurfaces

Ming Kang, Kai Ming Lau, Tsz Kit Yung, Shengwang Du, Wing Yim Tam, Jensen Li

Hong Kong University of Science and Technology (China)

We propose to combine coherent and wavefront control with metasurfaces to carry out designer unitary

transformations on SU(2) and SU(4). The developed framework will be useful for carrying out mathematical operations and quantum information processing.

09:00 - 10:55 — Lawrence Bragg Room

Session 4A12

Light-matter interactions in new materials and meta-architectures

Organized by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

Chaired by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

09:00 : **Invited talk**

Phase change reprogrammable visible photonics

Li Lu¹, Parikshit Moitra², Ramon Paniagua-Dominguez², Vytautas Valuckas², Sander F. G. Reniers³, Harish N. S. Krishnamoorthy⁴, Arseniy I. Kuznetsov², Yuqing Jiao³, Cesare Soci⁴, Robert E. Simpson¹

¹Singapore University of Technology and Design (Singapore), ²A*STAR (Singapore), ³Eindhoven University of Technology (The Netherlands), ⁴Nanyang Technological University (Singapore)

Phase change materials (PCMs) play an important role in tuneable photonics applications. Here, we designed reprogrammable hyperbolic metamaterials, nanoantenna beam-steering metasurfaces, and tuneable waveguide devices that operate at visible and near infrared wavelengths. All three devices exploit the large change in refractive index associated with the amorphous-crystalline phase transition in antimony trisulphide (Sb₂S₃). This particular phase change material has a 2 eV band gap, which deems it transparent to visible and near infrared light.

09:20 : **Invited talk**

Hexagonal boron nitride for integrated quantum photonics

Sejeong Kim¹, Johannes E. Fröch², Ngoc My Hanh Duong², Igor Aharonovich²

¹University of Melbourne (Australia), ²University of Technology Sydney (Australia)

Integrated quantum photonic circuitry is an emerging topic that requires efficient coupling of quantum light sources to waveguides and optical resonators. Here, we present the hBN quantum emitters integrated into photonic waveguides and photonic cavities.

09:40 : **Collective response of liquid crystal based metasurfaces**

Raviteja R. Mosali¹, Madhunika Atmakuri¹, Sushanth R. Amanaganti¹, Miha Ravnik², Jayasri Dontabhaktuni¹

¹Mahindra Ecole Centrale (India), ²University of Ljubljana and Jozef Stefan Institute (Slovenia)

In our earlier work, we studied the collective response of high refractive index cubic Te metasurface using full-wave electromagnetic simulations. In the present work, we investigate the effect of anisotropy of lattice constant and anisotropy of the medium on the collective response and further, we also study the effect of quasicrystalline symmetry on the electromagnetic response of Penrose tiling made of pentagonal shaped sub-micron dielectric structures.

09:55 : **Invited talk**

Metal-Insulator-Metal cavities for light enhancement and modulation

Vincenzo Caligiuri¹, Aniket Patra¹, Giulia Biffi², Renuka Devi Pothuraju², Antonio De Luca¹, Roman Krahne²

¹University of Calabria (Italy), ²Istituto Italiano di Tecnologia (Italy)

Metal-Insulator-Metal (MIM) nanocavities that sustain epsilon-near-zero (ENZ) resonances constitute a versatile platform for light modulation and amplification. They act as optical quantum wells for photons, and an analogy to quantum mechanics facilitates the analytic prediction of the resonances. Multiple cavities lead to coupled resonances, and we demonstrate how a MIMIM system can enhance both the absorption and emission of a dye. Our approach can be extended to ENZ bands, which can be described by the Kronig-Penney model.

10:15 : **Invited talk**

Controlling light with magneto-optical gratings

Ioan-Augustin Chioar¹, Merlin Pohlit¹, Christina Vantaraki¹, Richard Rowan-Robinson¹, Bjorgvin Hjorvarsson¹, Evangelos Papaioannou², Vassilios Kapaklis¹

¹Uppsala University (Sweden), ²Martin-Luther-Universität (Germany)

We report on the use of a continuous magnetic film of YIG for the control of light with mesoscale magnetic textures. We investigate in real and reciprocal space, the connection between the magnetic domain structure developed and the resulting scattering pattern of the transmitted light.

10:35 : **Invited talk**

Active tuning of thermal radiation in the far-field and near-field range with emerging low-dimensional materials

Georgia Theano Papadakis

ICFO - The Institute of Photonic Sciences (Spain)

Controlling the flow of a thermal current is of critical importance for any application requiring thermal regulation. Previous approaches focused largely on modulating thermal conductance via tailoring the propagation of acoustic phonons. In contrast, in this talk, I consider means of controlling the flow of heat mediated by thermal radiation. I will discuss how leveraging the unique properties of low-dimensional materials like graphene and monolayer hexagonal boron nitride yields significant tuning of radiative heat transfer.

09:00 - 11:00 — **Rene Descartes Room**

Session 4A13

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

09:00 : **Invited talk**

Dynamic Control of Terahertz Polarization Based on Babinet Inversion of Anisotropic Metasurfaces with Vanadium Dioxide

Toshihiro Nakanishi¹, Yosuke Nakata², Yoshiro Urade³, Kunio Okimura⁴

¹Kyoto University (Japan), ²Osaka University (Japan), ³RIKEN (Japan), ⁴Tokai University (Japan)

We propose anisotropic metasurfaces, which realize dynamic polarization control by switching between two states of the structures. We have to design only one of the two states linked through Babinet's principle, because the other state automatically satisfies the required conditions. We demonstrate a reconfigurable polarizer and a reconfigurable quarter-wave plate in terahertz regions utilizing insulator-to-metal transition of vanadium dioxide in the metasurfaces. The polarization control can be realized by switching the local conductivity of the vanadium dioxide.

09:20 : **Invited talk**

Photo thermal energy conversion in mid infrared metasurfaces

Yoshiaki Nishijima, Naoki To

Yokohama National University (Japan)

Mid infrared wavelength is unique to apply the detection of the small amount of molecules with molecular vibration or rotational modes. Especially absorption type of meta surface can be used for the mid infrared light source, detectors according to the Kirchhoff's thermal radiation law. Here we summarized the recent progress of plasmonic meta surfaces in mid infrared wavelength region. The experimentally measured optical properties were compared with simulations by finite difference time-domain calculations. Also, we demonstrate applications of these structures for the plasmonic IR-light sources and detectors and another sensing devices.

09:40 : **Invited talk**

Metamaterial-based Production of Acoustic Orbital Angular Momentum in Three- and Two-dimensional Systems

Bin Liang¹, Jianchun Cheng¹, Jing-jing Liu¹, Xue Jiang¹, Yifan Zhu¹, Xuefeng Zhu², Yong Li³, Yun Jing⁴, Likun Zhang⁵, Chengwei Qiu⁶

¹Nanjing University (China), ²Huazhong University of Science and Technology (China), ³Tongji University (China), ⁴North Carolina State University (USA), ⁵University of Mississippi (USA), ⁶National University of Singapore (Singapore)

In this talk, I will review the recent advance in metamaterial-based production of acoustic orbital angular momentum (OAM) in three-dimensional systems with a planar and thin device and in two dimensional systems with an irregularly-shaped and compact enclosure.

10:00 : Invited talk

Plasmonic nanolasers modulated by current on graphene-insulator-metal structures

Tien-Chang Lu, Heng Li, Zhen-Ting Huang

National Chiao Tung University (Taiwan)

We propose a plasmonic nanolaser on graphene-insulator-metal structures and experimentally demonstrate the current modulated lasing behavior. Graphene serves as a two-dimensional material with high electron mobility, which is beneficial to external current injection. When the current is applied, it is obvious to observe the peak intensity dissipation and blue-shifted of the lasing signal. This work exhibits the great potential in active plasmonic devices.

10:20 : Invited talk

Optical Waveguide Using Off- Γ Bound States in the Continuum in One Dimensional Grating Structures

Yuto Moritake, Masaya Notomi

Tokyo Institute of Technology (Japan)

We proposed and numerically investigated filtering waveguides using off- Γ bound states in the continuum (BICs) in one dimensional grating structures. By using propagating feature of off- Γ BIC, light transmission is guided only around the BIC frequency without radiation leakage. The designed structure is composed of Si grating structure integrated with optical waveguide. Numerical simulations confirm the proposed filtering function at off- Γ BIC frequency.

10:40 : Invited talk

High-Performance Laser Based on Topological Photonic Crystal

Xiao Hu

NIMS (Japan)

We find a new light confinement mechanism emerging from photonic topology, namely the parity of eigen modes in a topological photonic crystal is different from those in a trivial photonic crystal below the band edge, which prevents light to transport freely across the interface. We then develop a high-performance laser with vertical cavity surface emission, which meets the IEEE and VCSEL standards.

Coffee Break and Exhibit Inspection

Session 4P1

Poster session VII

11:00 - 11:40

P1: The idea and technical realization of the structure for the observation of inverse Faraday effect.

Piotr A. Drozd, Valentin Kachorovskii, Pawel Prystawko, Mateusz Slowikowski, Maciej Filipiak, Dmitri Yavorski, Maria Szola, Wojciech Knap

CENTERA Laboratories (Poland)

In this work, the idea of the observation of Inverse Faraday Effect (IFE) is presented. The mechanism of the effect is described and the main conclusions from theoretical calculations of IFE in two dimensional electron gas (2DEG) are presented. finally, the practical realization of the structure for IFE observation is presented. The structure is based on the GaN/AlGaIn HEMT structure with back - gate layer. The processing as well as the basic characterization of the structure will be presented.

P2: Spherically-shaped WGM resonators for lasers and biodetection

Piotr Paszke¹, Rafal Nowaczynski², Hancza Barbara Surma¹, Piotr Piotrowski¹, Kamil Szlachetko¹, Joel Bellessa³, Clementine Symonds³, Vincent Toanen³, Andrea Csaki⁴, Dorota Anna Pawlak¹

¹University of Warsaw (Poland), ²Warsaw University of Technology (Poland), ³Université Claude Bernard Lyon 1 (France), ⁴Leibniz Institute of Photonic Technology (Germany)

This work presents the results of doping glasses with both silver nanoparticles and quantum dots, fabrication of WGM resonators based on these active glasses and their properties. The authors are mainly interested in applications of such resonators as biosensors, therefore the biofunctionalization tests of glass surfaces was also done.

P3: Photonic liquid crystal fibers with gold nanoparticles-doped cubic blue phases for enhanced electric field tunability and thermal stability

Kamil Orzechowski¹, Wiktor Lewandowski², Olga Strzezysz³, Martyna Tupikowska², Tomasz R. Wolinski¹

¹Warsaw University of Technology (Poland), ²University of Warsaw (Poland), ³Military University of Technology (Poland)

In this work, optical properties of photonic crystal fibers infiltrated with gold nanoparticles-doped cubic blue phases liquid crystals are demonstrated. It is presented that the investigated complex photonic systems can provide promising tunable properties for particular wavelengths in the visible light spectrum. Moreover, the presence of gold nanoparticles with an appropriate organic coating in a liquid crystal matrix can enhance the external electric field sensitivity and temperature stability of the examined photonic liquid crystal fibers.

P4: Dynamic and chiral plasmonics using freezing-directed self-assembly of nanoparticles

Wiktor Lewandowski

University of Warsaw (Poland)

Nanostructures with chiral symmetry and plasmonic properties attract considerable interest. Here, we present endeavors towards achieving thin films of chiral, reconfigurable nanostructures. This approach relies on mixing liquid crystals with gold nanoparticles, melting and freezing composites. Due to the soft character of liquid crystals, these nanoarchitectures exhibit helical, hierarchical and switchable structure. With this method we achieved durable, configurable, chiral plasmonics systems in the bulk.

P5: Plasmonic Enhancement of Second-Harmonic Generation with film-Coupled Nanopatch Antennas
Ahsan Noor¹, Anoop R. Damodaran², In Ho Lee², Stefan Alexander Maier³, Sang Hyun Oh², Antonella D'Orazio⁴, Cristian Ciraci⁵

¹Politecnico Di Bari (Italy), ²University of Minnesota, Minneapolis (USA), ³Ludwig-Maximilians Universitat Munchen (Germany), ⁴Politecnico di Bari (Italy), ⁵Istituto Italiano di Tecnologia (Italy)

Field enhancements associated with resonant excitation of plasmonic structures have shown tremendous potential to improve the efficiencies of nonlinear wave-mixing processes at the nano-scale. In this work, we present an optimal mode-matched second-harmonic generation from hybrid plasmonic film-coupled nanopatch antennas.

P6: Local intensity enhancement and Purcell factor in hyperbolic metamaterials - spontaneous emission engineering

Karol Sielezin, Alessandro Pianelli, Michal Dudek, Marek Olifierczuk, Rafal Kowrdziej, Janusz Parka
Military University of Technology (Poland)

In this work, we theoretically study the quantum mechanisms that are an effective tool for analyzing the increase in the photonic density of states (PDOS) in relation to hyperbolic metamaterials (HMMs). Numerical results of the Purcell factor of HMM stack built on the basis of alternating layers of silica and silver are reported. We theoretically show that interplay of plasmon polaritons leads to an increase in the spontaneous emission and, as a result, to an increased PDOS.

P7: All-dielectric metasurface linear polarizer for visible wavelengths utilizing the first Kerker condition

Amr Mohamed Soliman, Calum Williams, Timothy Wilkinson

University of Cambridge (United Kingdom)

Polarizers are indispensable elements integrated within a myriad of optical setups across many applications, from imaging assemblies to display technology. Commercially available polarizers are typically characterized by a low damage threshold, low extinction coefficient, and low operating bandwidth. In this paper, we present

an all-dielectric metasurface linear visible polarizer with high extinction ratio. The polarizer consists of sub-wavelength titanium dioxide (TiO₂) structures which utilize Kerker's first condition in order to generate the desired transmission response across the visible spectrum. In addition, the reported all-dielectric polarizer has a high damage threshold and can be tuned to different wavebands through geometry modification.

P8: Hyperuniform Disorder for Tailored Light Scattering Metasurfaces

Peter M. Piechulla¹, Bodo Fuhrmann¹, Evgeniia Slivina², Carsten Rockstuhl², Ralf B. Wehrspohn¹, Alexander N. Sprafke¹

¹Martin Luther University Halle-Wittenberg (Germany), ²Karlsruhe Institute of Technology (Germany)

Nanoparticle arrays exploited for scattering commonly feature either periodic or random arrangements. For the periodic case, scattering is governed by spatial correlations, expressed by the structure factor. For the random case, structural correlations cancel each other out and scattering is governed by properties of the individual scatterer, expressed by the form factor. In contrast to these extremes, hyperuniform disorder enables both structure and form factor to impact the resulting scattering pattern, offering novel means to tailor light scattering.

P9: Localization of Surface Acoustic Waves in 2D Disordered Phononic Crystals Using Pump and Probe Spectroscopy

Thibault Deletang¹, Bernard Bonello¹, Rock Akiki², Bahram Djafari-Rouhani², Yan Pennec², Eric Lheurette²

¹Université Pierre et Marie Curie (France), ²Université de Lille (France)

We report here some experimental results and reflections on localization. Some advancements have been made on the sample fabrication and the experimental set-up which allows us to study which allows us to study the dynamics of structure vibration related to the propagation of a surface wave in a disordered medium using pump and probe spectroscopy.

P10: Light amplification by silver nanoparticles surface plasmon resonance

Jelena Mikelsona, Aivars Vembris

University of Latvia (Latvia)

Organic materials have several advantages in the fields of an organic light-emitting diode organic photovoltaics, etc., but in the same time they have lower stability and efficiency in comparison to inorganic compounds. It can be improved by introducing metallic nanoparticles in the organic media. In this work we study silver nanoparticles synthesis in aqueous solution, their transfer to organic solutions, their impact on luminescent material's photoluminescence quantum yield.

P11: High refractive index dielectric rings in liquid crystals as tunable metasurfaces

Madhunika Atmakuri, Pratiksha A. Sakhare, Sushanth R. Amanaganti, Jayasri Dontabhaktuni

Mahindra University (India)

In the present work, we study the resonant characteristics of high refractive index (RI) circular and square rings using full wave electromagnetic simulations. Effect of geometry, Thickness of structures and RI on the EM response is studied in the sub-wavelength regime. This rings when placed in birefringent liquid crystal medium orientation of liquid crystal molecules are observed to enhance certain modes over the other giving rise to tunable metasurfaces. Also, electric dipole, quadrupole and toroidal modes are observed in air.

P12: Resonance characteristics of sub-wavelength high refractive index dielectric metasurfaces as a function of lattice constant

Jayesh Sadasivan¹, Sivarama Krishnan², Jayasri Dontabhaktuni¹

¹Mahindra University (India), ²Indian Institute of Technology Madras (India)

In our earlier work, we had studied the collective response of high refractive index dielectric Te metasurfaces of small periodicity using electromagnetic simulations. In this work, we perform a detailed investigation of the response spectrum as a function of increasing periodicity. We observe the suppressions of electric dipolar modes, observation of novel modes like hybrid modes and BIC. Some of the states observed at the smaller periodicity regimes were found to be independent of the lattice constant.

P13: Observation of localized states induced by curved acoustic topological insulators

Hong-Wei Wu, Jia-Qi Quan, Yun-Kai Liu, Yi-Ming Pan

Anhui University of Science and Technology (China)

In this work, we design a two-dimensional curved acoustic topological insulator by perforating on a curved

rigid plate. We experimentally demonstrate that a topological localized state stands erect in the bulk gap, and the corresponding pressure distributions are confined at the position with the maximal curvature. The robustness of the localized state is demonstrated by introducing defects near the localized position. The interaction between the geometrical curvature and topology provides a novel scheme for manipulating and trapping wave propagation.

P14: Giant enhancement of chiral selective many-body correlation among emitters coupled with spirally stacked metal structures

Hirofumi Shiraki¹, Nobuhiko Yokoshi¹, Hajime Ishihara²

¹Osaka Prefecture University (Japan), ²Osaka Prefecture University and Osaka University (Japan)

We have studied the anomalously enhanced cooperative effect by metal structures sustaining the localized surface plasmons. In this contribution, we consider chiral spatial structures of the metal and emitter arrangements, and calculate the emission intensity of superfluorescence. The results show the remarkable difference by chirality in emission intensity, correlation between the emitters, and polarization distribution. We can expect that these results will lead to a new methodology in chiral research.

P15: Resonance-order dependent PIT in orthogonally arranged nanoscale cavities

Naoki Ichiji, Atsushi Kubo

University of Tsukuba (Japan)

We investigate the Plasmonic Induced Transparency (PIT) in a resonator structure consisting of two orthogonally arranged metal-insulator-metal (MIM) nanocavities by FDTD simulation and a classical mechanical model. The model calculations show the PIT caused in the orthogonally arranged cavities possesses a clear resonance order dependence. These results demonstrate the order selective spectrum modulation effect in nanoscale resonator structures.

P16: Tunability of Epsilon-Near-Zero behavior in a Self-assembled Liquid Crystal - Nanoparticle Hybrid Metamaterial

Amit Bhardwaj

Centre for nano and soft matter sciences (India)

We report experimental evidence of tunable epsilon-near zero (ENZ) behavior in a self-assembled lamella superstructure consisting of Au nanoparticles capped with a photo-active chiral liquid crystal ligand. Upon irradiation with UV, the LSPR peak of Au red-shifts by ~ 10 nm which restores with white light. The effective permittivity indicates ENZ behavior in the visible spectrum with a band width of ~ 45 nm which gets enhanced by a factor of 1.6 on UV illumination. Theoretical calculations based on effective medium approach, support the experimental findings.

P17: Wideband and High-gain Metasurface-Based Circularly Polarized Antenna

Kam Eucharist Kedze, Ikmo Park

Ajou University (Korea)

A metasurface-based circularly polarized microstrip patch antenna with wide bandwidth and high gain is presented. The antenna structure is a square modified microstrip patch that is sandwiched between a metasurface and the ground plane. The microstrip patch and cross slot, in conjunction with the metasurface, generate multiple resonances and AR minimum points that are combined to produce large impedance, AR, and 3-dB gain bandwidths.

P18: Engineering Plasmon-Enhanced Upconversion via Transformation Optics

Avi Mathur, Abhishek Mukherjee, Jasvith Raj Basani, Kannan Ramaswamy

Birla Institute Of Technology and Science (BITS) (India)

Plasmon Enhanced photon upconversion via nanoparticles has shown promise experimentally with a vast multitude of geometries. Applying transformation optical techniques to transform the geometry into a structure that is more feasible to fabricate, is an idea that is attractive in its prospects. These techniques are based on the invariance of the basic laws governing the light-matter interactions and also scale up the output in comparison with the original geometry, replacing the initially used materials with the transformed ones.

P19: Longitudinal and Transverse Surface Plasmon-Enhanced Upconversion in Nanopillar shaped geometries

Abhishek Mukherjee, Avi Mathur, Jasvith Raj Basani, Kannan Ramaswamy

Birla Institute Of Technology and Science (BITS) (India)

Upconversion Fluorescence, which refers to a process of converting low energy photons to higher energies, has received an explosion of interest owing to its potential use in applications such as temperature sensing, biological imaging, etc. A feasible strategy to improve upconversion efficiencies has been manipulating light-matter interaction at the position of an upconverting emitter. Thus, surface plasmon resonance has come into the spotlight. This work aims to demonstrate a nanopillar geometry which can exhibit longitudinal and transverse plasmon resonances.

P20: Plasmonic color filters: angular dependence

Katarína Rovenská¹, Filip Ligmajer¹, Jan Chocho², Jiří Liška¹, Tomáš Šikola¹

¹Brno University of Technology (Czech Repub), ²ON Semiconductor (Czech Repub)

Plasmonic color filters promise a potential solution to miniaturization of color-filtering elements in modern devices. Yet, the dependence of plasmonic filters' spectral responses on the incident angle of light rays has been addressed minimally. In this work, we analyze the angular dependence in plasmonic color filters consisting of nanoholes in metallic film. For these purposes, various plasmonic color filtering arrays (differing in structure size, shape, periodicity, etc.) were characterized with transmission optical spectroscopy for nonzero incident light angles.

P21: A Software for Simulation and Computer Aided Inverse Design of Nanophotonic Metamaterials Based on Periodically Patterned Metasurfaces

Roy Avrahamy¹, Moshe Zohar², Benjamin Milgrom³, Mark Auslender¹

¹Ben Gurion University of the Negev (Israel), ²Shamoon College of Engineering (Israel), ³Jerusalem College of Tech (Israel)

A software for simulation and inverse design of micro- and nano-optical devices embedding smooth and periodically patterned layers is described. With this tool, a ~ 100 % efficient, silicon (Si) based, photodetector (PD) for a near-infrared (NIR) wavelength is designed, and on this example the topological photonics features are shown.

11:40 - 12:40 — Victor Veselago Room

Session 4A14

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

11:40 : Invited talk

Transverse Spin States at Momentum Topological Interfaces utilizing Optical Anisotropy

Xianji Piao, Sunkyu Yu, Namkyoo Park

Seoul National University (Korea)

Here we investigate the topological interface between different momentum geometry, as an optical analogy of the quantum spin Hall effect. By exploiting the alteration of optical anisotropy, the spatial transition of the differential geometry of momentum states are demonstrated. We show that this topological interface leads to the excitation of a transverse-spin state with robust spin-dependent unidirectional transport.

12:00 : Invited talk

A New Type Terahertz Quantum Cascade Laser Using Graphene-based van der Waals Heterostructures

Taiichi Otsuji

Tohoku University (Japan)

This paper reviews recent advances in the research for graphene-based van der Waals heterostructures towards a new type of terahertz quantum-cascade lasers.

12:20 : Invited talk

Ultrafast THz Absorption Modulation in a Graphene-Based Thin Absorber

Anastasios Koulouklidis¹, Anna Tasolamprou¹, Christina Daskalaki¹, Eydokia Kyriakou¹, M. Said Ergoktas², George Kenanakis¹, George Deligeorgis¹, Maria Kafesaki¹, Coskun Kocabas², Eleftherios Economou¹, Costas Soukoulis³, Stelios Tzortzakis⁴

¹IESL-FORTH (Greece), ²University of Manchester (United Kingdom), ³Iowa State University (USA), ⁴Texas AM Universtity at Qatar (Qatar)

We study experimentally and theoretically an ultrafast, optically tunable graphene-based thin film absorption modulator for operation in the THz regime. Graphene hot carriers generated by an ultrafast near-IR pulse, induce negative photoconductivity with a decay time of 2.79ps. As a result, a modulation of 40 % in the THz absorption at 2.17THz is observed in the structure. Additionally, we show that the modulation can be further increased at 60 % by simultaneously varying the conductance of graphene via ionic gating.

11:40 - 12:40 — Allan Boardman Room

Session 4A15

Metasurfaces, flat optics, FSS and HIS

11:40 : High Efficiency Titanium Dioxide Huygen's Metasurfaces In UV

Yu Cheng Chou, Ta-Jen Yen

National Tsing Hua University (Taiwan)

Metasurfaces possess the capability to manipulate multi aspects of light and have been used to demonstrate wavefront engineering devices. In this project, we choose Huygens' metasurfaces with high efficiency and small thickness to wavelength ratio to achieve wavefront shaping. Overlap of electric and magnetic resonances in titanium dioxide nanodisk employs Huygens' condition which gives our device a full phase coverage and close unity transmission in UV. Such properties enable an efficient light focusing flat optical device.

11:55 : Beam-type Elastic Metagratings for Selective Reflections of Longitudinal Waves

Shin Young Kim¹, Woorim Lee¹, Joong Seok Lee², Yoon Young Kim¹

¹Seoul National University (Korea), ²Chungnam National University (Korea)

Metagratings have recently received much attention in electromagnetic and acoustic wave research fields. Governed by the diffraction grating theory, metagratings enable precise wavefront steering by suppressing undesired high-order scattering modes. Recently, we proposed a novel elastic metagrating model using a periodic arrangement of beam-type members in order to realize efficient steering of longitudinal waves [1]. Based on analytical modeling of the beam-type elastic metagratings, anomalous reflections and asymmetric splitting of longitudinal waves were realized successfully in numerical and experimental studies.

12:10 : Optically tunable nanoantennas for visible range based on vanadium dioxide

Filip Ligmajer, Peter Kepič, Martin Hrtoň, Tomáš Šikola

Brno University of Technology (Czech Repub)

Implementing dynamic tunability into the design of metasurfaces is one of the major challenges of the field today. In our contribution we will review various approaches to incorporation of vanadium dioxide into metasurfaces. In particular, we will explore utilization of VO₂ nanostructures as both Mie resonant and also propagation-phase building blocks of tunable metasurfaces.

12:25 : All-dielectric metasurface doublet enabling beam steering and polarizing beam splitting

Changyi Zhou¹, Woo-Bin Lee¹, Chul-Soon Park¹, Song Gao¹, Duk-Yong Choi², Sang-Shin Lee¹

¹Kwangwoon University (Korea), ²Australian National University (Australia)

Multifunctional metasurfaces, fulfilling a variety of tasks, have attracted drastically growing interest. Here we proposed and embodied an all-dielectric metasurface doublet (MD) by vertically concatenating twosome arrays of rectangular nanoresonators on either side of a quartz substrate, in which distinct phase profiles are encoded for orthogonally polarized light. Bifunctional beam manipulation including enlarged steering and polarizing beam splitting was enabled by the MD. The superior performance of the proposed device paves the way to large-scale photonic integrated applications.

11:40 - 12:40 — Tatsuo Itho Room**Session 4A16****Photonic Crystals****11:40 : Self-assembled Photonic Crystals for Colorful Radiative Coolers****Hyeon Ho Kim, Eunji Im, Seungwoo Lee***Korea University (Korea)*

Recently, radiative cooling has been actively studied, as objects can be cooled without using additional energy. Here, we demonstrate a new class of radiative cooler, this is silica photonic crystals, which are used for the structural colorization. The intrinsic phonon vibration of silica leads to the absorption of mid-infrared (mid-IR) waves, which in turn cool the objective in a radiative way. At daytime, these assembled photonic crystals effectively reduce the temperature of the crystalline-silicon (c-Si), while maintaining appearing colors.

11:55 : Enhancement of Laser-Induced Damage Threshold by 3D Photonic Crystal Structure**Lei Pan, Hongbo Xu, Ruizhen Lv, Yao Li***Harbin Institute of Technology (China)*

Enhancement of laser-induced damage threshold by 3D photonic crystal (3D-PC) structure is demonstrated using both experimental and simulation methods. A polystyrene 3D-PC with a reflection peak at 1064nm is fabricated. The laser-induced damage threshold (LIDT) of 3D-PCs is as 2.4 times high as the LIDT of disordered polystyrene films. The simulation results show that the electric field is contained in the pores of 3D-PCs while is reduced in the solid part, which may be responsible for the high LIDT.

12:10 : Laser emission from atoms placed in vacuum**Iliya Doronin¹, Alexander Zyablovsky¹, Evgeny Andrianov¹, Alexander Pukhov¹, Yurii Lozovik¹, Alexey Vinogradov¹, Alexander Lisiansky²**¹*Dukhov Research Institute of Automatics (Russia)*, ²*Queens College of the City University of New York (USA)*

We consider pumped active atoms placed in a free space. We show that such atoms can emit coherent laser-like radiation at sufficient pump rate. This is possible due to the fact that active atoms interact with electromagnetic field of vacuum with a finite strength, even though there are no distinct modes of electromagnetic field. Our results suggest the possibility of creating new types of low-cost laser devices, as well as open new perspective on fundamental physics behind lasers.

12:25 : Light trapping with subwavelength compound parabolic concentrators**Ashsih Prajapati, Gilad Marko, Gil Shalev***Ben Gurion University of the Negev (Israel)*

Light trapping and the broadband absorption of the solar radiation is of interest to solar energy applications. In the current work, we report a new paradigm for light trapping, that is light trapping based on arrays of subwavelength nonimaging light concentrators (NLCs). We numerically show that NLC arrays provide ~50 % broadband absorption enhancement of the solar radiation compared with that of optimized nanopillar arrays. We show that CPC arrays (in contrast to nanopillar arrays) function as anti-transmission layers.

11:40 - 12:40 — Christian Huygens Room**Session 4A17****New materials for photonics****11:40 : Guided mode resonant gratings with van der Waals materials: A platform for high-Q resonances and its applications****Qilin Hong, Xingqiao Chen, Xiaodong Yuan, Zhihong Zhu, Jianfa Zhang***National University of Defense Technology (China)*

A type of remarkably high-quality resonant nanostructure based on atomically thin two-dimensional materials

is proposed. It is shown theoretically and numerically that the excitation of leaky modes in the proposed structures can achieve resonances with extremely narrow linewidths down to 0.0005 nm and high Q-factors up to millions in the telecom range. This new type of 2D material resonant nanostructure can be employed for a variety of applications ranging from lasers, filters and polarizers to nonlinear optical devices.

11:55 : Graphene-metal hybrid for strongly enhanced terahertz harmonics

Jan-Christoph Deinert¹, David Alcaraz Iranzo², Hassan A. Hafez³, Frank Koppens⁴, Mischa Bonn³, Dmitry Turchinovich⁵, Michael Gensch⁶, Sergey Kovalev¹, Klaas-Jan Tielrooij⁷, Klaas-Jan Tielrooij⁷

¹Helmholtz-Zentrum Dresden-Rossendorf (Germany), ²ICFO - Institut de Ciències Fòniques (Spain), ³Max-Planck-Institut für Polymerforschung (Germany), ⁴The Barcelona Institute of Science and Technology (Spain), ⁵Universität Bielefeld (Germany), ⁶Technische Universität Berlin (Germany), ⁷Catalan Institute of Nanoscience and Nanotechnology (ICN2) (Spain)

By combining the peculiar ultrafast heating-cooling dynamics of graphene electrons with the enhancement of incident electromagnetic fields by a metallic grating, we have created a hybrid structure with an unprecedentedly large third order (sheet) susceptibility in the terahertz range. In particular, we observe terahertz (THz) third harmonic generation with a field conversion efficiency above 1%. The nonlinear enhancement furthermore allows for the observation of signatures of higher harmonics (up to 9th order) using a table-top laser setup.

12:10 : Local Variations of Light Absorption and Emission in Monolayer WS₂ Flakes

Marzia Ferrera¹, Michele Magnozzi¹, Theo Pflug², Simona Pace³, Lorenzo Ramò¹, Markus Olbrich², Paolo Canepa¹, Hasret Agircan⁴, Alexander Horn², Stiven Forti⁵, Ornella Cavalleri¹, Camilla Coletti³, Francesco Bisio⁶, Maurizio Canepa¹

¹Università di Genova (Italy), ²Laserinstitut Hochschule Mittweida (Germany), ³Istituto Italiano di Tecnologia (Italy), ⁴Istanbul Technical University (Turkey), ⁵Center for Nanotechnology Innovation IIT@NEST (Italy), ⁶CNR-SPIN (Italy)

The investigation of the local excitonic response of two-dimensional transition metal dichalcogenides is crucial both for fundamental research and in view of their implementation in novel optoelectronic devices. In this work, we compare the photoluminescence spatial patterns of monolayer WS₂ flakes grown by chemical vapor deposition with their position-resolved dielectric function probed by imaging spectral ellipsometry. The two datasets show both correlated and uncorrelated spatial patterns. Micro-structural variations within the flakes are deemed responsible for the observed discrepancies.

12:25 : The role of trion excitation in the shear-induced optical bandgap tunability of MoS₂

Assaf Ya'akovovitz, Avi Niv

Ben-Gurion University of the Negev (Israel)

With high strength, valley polarization, and direct bandgap, Mono-layered MoS₂ is becoming an essential member of a growing family of materials that will shape our optoelectronic future. In this talk, we will describe bandgap tuning under shear strain. Our study reveals an intricate connection between the global electronic structure and the localized excitons and trions of this material.

11:40 - 12:40 — Augustin Fresnel Room

Session 4A18

Plasmonics and nanophotonics

11:40 : Photothermal characteristics in silicon nanoparticles

Yongqian Li¹, Xingyu Yang¹, Yidyu Yu¹, Binbin Wang², Xiaoying Li¹, Rafael Salas-Montiel²

¹Key Laboratory of Micro/Nano Systems for Aerospace of Ministry of Education (China), ²Université de technologie de Troyes (France)

Photothermal characteristics of nonspherical dielectric nanoparticles was reported. The significant nanoheating effect of silicon cuboid nanoparticles is attributed to the resonant modes of electric and magnetic resonances of different orders. Especially, the remarkable photothermal effect arises from the hybrid modes, the overlapping of different resonant modes, due to the varying ratio of cuboid nanoparticles. The thermal radiation of these nanoparticles due to interactions and interference between both electric and optically-induced

magnetic resonances of different orders are investigated.

11:55 : Nanoscaled VO₂ insulator-to-metal transition controlled by plasmonic single-nanoantenna

Luca Bergamini¹, Bigeng Chen², Daniel Traviss², Yudong Wang², C. H. de Groot², Jeffrey M. Gaskell³, David W. Sheel³, Nerea Zabala¹, Javier Aizpurua⁴, Otto L. Muskens²

¹UPV-EHU (Spain), ²University of Southampton (United Kingdom), ³University of Salford (United Kingdom), ⁴CSIC-UPV/EHU and DIPC (Spain)

Resonant plasmonic nanoantennas are known to concentrate light at the nanoscale around its surface. VO₂ is of interest for its insulator-to-metal transition when heated up above the relatively low critical temperature of 68°C. In this study we show how the plasmonic feature of a single antenna can be used to induce and control the insulator-to-metal transition in a VO₂ film.

12:10 : Non-classical light source with single photon and squeezing properties in a nanoscale photonic-crystal-plasmonic system

Lingxiao Shan, Juanjuan Ren, Qi Zhang, Yun Ma, Xueke Duan, Qihuang Gong, Ying Gu

Peking University (China)

Single photon and squeezing properties are important in on-chip quantum applications. Single photon sources were investigated in microstructures like nanowires. However, on-chip squeezing light sources remain exploring. Here, a photonic-crystal-plasmonic nanostructure is proposed with strong mode-emitter coupling from a band-edge mode. Our calculation shows non-classical light is achieved with degree of squeezing 0.46 dB and $g^2(0) < 0.1$ under strong coupling. Besides 70% emission are channeled in the 2D line defect. The system provides possibilities to on-chip versatile non-classical light sources.

12:25 : Hot electrons remote excitation and their ultrafast harvesting

Romain Hernandez¹, Renato Juliano-Martins¹, Mario Lodari², Michele Celebrano², Marco Finazzi², Lamberto Duo², Giovanni Isella², Marlène Petit¹, Adrian Agreda¹, Jean-Claude Weeber¹, Alexandre Bouhelier¹, Monica Bollani², Olivier Demichel¹, Paolo Biagioni², Benoit Cluzel¹

¹Laboratoire Interdisciplinaire Carnot de Bourgogne (France), ²Politecnico di Milano (Italy)

Generation, collection and relaxation of hot electrons within plasmonic devices are studied with two approaches. The first consists in investigating with a hyperspectral nearfield imaging technique the delocalized hot electrons produced along a Au waveguide by a propagative surface plasmon. The second, in producing hot electrons with localized surface plasmons within a Au grating on top of a silicon-on-insulator substrate. This Schottky barrier hot electron photodetector allows studying the absorption and transport dynamics of hot electrons.

11:40 - 12:40 — Ibn Al-Haytham Room

Session 4A19

Metamaterials and metasurfaces

11:40 : Multi-band middle-infrared-compatible camouflage with thermal management via simple photonic structures

Ziquan Xu¹, Meiyang Pan¹, Huanzheng Zhu¹, Min Qiu², Qiang Li¹

¹Zhejiang University (China), ²Westlake University (China)

We demonstrate two simple photonic structures for visual and MIR camouflage with thermal management utilizing thin-film interference. The additional laser camouflage is realized by further incorporating simple wavelength-scale grating structures. The fabricated structures exhibit dual-band MIR camouflage by low emission, radar camouflage to carbon dioxide laser by high absorption at 10.6 micron, visual camouflage by structural colors and thermal management by high emission in 5-8 micron band.

11:55 : Tunable Fano Resonance in a Liquid Crystal Colloidal Metamaterial

Amit Bhardwaj

Centre for nano and soft matter sciences (India)

A colloidal metamaterial is realized by dispersing submicron-sized high-refractive-index dielectric resonators

in a nematic liquid crystal medium. Darkfield hyperspectral imaging reveals that when the NLC molecules reorient on application of an ac electric field a doughnut-shaped scattering pattern is obtained, indicating the occurrence of Fano resonance. The theoretical simulation based on the "Multi-pole Fano interference model" confirms the experimental findings. With increasing voltage, the value of Fano parameter q decreases and approaches unity corresponding to an ideal Fano shape.

12:10 : **Meta-lenses Using Metal-Dielectric-Metal Plasmonic Waveguide at Optical Frequency**

Boopalan Ganapathy, Jayanth N. V., Subramaniam Chittur K.

Vellore Institute of Technology (India)

Meta-lenses are proved to be effective in different regions of the electromagnetic spectrum. The study presented here is on meta-lenses working at an optical frequency. Plano-concave meta-lenses are designed at red wavelength (660 nm) of the visible region. Metal-dielectric-metal plasmonic parallel plate waveguide is used as the unit cell to emulate an Epsilon-Near-Zero medium. Zoning technique is also presented that has an advantage of less volume compared to the non-zoned one.

12:25 : **Switching the Optical Chirality by Magnetic fields in Magnetoplasmonic Metasurfaces**

Jun Qin¹, Longjiang Deng¹, Tongtong Kang¹, Gaspar Armeltes², Lei Bi¹

¹*University of Electronic Science and Technology of China (China)*, ²*Instituto de Micro y Nanotecnologia (INM-CNM-CSIC) (Spain)*

We report a magneto-optical metasurface device using low loss Ce:YIG thin films for active chiroptical photonic device applications. A far field modulation of the circular dichroism from $-0.6^\circ \pm 0.2^\circ$ to $+1.9^\circ \pm 0.1^\circ$ at 950 nm wavelength is observed under applied magnetic fields, enabling efficient control of optical chirality both in the far field and near field at the subwavelength scale.

11:40 - 12:35 — **Gaston Floquet Room**

Session 4A20

Metasurface for information processing

Organized by: Jensen Li and Xianzhong Chen

Chaired by: Jensen Li and Xianzhong Chen

11:40 : **Invited talk**

Terahertz Active Metasurface

Yan Zhang, Xinke Wang, Teng Wang, Jinying Guo

Capital Normal University (China)

Metasurface provides a number of approaches to manipulate the wavefront of light. Active control of the metasurface will bring more fascinate applications. We introduce two terahertz active metasurface devices which are fabricated with vanadium dioxide. With thermal exciting, the functions of devices can be switched on and off. The performances of the devices are characterized with a focal plane terahertz imaging system. It was found that the fabricated devices can achieve the preset functions well.

12:00 : **Invited talk**

Optical Metasurface for Engineered Polarization Profile

Xianzhong Chen, Yuttana Intaravanne

Heriot-Watt University (United Kingdom)

The unprecedented capability of optical metasurfaces in the manipulation of the light's polarization at sub-wavelength resolution has provided an unusual approach for arbitrary manipulation of polarization profiles. A compact metasurface platform has been demonstrated to arbitrarily engineer a polarization profile that is very difficult or impossible to realize with conventional optical elements. We have developed various metasurfaces to engineer the polarization profile of a light beam.

12:20 : **3D-printed complex-amplitude metasurface for orbital angular momentum holography**

Haoran Ren¹, Xinyuan Fang², Jaehyuck Jang³, Johannes Burger¹, Junsuk Rho³, Stefan A. Maier¹

¹Ludwig-Maximilians-Universität München (Germany), ²University of Shanghai for Science and Technology, (China), ³Pohang University of Science and Technology (POSTECH) (Korea)

Metasurface holograms, consisting of subwavelength structures on a flat surface for wavefront shaping, promise new applications in information technologies. To increase the bandwidth of a metasurface hologram, different degrees of freedom of light and in particular, orbital angular momentum with an infinite quantum number hold great promise. Here, we demonstrate the design and 3D laser printing of a large-scale complex-amplitude metasurface hologram for high-dimensional orbital angular momentum-multiplexing holography.

11:40 - 12:40 — Lawrence Bragg Room

Session 4A21

Light-matter interactions in new materials and meta-architectures

Organized by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

Chaired by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

11:40 : **Invited talk**

A Deep Learning Approach to the Forward Prediction and Inverse Design of Plasmonic Metasurface Structural color

Mehdi Keshavarz Hedayati, Nathan Roberts

Durham University (United Kingdom)

Here, optimized Deep Neural Network models are presented to enable the forward and inverse mapping between metamaterial and corresponding color. The forward model can predict color with *textgreater*96% accuracy, with a 105 order of magnitude decreases in computational time when compared to simulations. For the first time, the use of synthetic training data for self-learning is reported which results in a 15% improvement in training accuracy of the inverse model. The findings enable the discovery of new photonic materials.

12:00 : **Invited talk**

Space-time metamaterials: homogenization theory, giant bianisotropy and light drag without moving media

Paloma A. Huidobro¹, Mario Silveirinha¹, Emanuele Galiffi², John Pendry²

¹University of Lisbon (Portugal), ²Imperial College London (United Kingdom)

Here I will present a theory of homogenisation of space-time metamaterials, which yields expressions for the effective permittivity, permeability and magnetoelectric coupling in the long wavelength limit. The derived parameters show that synthetic motion can result in giant bianisotropy and the dragging of electromagnetic fields down to the quasistatic limit and without any moving matter

12:20 : **Invited talk**

Watching operating Li:ion batteries by Raman through hollow-core optical fibres

Ermanno Miele, Wesley M. Dose, Ilya Manyakin, Michael H. Frosz, Philip St.J. Russell, Clare P. Grey, Jeremy J. Baumberg, Tijmen G. Euser

University of Cambridge (United Kingdom)

We demonstrate a fibre-optic sensing method capable of monitoring chemical changes within Li:ion cells under real working conditions. Our technique is based on optofluidic single-ring hollow-core fibres, that uniquely allow light to be guided at the centre of a microfluidic channel. We integrate the fibres into working Li:ion cells, use them to take sub-microlitre samples of the electrolyte liquid, and analyse these by background-free Raman spectroscopy to identify early signs of battery degradation.

11:40 - 12:40 — Rene Descartes Room

Session 4A22

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

11:40 : **Invited talk****Fabrication of ultrathin multilayer structures and their characterization****Andrei V. Lavrinenko, Johneph Sukham, Maryam Mahmoodi, Osamu Takayama, Seyed Hassan Tavasoli, Radu Malureanu***Technical University of Denmark (Denmark)*

Multilayer structures have been in the focus of research for decades. They exhibit various optical properties starting from anti-reflection coating, Bragg gratings and finishing with hyperbolic metamaterials. Meantime they accept quite simple theoretical analysis and straightforward modeling. In spite of such intensive study there are still numerous questions about quality of fabrication processes, accessible parameters ranges and adequate models accurately explaining their performance. We report on our activity in fabrication of ultrathin dielectric and metal-dielectric layers, and their characterization.

12:00 : **Invited talk****Scattering properties of Parity-Time symmetric chiral metamaterials****Maria Kafesaki¹, Ioannis Katsantonis², Sotiris Droulias², Costas Soukoulis³, Eleftherios Economou²**¹FORTH-IESL and University of Crete (Greece), ²FORTH-IESL and Univ. of Crete (Greece), ³FORTH-IESL and Iowa State University (USA)

Combining parity-time (PT) symmetry and chirality one can achieve a variety of novel electromagnetic properties and effects. Here we demonstrate some of those effects in a simple bi-layer PT-symmetric chiral structure illuminated by a plane wave. Phenomena such as asymmetric transmitted wave ellipticity, mixed PT-related phases, simultaneous coherent perfect absorption and lasing of circularly polarized waves, etc., are numerically demonstrating. All those phenomena, realizable with realistic chiral metamaterials, empower PT-symmetric chiral systems with unique possibilities regarding electromagnetic wave control.

12:20 : **Invited talk****Polarisation study of the light scattered by nanoporous plasmonic microparticles****Pritam Khan, Grace Brennan, Daragh Rice, Syed A. M. Tofail, Ning Liu, Christophe Silien***University of Limerick (Ireland)*

Nanoporous microparticles exhibit large surface-to-volume ratio and can detect chemicals and biomolecules. Using dark-field microscopy, we observed that, when the nanoporous microparticles are illuminated by circularly polarized lasers, the scattered light polarization is inverted. The inversion is interesting as the microparticles are (nearly-)spherical, and the nanopores do not exhibit symmetry. We observed that the scattered light polarization is tunable when plasmonic nanoparticles and short molecules are in the pores. We discuss these findings and their exploitation as sensors.

Lunch

12:40 - 14:00

14:00 - 16:00 — **Victor Veselago Room**

Session 4A23

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

14:00 : **Invited talk****The Revolutionary Advent of Magnetless Nonreciprocal Metasurfaces**

Christophe Caloz¹, Toshiro Kodera²

¹*Katholieke Universiteit Leuven (KU Leuven) (Belgium)*, ²*Meisei University (Japan)*

We present a narrative overview of our work on magnetless nonreciprocal metasurfaces over the past decade, showing how the spacetime telescoping of the two emerging areas of magnetless nonreciprocity and metasurface transformation has led to the revolutionary technology of magnetless nonreciprocal metasurfaces.

14:20 : Invited talk

Photonic entanglement based quantum metrology

Laurent Labonté¹, Mattis Reisner¹, Florent Mazeas¹, Djeylan Aktas¹, Florian Kaiser¹, Romain Dauliat², Philippe Roy², Raphael Jamier², Sebastien Tanzilli¹

¹*Université Côte d'Azur (France)*, ²*Université de Limoges (France)*

During the conference, we show how we take advantage of these capabilities to gather optical fiber photonic engineering with quantum optics. More specifically, we aim at presenting two quantum-based methods for (i) high-accuracy (10⁻⁵) and dispersion-free measurement of refractive index difference and (ii) chromatic dispersion measurement based on the concept of quantum white-light interferometry that allows absolute measurement of chromatic dispersion with 2.5 times improved accuracies compared to state-of-the-art realizations at telecom wavelengths.

14:40 : Invited talk

Nanomechanics with plasmonic nanoantennas

Hilario D. Boggiano¹, Rodrigo Berte², Alberto F. Scarpettini³, Emiliano Cortés², Stefan A. Maier², Andrea V. Bragas¹

¹*Universidad de Buenos Aires (Argentina)*, ²*Ludwig-Maximilians-Universität München (Germany)*, ³*Universidad Tecnológica Nacional (Argentina)*

The measurement of the mechanical properties of thin films is a key factor both for their integration into large-scale devices and to determine their duration as advanced coatings, sensors or optoelectronic components among other applications in the current technological landscape. Here we introduce a new method using optical antennas as nanomechanical sensors to determine the nanoscale mechanical response of polymeric films in the GHz range, although the method can be extended to a wide range of materials

15:00 : Invited talk

Transition to strong coupling regime for quantum emitters coupled to a plasmonic resonator

Tigran V Shahbazyan

Jackson State University (USA)

We present a model for exciton-plasmon coupling based on energy exchange between emitters and localized surface plasmons in metal-dielectric structures. Plasmonic correlations between emitters give rise to a collective state exchanging its energy with a resonant plasmon mode. By accurately defining the plasmon mode volume, we relate the QE-plasmon coupling to collective energy transfer rate. For QEs distributed in extended region enclosing a plasmonic structure, the ensemble QE-plasmon coupling saturates to a universal value independent of system size and shape.

15:20 : Invited talk

Magneto-optical control with non-reciprocal surface waves for nanoscale thermotronics

Svend-Age Biehs

Carl von Ossietzky Universität (Germany)

Magneto-optical materials have been proposed as promising candidates for an active control of nanoscale heat radiation. Unexpected effects like the thermal radiative Hall effect, persistent currents, giant magneto-resistance, and circular heat fluxes, persistent angular momentum and spin of thermal radiation have been highlighted. We review recent developments in this new direction of magneto-optical thermotronics focusing on the rectification of nanoscale heat fluxes by means of non-reciprocal surface wave and reveal the spin coupling mechanism behind this effect.

15:40 : Invited talk

Long-range FRET-mediated exciton diffusion in highly efficient self-assembled monolayers of cesium lead halide perovskite nanocrystals

Monica Lorenzon, Matthew Jurow, Min Ji Hong, Yi-Hsien Lu, Edward S. Barnard, Miquel Salmeron, Yi

Liu, Alexander Weber-Bargioni, Erika Penzo, Adam Schwartzberg

Lawrence Berkeley National Laboratory (USA)

Colloidal perovskite nanocrystals (PNCs) are solution-processable functional materials whose emission can be easily tuned via both size and composition. In this work, we investigate exciton diffusion in a self-assembled, close-packed monolayer of inorganic PNCs and measure a record diffusion length of 480nm, one order of magnitude large than what previously observed for traditional cadmium-based quantum dots. Moreover, we report a detailed investigation of the effect of a ceramic protective coating on our PNC monolayers, deposited via atomic layer deposition.

14:00 - 15:15 — Allan Boardman Room

Session 4A24

Chiral and hyperbolic metamaterials

14:00 : Active Chiral Metasurfaces via Colloidal Self-Assembly

Andreas Fery¹, Patrick T. Probst², Martin Mayer², Vaibhav Gupta², Anja Maria Steiner², Gunter K. Auernhammer², Tobias A. F. Koenig²

¹*Technische Universität Dresden (Germany)*, ²*Leibniz-Institut für Polymerforschung Dresden e.V. (Germany)*

Active chiral metasurfaces enable continuous on-chip polarization engineering, detection and encoding. Here, we introduce a facile bottom-up approach that produces circular dichroism of up to 11 degrees in the visible-near-infrared spectral region. This pronounced effect surpasses previous colloidal approaches by two orders of magnitude. Furthermore, the presented design of stacked particle chain arrays allows in-situ re-stacking and local compression to tune dynamically all aspects of circular dichroism: sign, magnitude and spectral position.

14:15 : Towards dynamic and chiral plasmonics using liquid crystals

Wiktor Lewandowski

University of Warsaw (Poland)

Nanostructures with chiral symmetry and plasmonic properties attract considerable interest. Here, we present endeavors towards achieving thin films of chiral, reconfigurable nanostructures, which resolve some of the current synthetic challenges. Here, synthesis relies on mixing (chiral) liquid crystals with gold nanoparticles. Due to the soft character of liquid crystals, these nanoarchitectures exhibit helical, hierarchical and switchable structure. Durability, configurability and tailorability of such nanomaterials in the solid state, is particularly attractive for real world applications.

14:30 : Tailored Hyperbolic Metamaterials for Enhanced Superlensing Effects along the entire Visible Regime

Nadia Pinton¹, Vincenzo Caligiuri², Giulia Biffi¹, Antonio De Luca², Roman Krahn¹

¹*Istituto Italiano di Tecnologia (Italy)*, ²*University of Calabria (Italy)*

Superlensing effects have been established in presence of an Epsilon-Near-Zero (ENZ) resonance. However, ENZ engineered metamaterials in the visible spectral range have been elusive. In this paper, we demonstrate that, by using Hyperbolic Metamaterials (HMMs) composed of a periodic stack of metal/high-refractive index dielectric bilayers and varying the oxidation degree of the dielectric material, the ENZ resonance and thus the superlensing effects can be shifted in wavelength along the entire visible range.

14:45 : Giant third-harmonic dichroism in all-dielectric chiral metasurfaces based on quasi-bound states in the continuum

Marco Gandolfi, Andrea Tognazzi, Davide Rocco, Costantino De Angelis, Luca Carletti

CNR-INO (National Institute of Optics) (Italy)

We develop a new approach based on quasi-BICs to develop chiral metasurfaces exhibiting nonlinear circular dichroism (up to 99.9%) and high conversion efficiency. Tuning mode interference allows selective linear and nonlinear circular dichroism.

15:00 : Asymmetry leading to Circular Conversion Dichroism in Optical Regime**Ahsan S. Rana¹, Inki Kim², Muhammad Zubair¹, Junsuk Rho², Muhammad Q. Mehmood¹**¹University of the Punjab (Pakistan), ²Pohang University of Science and Technology (POSTECH) (Korea)

Asymmetric transmission (AT) of light is an intriguing phenomenon of chirality that can lead to fashion optical diodes. However, achieving AT in the optical regime is a monumental task because of dielectric losses. In this research, we present a novel approach that utilizes nano bars of hydrogenated amorphous silicon to obtain AT in the optical regime with the maximum at 633 nm. The design also uses dielectric loss to engineer circular conversion dichroism (CCD), which can completely eradicate backscattering.

15:15 - 16:00 — Allan Boardman Room**Session 4A25****Super-resolution imaging****15:15 : Exploring object sub-wavelength features using transformation-optics based imaging systems****Mircea Giloan¹, Robert Gutt², Gavril Saplacan³**¹Babes-Bolyai University (Romania), ²National Institute for Research and Development of Isotopic and Molecular Technologies (Romania), ³Company for Applied Informatics (Romania)

The information about the fine sub-wavelength details of an object are carried by waves with high transverse wave vectors which become evanescent and decay exponentially inside a classical material. The proposed optical imaging systems made of transformation-optics inspired lenses are capable to process the waves with high transverse wave vectors and enable their contribution to the reconstructed image. The capability of these optical imaging systems to perform magnified images of sub-wavelength features of arbitrary objects is theoretically analyzed.

15:30 : Plano-Convex-Microsphere (PCM) super lens for direct laser nano-fabrication and optical super-resolution imaging in far-field**Bing Yan¹, Liyang Yue¹, Rakech Dhama¹, Daniel Siebadji Tchuimeni¹, Xibin Yang², Daxi Xiong², Zengbo Wang¹**¹Bangor University (United Kingdom), ²Chinese Academy of Sciences (China)

This paper proposes and demonstrates a high-performance all-dielectric compound superlens, formed by integrating a conventional Plano-Convex lens with a high-index Microsphere lens. We call such lens the Plano-Convex-Microsphere (PCM) lens, which is developed for far-field super-resolution applications. The compound PCM lens is highly versatile and can be simply adapted into an existing optical system to realize super-resolution function. Here, the super-resolution features of the PCM lens were demonstrated for two applications: direct laser nanofabrication and white light nano-imaging.

15:45 : Investigation of proximity effects in light funnel arrays using near-field optical microscopy**Ankit Chauhan, Gil Shalev***Ben Gurion university of the Negev Beer Sheva (Israel)*

Surface arrays of silicon light funnels (LF) were suggested as a promising platform to produce broadband absorption that is considerably superior to that of the well-known nanopillar (NP) arrays, for example. The current study explores the underlying mechanism of broadband absorption in LF arrays. To this end the optical near-field of LF and NP arrays is experimentally probed. We show that in LF arrays the near-field increases as the array period decreases in contrast with NP arrays.

14:00 - 16:00 — Tatsuo Itho Room**Session 4A26****Theory, Fabrication, Experiments, and Applications**

14:00 : Bound states in the continuum in asymmetric waveguides: role of proportionate coupling**Nikolay Shubin, Vladimir Kapaev, Alexander Friman, Alexander Gorbatshevich***Russian Academy of Sciences (Russia)*

We perform theoretical analysis of bound states in the continuum (BIC) formation in a resonator coupled to two waveguides. Analytical description provides clues on BIC conditions - a single point in the parameter space, when exact numerical calculations become cumbersome. The Friedrich-Wintgen mechanism can be realized in asymmetric system with proportionate couplings to waveguides. The derived conditions are universal and can be implemented to electronic or electromagnetic waveguides. As an example we present BIC in an asymmetric quantum billiard.

14:15 : A low-dimensional nonlinear eigenproblem for the complete complex bandstructure and microscopic fields of arbitrary two-domain metamaterials**Antonio Günzler, Cedric Schumacher, Matthias Saba***University of Fribourg (Switzerland)*

Homogenization theories for plasmonic metamaterials usually start with crude approximations that are valid in certain limits in zero order, such as small frequencies, wave vectors and material fill fraction. We here instead present a rigorous solution to Maxwell's equations in binary periodic materials employing a combined Green's-Galerkin procedure to obtain the complex-valued eigenmodes of the material. Our theory can be generalized and remains valid in regimes outside of those accessed by standard approaches.

14:30 : Electromagnetic Modeling of finite Fragments of Metamaterials and Metasurfaces based on Method of Minimal Autonomous Blocks**Yauheni Arlou¹, Sergei Maly¹, Eugene Sinkevich²***¹Belarusian State University (Belarus), ²Belarusian State University of Informatics and Radioelectronics (Belarus)*

Technique to model finite fragments of frequency-selective screens, metasurfaces and flat layers of metamaterials based on method of minimal autonomous blocks is proposed. Use of multichannel macroblocks (scattering matrices) corresponding to unit cells of the material and surrounding space is the basis. The technique significantly decreases requirement to RAM compared to other methods. It can model arbitrary excitation modes of finite structures by local and remote sources. Modeling results of finite fragment of frequency-selective surface are given.

14:45 : Fluorescent Multi-layered films for Label-Free Detection of Volatile Organic Compounds**Heba Megahd¹, Paola Lova¹, Andrea Pucci², Davide Comoretto¹***¹University of Genova (Italy), ²University of Pisa (Italy)*

The detection of vapors is fundamental in many different applications, such as air pollution analysis, industrial process monitoring and breath analysis. This work utilizes a styrene copolymer with fluorescent molecular rotors exhibiting aggregation-induced emission as a promising vapor sensor. Capping thin films of such copolymers with different polymers provides a quick and selective means of detecting volatile organic compounds through fluorescence quenching.

15:00 : Near-field luminescence of two-dimensional semiconductors**Vlastimil Krapek, Petr Dvorak, Lukas Kejik, Martin Konecny, Zoltan Edes, Martin Hrton, Michal Kvapil, Michal Horak, Tomas Sikola***Brno University of Technology (Czech Republic)*

Two-dimensional semiconductors are ideal light sources for on-chip integration. They exhibit strong luminescence, and are capable of single-photon emission. Since the wavelength of the light is considerably larger than the physical dimensions of the emitter, near-field handling of the emission with a deeply subwavelength spatial resolution would be of great importance. Here we present fully near-field photoluminescence study of two-dimensional semiconductors, with a surface plasmon interference device used for the excitation and scanning near-field optical microscopy for the collection.

15:15 : Efficient Implementation of Active Exterior Cloak in Three Dimensions**Cheuk-Him Yeung, Tom Shearer, William J. Parnell***University of Manchester (United Kingdom)*

We describe a cloaking strategy for the 3D Helmholtz equation by using a number of multipolar sources to construct a silent region which shields any object inside from the incoming wave and minimizes the scattering.

Their amplitudes are expressed in a surface integral as we solve the Kirchhoff-Helmholtz integral equation by applying the addition theorem for Bessel functions. The positioning of the sources at the vertices of a Platonic solid simplifies the integral and allows for efficient numerical evaluation.

15:30 : Spark discharge synthesis of noble metals and GeSi nanoparticles for UV-vis-NIR plasmonics

Anna Lizunova, Vladislav Borisov, Dana Malo, Kirill Khabarov, Viktor Ivanov

Moscow Institute of Physics and Technology (Russia)

Noble metals (Pt, Au and Ag) and alloy GeSi aerosol nanoparticles were synthesized by spark discharge in pure argon atmosphere with additional in-flow sintering of nanoparticles at temperatures from 25 to 950 °C. The size, morphology, crystal structure and optical properties of obtained nanomaterials were investigated. It has been established that the thermal treatment of aerosol nanoparticles leads to crucial change in absorption properties and vary the morphology from branched agglomerates to spherical nanoparticles.

15:45 : Synthesis and characterization of palladium nanoparticles by laser ablation in liquids

Mónica Fernández-Arias¹, Ana M. Vilas-Iglesias¹, Mohamed Boutinguiza¹, Daniel Rodríguez², Felipe Arias-González³, Pablo Pou¹, Antonio Riveiro¹, Javier Gil³, Juan Pou¹

¹University of Vigo (Spain), ²UPC-Barcelona TECH (Spain), ³Universitat Internacional de Catalunya (Spain)

The outstanding catalytic and electronic properties of palladium nanoparticles, make them useful in a wide variety of applications including the biomedical field. In this work, palladium nanoparticles are obtained by laser ablation in water and methanol with two different laser sources (a Green nanosecond and a IR picosecond laser) and their physical-chemical properties are analyzed.

14:00 - 15:15 — Christian Huygens Room

Session 4A27

Mesoscale Plasmonics, Nanophotonics and Acoustics

Organized by: Igor V. Minin and Oleg V. Minin

Chaired by: Igor V. Minin and Oleg V. Minin

14:00 : Invited talk

Super-enhancement focusing of Teflon sphere in terahertz band

Liyang Yue¹, Zengbo Wang¹, Bing Yan¹, Oleg Minin², Igor Minin²

¹Bangor University (United Kingdom), ²Tomsk State University (Russia)

A Teflon sphere can focus the light near its shadow surface. In this paper, two circular hotspots having the extremely large field-intensity were discovered around the poles of a specifically sized Teflon sphere irradiated by a plane wave in terahertz band using an analytical algorithm. A huge contribution of scattering amplitude from a single order of mode in electric-field or magnetic-field is considered as the main factor to trigger this phenomenon of super-enhancement focusing.

14:20 : Dielectric Microstructures for Extended Photonic Nanojet Generation

Ksenia Sergeeva, Aleksandr Sergeev

Russian Academy of Sciences (Russia)

The functional aluminum oxide microstructures of various geometry for the generation of extended photonic nanojets in transmission, as well as in reflection modes, were developed. The numerically calculated parameters of photonic nanojets are in good agreement with the experimentally obtained ones. The preliminary studies of the photoluminescence enhancement of sensitive layers under excitation by photonic nanojets revealed up to four-time an increase in its emission intensity.

14:35 : Invited talk

High Intensity Photonic Nanojets from a Gradient Low Refractive Index Profile

Alexander Littlefield, Jinlong Zhu, Lynford Goddard

University of Illinois at Urbana-Champaign (USA)

Designs for photonic nanojet (PNJ) generators are examined. The maximum intensity of the PNJ is used as an objective for optimization under the constraint of maximum refractive index and fixed generator size. Building on the work of Patel et al. with a generator of a step-index profile across tangent microspheres, we propose a gradient index (GRIN) design. Our design has a higher intensity focus yet requires lower index contrast. This eases fabrication while improving resolution in applications including photolithography.

14:55 : Invited talk

Zero-group-velocity mechanical metamaterial

Rafael A. Mendez-Sanchez¹, Bryan Manjarrez¹, Enrique Flores², Arturo Arreola², Gabriela Baez²

¹Universidad Nacional Autonoma de México (Mexico), ²Universidad Autonoma Metropolitana-Azcapotzalco (Mexico)

A quasi-one-dimensional mechanical metamaterial, with a zero-dispersion band, was designed. Each metaatom consists of an aluminum plate weakly connected to neighbouring metaatoms through aluminum beams. The band structure of the perturbative metamaterial was calculated using COMSOL. Depending on the location of the beams a flat band, separated of all other bands, can be engineered. The resulting metamaterial has a group velocity of less than 1 % of the velocity of compressional waves in aluminum.

14:00 - 16:00 — Augustin Fresnel Room

Session 4A28

Plasmonics and nano-optics

14:00 : Silver nanowires - remotely excited (bio)sensors

Michal Cwik¹, Karolina Sulowska², Dorota Buczynska¹, Ewa Rozniecka¹, Martyna Domagalska², Sebastian Mackowski², Joanna Niedziolka-Jönsson¹

¹Polish Academy of Sciences (Poland), ²Nicolaus Copernicus University (Poland)

We synthesized silver nanowires (AgNWs) with varying diameters and lengths by changing reducing agent. We found that choice of the reducing agent leads to nanowires with desired dimensions and thus plasmonic properties. For hybrid nanostructures consisting of AgNWs and photoactive proteins we observe that with increase of AgNWs diameter surface plasmon propagation length increases, yet fluorescence enhancement decreases. We show that synthesized AgNWs can be utilized for remote excitation of molecules localized hundred of microns from excitation spot.

14:15 : A refractive index sensor based on a Au inverted honeycomb lattice

Javier Rodriguez Alvarez¹, Lorenzo Gnoatto², Marc Martínez Castells³, Albert Guerrero⁴, Xavier Borrísé⁵, Arantxa Fraile Rodríguez¹, Xavier Batlle¹, Amílcar Labartaa¹

¹IN2UB - University of Barcelona (Spain), ²University of Barcelona (Italy), ³University of Barcelona (Spain), ⁴Institut de Microelectrónica de Barcelona (IMB-CNM, CSIC) (Spain), ⁵Catalan Institute of Nanoscience and Nanotechnology (ICN2) (Spain)

We present an efficient refractive index sensor consisting in a Au inverted honeycomb lattice. Numerical simulation showed high sensitivity values up to 99 nm/RIU for test layers of 50 nm. In addition, the figure of merit of the sensor detecting slight changes of the refractive index of a water medium at a fixed wavelength was 199 RIU-1. As an experimental proof of concept, the heterostructure was manufactured by electron beam lithography and the measured optical response agreed with the simulations.

14:30 : Towards perfect metallic behavior in optical resonant absorbing nanostructures

Clément Verilhac, Mathilde Makhsiyani, Riad Haidar, Jérôme Primot, Patrick Bouchon

Université Paris-Saclay (France)

Looking for a perfect metallic behavior is a crucial research line for metamaterials scientists. We propose a strategy based on a contrast of dielectric index to control losses in metals within waveguides and resonant nanostructures. This permits to tune the quality factor of the guided mode and of the resonant absorption over at least four orders of magnitude. This concept is applied to a practical design to finely control the localization of dissipation in an absorbing photonic structure.

14:45 : Plexcitons in Thermo-responsive Dextran-Graft-PNIPAM / Au Nanoparticles / CdTe Quantum Dots Nanosystem: Temperature Effects**Oleg Yeschhenko¹, Pavlo Khort¹, Nataliya Kutsevol¹, Olga Kapush², Volodymyr Dzhanan¹**¹Taras Shevchenko National University of Kyiv (Ukraine), ²National Academy of Sciences of Ukraine (Ukraine)

The temperature driven plasmon-exciton coupling in thermo-responsive dextran-graft-PNIPAM / Au nanoparticles / CdTe quantum dots (D-g-PNIPAM / Au NPs / CdTe QDs) nanosystem was studied. A large (0.84 eV) splitting of plexciton states was observed. The heating-cooling cycle of the aqueous solution of the studied nanosystem leads to a reversible quenching-recovery alteration of the QD photoluminescence. The quenching was rationalized as a result of an increased probability of nonradiative resonance energy transfer (RET) from CdTe QDs to Au NPs.

15:00 : Up-conversion luminescence activated by surface plasmon polaritons**Maciej Cwierzona, Karolina Sulowska, Michal Zebrowski, Sebastian Mackowski, Dawid Piatkowski***Nicolaus Copernicus University (Poland)*

In this work we discuss remote activation and detection of up-conversion photoluminescence, coming from submicron Er³⁺/Yb³⁺ co-doped nanocrystals (NCs) droplet, deposited locally at one end of long single silver nanowire. We show that different polarization of the laser beam as well as diameter of the nanowire change optical response of the nanocrystals in this polariton-mediated, remote up-conversion process.

15:15 : Semiconductor-based nanostructures for spectral filtering**Clément Maes¹, Grégory Vincent¹, Fernando Gonzalez-Posada Flores², Laurent Cerutti², Riad Haïdar¹, Thierry Taliercio²**¹ONERA (France), ²Université de Montpellier (France)

We present a theoretical study of a nanostructured guided-mode resonant (GMR) spectral filter operating in the long-wave infrared (LWIR) wavelength range. The component is made of III-V semiconductors: heavily n-doped InAsSb for the grating and GaSb for the waveguide of the GMR resonator.

15:30 : Spin-orbit coupling in dielectric nanostructures: a new recipe to realize sorting and boost diffusion at the nanoscale**Adria Canos Valero¹, Denis Kislov¹, Egor A. Gurvitz¹, Hadi K. Shamkhi¹, Alexander A. Pavlov², Dmitrii Redka³, Sergey Yankin⁴, Pavel Zemanek⁵, Alexander S. Shalin¹**¹ITMO University (Russia), ²INME - RAS (Russia), ³Electrotechnical University (Russia), ⁴LLC COMSOL (Russia), ⁵Institute of Scientific Instruments (Russia)

State-of-the-art microfluidics requires a variety of controllable components to perform operations inside microchambers. Brand-new nanophotonic approaches can significantly enhance existing capabilities via light-matter interactions. We propose a novel concept featuring dual on-chip functionality: boosted optically-driven diffusion and sorting. Well-designed high-index nanoantennae mediate spin-orbit coupling from a laser beam, producing subwavelength optical nanovortices that drive spiral motion of plasmonic nanoparticles, inducing fluid flow, enabling moving-part-free nanomixing. Moreover, we achieve precise sorting of gold NPs, demanded for on-chip separation and filtering.

15:45 : Plasmonic nano-patterning on optical fibers for optical neural interfaces**Rosa Mach-Batlle¹, Filippo Pisano¹, Di Zheng¹, Leonardo Sileo¹, Massimo De Vittorio², Ferruccio Pisanello¹, Cristian Ciraci¹**¹Istituto Italiano di Tecnologia (Italy), ²Università del Salento (Italy)

Optogenetics makes it possible to control the activity of neurons by light. Optical fibers are the most common device for delivering light to the brain, but they do not offer much control over the light distribution and intensity reaching the brain tissue. Here, we explore the possibility of designing a plasmonic nano-patterning at the fiber facet able to enhance the interaction between light and brain tissue in a controlled manner, which could lead to significant advances in the field.

14:00 - 16:00 — Ibn Al-Haytham Room

Session 4A29

Plasmonic Nanomaterials for Bio-diagnostics, Environmental Monitoring and Food Safety

Organized by: Lucia Petti

Chaired by: Lucia Petti and Massimo Rippa

14:00 : **Invited talk****Pixeled metasurface for multiwavelength detection****Valentina Di Meo¹, Alessio Crescitelli¹, Massimo Moccia¹, Annamaria Sandomenico², Angela M Cusano³, Marianna Portaccio⁴, Maria Lepore⁴, Vincenzo Galdi¹, Emanuela Esposito¹**¹University of Sannio (Italy), ²Institute of Biostructure and Bioimaging, National Research Council (Italy), ³CeRICT srl (Italy), ⁴University of Campania "Luigi Vanvitelli"(Italy)

We present a plasmonic biosensor based on SEIRA spectroscopy, which exploits resonant coupling between plasmonic nanoantennas and vibrational excitation of small molecules. Our platform features a large-area metasurface made of gold nanoantennas fabricated on a silicon substrate, comprising different macroregions "pixels". A single chip is capable of performing analysis from the region of functional groups to that of fingerprint. We detect a concentration as low as 86 pmol/L of a small molecule which is the best marker of vitamin D3.

14:20 : **Invited talk****Hybrid Plasmonic Nanomaterials: Functional Platforms for Bio and Food****Bruno Miranda, Ilaria Rea, Principia Dardano, Carlo Forestiere, Luca De Stefano***ISASI-CNR (Italy)*

We design, fabricate and analyze a hybrid nanoplasmonic device by using gold nanoparticles and polyethylene glycole. We characterize the absorption spectra of spherical gold nanoparticles embedded in polyethylene glycol diacrylate. The designed wearable platforms could detect specific target analytes in localized surface plasmon resonance and fluorescence modes. The proposed fabrication strategy represents a good candidate for becoming a functional platform for Bio and Food screening.

14:40 : **Invited talk****Photometric station for in-vitro diagnostic analysis using organic-based opto-electronic devices and photonic crystals****Giuseppe Nenna¹, Maria Grazia Maglione¹, Pasquale Morvillo¹, Tommaso Fasolino¹, Anna De Girolamo Del Mauro¹, Rosa Ricciardi¹, Riccardo Miscioscia¹, Giuseppe Pandolfi¹, Giovanni De filippo¹, Francesco Pascarella¹, Carla Minarini¹, Rosita Diana¹, Massimo Rippa², Lucia Petti², Paolo Tassini¹, Salvatore Aprano³, Giorgio Allasia³**¹ENEA, Italian National Agency for New Technologies (Italy), ²Università di Napoli Federico II (Italy), ³FOS S.p.A. (Italy)

This work describes, for the first time, the use of innovative organic optoelectronic devices and photonic crystals to prepare an absorbance apparatus for plate readers, for the invitro detection of antibodies or antigens in biological samples.

15:00 : **Invited talk****Towards label-free, implantable neuro-plasmonic probes****filippo Pisano¹, Antonio Balena¹, Muhammad Fayyaz Kashif², Marco Pisanello¹, Antonio Quattieri¹, Leonardo Sileo¹, Tiziana Stomeo¹, Antonella D'Orazio², Marco Grande², Ferruccio Pisanello¹, Massimo De Vittorio¹**¹Istituto Italiano di Tecnologia (Italy), ²Politecnico di Bari (Italy)

Optical fibers are widely applied to stimulate and monitor neural activity in deep brain regions expressing light-sensitive genetically-encoded sensors and actuators. Recent progress has demonstrated that these probes allow for endoscopic imaging and depth-resolved tissue interrogation. However, label-free sensing of neural events in vivo is still beyond the reach of currently available technologies. To remediate to this shortcoming, we have developed novel methods to fabricate multiple plasmonic bio-sensors on the optically active surface of flat-cleaved and tapered optical fibers.

15:20 : **Invited talk**

Plasmonic nanostructures for label-free detection of water contaminants

Massimo Rippa¹, Ambra Vestri¹, Radoslaw Kolkowski², Joseph Zyss³, Jun Zhou⁴, Lucia Petti¹

¹*Institute of Applied Sciences and Intelligent Systems .E. Caianiello. of CNR (Italy)*, ²*Center for Nanophotonics, AMOLF (The Netherlands)*, ³*Institut d'Alembert (France)*, ⁴*Ningbo University (China)*

In this work we design, fabricate and characterize novel periodic arrangements of gold nanoelements. We tested the detection performance of our nanostructures analyzing different concentrations in water of fipronil and imidacloprid insecticides using both the Localized Surface Plasmon Resonance (LSPR) and Surface Enhanced Raman Spectroscopy (SERS) sensing approach.

15:40 : **Invited talk**

Bioanalytics using plasmonic nanostructures

David Zopf, Angelina Pittner, Philipp Muller, Cornelia Reuter, Matthias Thiele, Ondrej Stranik, Andrea Csaki, Wolfgang Fritzsche

Leibniz Institute of Photonic Technology (IPHT) (Germany)

Novel requirements for bioanalytical methods emerge due to trends such as personalized medicine. Upon binding of molecules, the localized surface plasmon resonance (LSPR) of these structures is changed, and can be used as sensoric readout. We will show how plasmonic nanostructures provide sensing capabilities with the potential for ultrasensitive and robust assays in a high parallelization, and without the need for markers.

14:00 - 16:00 — Gaston Floquet Room

Session 4A30

Symposium II: New trends in nanophotonics and advanced materials

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

14:00 : **Invited talk**

Brillouin Spectroscopy in Optophononic Micropillars at 18 GHz

Anne Rodriguez, Edson Cardozo de Oliveira, Priya Priya, Abdelmounaim Harouri, Isabelle Sagnes, Luc Le Gratiet, Martina Morassi, Aristide Lemaitre, Loic Lanco, Pascale Senellart, Martin Esmann, Norberto Daniel Lanzillotti-Kimura

Centre de Nanosciences et de Nanostructures - C2N (France)

We measured the Brillouin spectrum on an elliptical optophononic micropillar resonator based on AIAs/GaAs superlattices designed to confine light and sound simultaneously. The ellipticity has associated two polarized modes used to discriminate the reflected laser and the Brillouin signal.

14:20 : **Invited talk**

Dynamics of Nanomechanical Metamaterials: Pico-vibrometry with Light and Electron Beams

Jun-Yu Ou, Tongjun Liu, Jinxiang Li, Dimitrios Papas, Eric Plum, Kevin F. MacDonald, Nikolay I. Zheludev

University of Southampton (United Kingdom)

We report on the detection and quantitative mapping of picometre (sub-atomic) amplitude, thermal (phonon-induced) and driven movements in photonic nanostructures, using light and electron beams. These techniques enable measurements of the dynamic mechanical properties that underpin the functionality of a growing range of micro/nano-opto-mechanical (meta)materials, devices, sensors and systems, and present new opportunities in the exploration of fundamental nonequilibrium (opto)mechanics.

14:40 : **Invited talk**

Slow-light dissipative Kerr solitons in coupled-cavity waveguides

Juan Pablo Vasco, Vincenzo Savona

Ecole Polytechnique Fédérale de Lausanne (Switzerland)

We study frequency combs and dissipative Kerr solitons in silicon coupled-cavity waveguides (CCW) with globally optimized dispersion at telecom wavelengths. The corresponding threshold for comb generation is

found to explicitly depend on the main CCW figures of merit, namely, mode volume, normal mode quality factor and slow-light group index. Our results set the CCW as a new paradigm for low-threshold comb generation via advanced dispersion engineering and slow-light enhancement.

15:00 : Invited talk

Topological Pillared Phononic Crystals: Edge States, Fano Resonance and Their Robustness Against Disorder

Wei Wang¹, Yabin Jin², Wan Wang², Bernard Bonello¹, Romain Fleury³, Daniel Torrent⁴, Yan Pennec⁵, Bahram Djafari Rouhani⁵

¹Sorbonne Université (France), ²Tongji University (China), ³EPFL (Switzerland), ⁴Universitat Jaume I (Spain),

⁵Université de Lille (France)

We study theoretically the topological properties of pillared phononic crystals. By breaking the space inversion symmetry in a honeycomb lattice, we show different topological phases emulating the analogs of quantum valley and spin Hall effects. Robust edge states with one-way propagation are demonstrated as well as a rich variety of refraction phenomena at the outlets. A robust topological Fano mechanical resonance is achieved in a pillared beam from the superposition of a dark and a bright edge mode.

15:20 : Invited talk

Enhancing chiral fields with arrays of achiral nanoparticles

T. V. Raziman

Eindhoven University of Technology (The Netherlands)

Chiral light-matter interaction forms the basis for molecular circular dichroism spectroscopy, optical spin manipulation, and optical torques. Magnifying chiral effects using nanophotonics requires preserving the chiral near field. We propose conditions and limits for enhancing chiroptical effects near achiral metasurfaces. We prove that a nanostructure cannot be universally optimal for different chiral applications. We also predict an analytical limit of maximum circular dichroism in highly evanescent Fourier orders. Our results establish guidelines for nanophotonic enhancement in diverse chiroptical applications.

15:40 : Invited talk

Integrated nonlinear photonics in AlGaAs-on-insulator devices

S. May¹, J. McPhillimy¹, C. Klitis¹, B. Guilhabert¹, M. Kues¹, M. D. Dawson², M. J. Strain², M. Clerici¹, M. Sorel¹

¹University of Glasgow (United Kingdom), ²University of Strathclyde (United Kingdom)

The heterogeneous integration of AlGaAs-on-insulator (AlGaAs-OI) has proven to be a powerful material platform for nonlinear optics. This talk will explore how chip-scale bonding and transfer printing techniques can be used for the fabrication of integrated photonic chips for highly efficient second and third order nonlinear interactions. Examples to be presented will include devices for second harmonic, super-continuum and four-wave mixing generation, as well as vertical geometries to engineer the interaction between different spatial modes.

14:00 - 16:05 — Lawrence Bragg Room

Session 4A31

Light-matter interactions in new materials and meta-architectures

Organized by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

Chaired by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

14:00 : Invited talk

Near-field directionality with higher order multipolar sources

Michela Florinda Picardi, Francisco Rodríguez Fortuño

King's College London (United Kingdom)

We reveal the near-field directionality properties of multipolar sources comprised of superpositions of electric

and magnetic dipoles and quadrupoles. We build a table of elementary sources comprised of the superposition of two multipoles, either both electric, both magnetic or one of each nature.

14:20 : Invited talk

Large-area optical metasurfaces: from plasmonic color routing to energy harvesting in 2D materials
Maria Caterina Giordano¹, Matteo Barelli¹, Andrea Mazzanti², Giuseppe Della Valle², Francesco Bua-
tier de Mongeot¹

¹ *Universitadi Genova (Italy)*, ² *Politecnico di Milano (Italy)*

The nanofabrication of nanoscale metasurfaces able to feature tunable optoelectronic response is crucial in view of sensing and flat-optics applications. Here the engineering of large-scale metasurfaces based either on self-organized plasmonic nanoantennas or on two-dimensional (2D) semiconductor layers will be shown. These templates enable accurate tuning of localized plasmon resonances and can feature broadband color routing properties with strong impact in flat-optics, photonics and sensing applications.

14:40 : Invited talk

Designing with spins: magnonic metamaterials based on nanoengineered spin textures
Edoardo Albisetti¹, Silvia Tacchi², Raffaele Silvani³, Giuseppe Scaramuzzi¹, Simone finizio⁴, Sebastian
Wintz⁴, Christian Rinaldi¹, Matteo Cantoni¹, Jorg Raabe⁴, Giovanni Carlotti³, Elisa Riedo⁵, Riccardo
Bertacco¹, Daniela Petti¹

¹ *Politecnico di Milano (Italy)*, ² *Istituto Officina dei Materiali del CNR (CNR-IOM) (Italy)*, ³ *Università di Perugia (Italy)*, ⁴ *Paul Scherrer Institut (Switzerland)*, ⁵ *New York University (USA)*

Magnonics seeks to control the excitation, propagation and transduction of spin waves, i.e. propagating perturbations in the arrangement of spins in magnetic materials, for information processing. Here, we present our work on the manipulation of spin waves in nanoengineered magnetic materials. first, we present a technique, tam-SPL, for nanopatterning spin textures in magnetic multilayers. Then, we discuss the use of spin textures for controlling the emission, propagation and interference of spin waves, aiming to develop energy-efficient wave-based computing platforms.

15:00 : Solar light with sub-microns hyperboloids non-imaging light concentrators arrays

Ashish Prajapati, Gil Shalev

Ben-Gurion University of the Negev (Israel)

Metamaterials based on arrays of subwavelength dielectric structures have recently proved to be a viable research tool towards the realization of various photonic devices. In the current study we introduce a new approach towards efficient light trapping and broadband absorption of solar radiation based on silicon surface arrays composed of subwavelength trumpet non-imaging light concentrators (henceforth, trumpet arrays). We show superior broadband absorption of solar radiation in trumpet arrays compared with that of the optimized NP arrays (~26% enhancement).

15:15 : Photo-magnetic recording with L-band ultrashort laser pulses in dielectric medium

Antoni Frej, Tomasz Zalewski, Andrzej Stupakiewicz

University of Bialystok (Poland)

We experimentally demonstrated the possibility of the potential use of ultrashort laser pulses at telecommunication L-band to all-optical magnetic recording. A single linearly polarized laser pulse near 1590 nm wavelength switch the magnetization between two states in Co-doped YIG thin transparent film. Changing the linear polarization of the laser pulse, we can write-erase magnetic area in the sample.

15:30 : Single-shot time-resolved imaging of all-optical ultrafast photo-magnetic switching

Tomasz Zalewski, Antoni Frej, Andrzej Stupakiewicz

University of Bialystok (Poland)

The understanding of fundamental mechanism allowing for all-optical magnetization switching in femtosecond time scale is a key for providing novel outperforming applications. Recently, it has been discovered that only by a single laser pulse, extremely fast (about 20 ps), reversible and repeatable photo-magnetic switching in Co-doped yttrium iron garnet films can be obtained. Here, we demonstrate the experimental technique enabling visualization of magnetization switching at femtosecond time scale using single-shot time-resolved magneto-optical imaging in YIG:Co films.

15:45 : Invited talk

Resonant Heat Transfer in Nanophotonics Driven Thermal Water Desalination**Alessandro Alabastri, Pratiksha D. Dongare, Oara Neumann, Peter Nordlander, Naomi J. Halas***Rice University (USA)*

Light absorbing nanoparticles can efficiently convert electromagnetic radiation into heat in nanoscale regions. This photothermal effect can be exploited to locally increase the temperature of water, promoting its evaporation and thus purification. Optothermal nonlinearities and resonant thermal phenomena have been uncovered to improve the efficiency of this process. Here we show how these concepts together can be engineered to achieve fresh water production rates up to ~ 20 L/(m².day) under standard solar irradiation.

14:00 - 15:30 — Rene Descartes Room**Session 4A32****Metasurfaces and thermal metamaterials****14:00 : Dynamic optical MEMS Metasurfaces****Paul Thrane¹, Chao Meng², Christopher A. Dirdal³, Jo Gjessing³, Fei Ding², Sergey L. Bozhevolnyi²**¹*University of Southern Denmark (Norway)*, ²*University of Southern Denmark (Denmark)*, ³*SINTEF Microsystems and Nanotechnology (Norway)*

We present a versatile platform for achieving dynamical metasurfaces by combining micro-electro-mechanical systems (MEMS) with plasmonic nanostructures. By varying the separation of a gold piezoelectric MEMS mirror and gold nanostructures we adjust the behavior of the reflective metasurface. The concept is demonstrated for several cases, including switchable gratings and lenses. Few optical MEMS metasurfaces have been demonstrated previously, and this platform has the advantage of allowing full freedom in the metasurface design and decoupling the MEMS and nanostructure fabrication.

14:15 : Ion-beam-doped transparent conductive oxides for metasurface applications**Alexander Koch¹, Hongyan Mei², Jura Rensberg¹, Martin Hafermann¹, Jad Salman², Chenghao Wan², Ray Wambold², Mikhail Kats², Carsten Ronning¹**¹*Friedrich Schiller University (Germany)*, ²*University of Wisconsin-Madison (USA)*

Doped transparent conductive oxides have gained a lot of attention for applications in plasmonics and nanophotonics due to their low optical loss, metal-like behavior, tailorable optical properties, and well-established fabrication procedures. N-type doped zinc oxide, like gallium-doped ZnO is attractive because its permittivity can be engineered over a broad wavelength range across the infrared. Here, we demonstrate how the optical properties of ZnO can be modified by doping with gallium using a commercial focused ion beam system and post-implantation annealing.

14:30 : Enhanced Evanescent field Confinement Driven by Bound States in the Continuum**Sergei Lepeshov, Andrei Bogdanov***ITMO University (Russia)*

Here, the enhancement of electromagnetic field confinement in an all-dielectric metasurface is demonstrated. The enhanced confinement is achieved when the polarization singularity, corresponding to accidental bound states in the continuum, moves to the domain of evanescent fields (under the light line). Such a hybridization of the bound states and evanescent waves results in the 70-fold increase of the electric field enhancement on the top of the metasurface and boosting of the electric field localization.

14:45 : Broadband high-efficiency achromatic meta-device based on phase and dispersion independently controlled metasurface**Wenye Ji, Paul Urbach***Delft University of Technology (The Netherlands)*

Achromatic devices have wide application prospect in radar or fields. However, chromatic aberration and limited bandwidth restrict their development. In this paper, we achieve broadband high-efficiency transmission achromatism based on metasurface. In order to verify the theory and design method, we design an achromatic anomalous deflector and fabricate the sample. The experimental results are in good agreement with theory. Our findings provide valuable theory and strategy for achromatic device design.

15:00 : Investigation of electromagnetic coupling between the antenna and split-ring-based metasurface in CMOS technology**Alexander Vladimirovich Chernyadiev¹, Dmytro But¹, Cezary Kolaciński¹, Kestutis Ikamas², Alvydas Lisauskas¹**¹*Institute of High Pressure Physics PAS (Russia)*, ²*Vilnius University (Lithuania)*

In this contribution we investigate the electromagnetic coupling between the 350~GHz resonant antenna and the metasurface constructed from an array of split-ring structures which are monolithically integrated using a 180~nm silicon-based CMOS technology. We examine how the coupling between these structures and the number of split-rings affects the high frequency impedance of the whole system. The efficient control of impedance is important for designing passive and active terahertz components, such as tunable detectors and sources.

15:15 : Designing and Fabrication of efficient nano-materials for thermoelectric generator**Wiqar Hussain Shah***International Islamic University (Pakistan)*

The electrical and thermal properties of the doped Tellurium Telluride chalcogenide nano-particles are mainly characterized by a competition between metallic (hole doped concentration) and semi-conducting state. nano-particles, prepared by solid state reactions in sealed silica tubes and ball milling method. These increasing behavior of Seebeck coefficient leads to high power factor which are increases with increasing temperature and Sn concentration, because of lowest electrical conductivity but its power factor increases well with increasing temperature.

Break

16:00 - 16:30

16:30 - 17:25 — Victor Veselago Room**Session 4A33****Symposium II: New trends in nanophotonics and advanced materials**

Organized by: Junsuk Rho, Hakjoo Lee and Namkyoo Park

16:30 : Invited talk**Plasmonic aerosols to govern light****Jake Fontana, Jeff Geldmeier, Paul Johns, Nicholas J. Greybush, Jawad Naciri***U.S. Naval Research Laboratory (USA)*

We show the experimental demonstration of a plasmonic aerosol and discuss the significance of uniting the fields of plasmonics and aerosols. We find that the aerosols are optically homogeneous, thermodynamically stable, with wide wavelength tunability, and extremely large sensitivities to their environment. Plasmonic aerosols may therefore provide a novel medium to govern light-matter interactions, thereby opening up innovative opportunities.

16:50 : Invited talk**Graphene Plasmonic Oscillators for Terahertz Light Emission****Yuyu Li, Pablo Ferreyra, Anna Swan, Roberto Paiella***Boston University (USA)*

We report the measurement of electrically-driven narrowband THz light emission from graphene. The underlying radiation mechanism involves the generation of hot carriers under current injection and their subsequent energy relaxation through the excitation of plasmonic resonances in graphene nanoribbons. Free-space THz radiation is then emitted by the resulting collective oscillations of the graphene electron gas. These results are technologically relevant for the development of highly miniaturized and broadly tunable THz radiation sources.

17:10 : Wavefront Control of Light Emission from Halide Perovskite Metamaterials

Yixin Chen, Jinze Cai, Xuezi Ma, Shoufeng Lan, Zi Jing Wong
(USA)

Wavefront Control of Light Emission from Halide Perovskite Metamaterials

16:30 - 17:30 — Tatsuo Itho Room

Session 4A34

Plasmonics and nano-optics

16:30 : Casimir induced instabilities at metallic surfaces and interfaces

Kun Ding¹, Daigo Oue¹, C. T. Chan², J. B. Pendry¹

¹Imperial College London (United Kingdom), ²Hong Kong University of Science and Technology (China)

Surface plasmons subject to a surface distortion split asymmetrically in energy resulting in a net lowering of zero-point energy. This is because surface plasmon eigenvalues are the square of frequencies, a statement generally true for electromagnetic excitations. We utilize the conformal mapping method to demonstrate asymmetric splitting under surface corrugations can lead to a decrease in zero-point energy of a single corrugated metallic surface and drive the instability of a metallic thin film and cavity.

16:45 : Resonant Plasmon Enhancement of Light Emission from CdSe-Based Nanocrystals on Au Nanodisk Arrays

Alexander Milekhin¹, Mahfujur Rahaman², Ilya A. Milekhin², Tatyana A. Duda³, Ekaterina E. Rodyakina³, Kirill Anikin³, Roman B. Vasiliev⁴, Volodymyr Dzhagan⁵, Alexander V. Latyshev⁶, Dietrich R.T. Zahn²

¹Novosibirsk State University (Russia), ²Chemnitz University of Technology (Germany), ³A. V. Rzhanov Institute of Semiconductor Physics (Russia), ⁴Moscow State University (Russia), ⁵V. E. Lashkaryov Institute of Semiconductor Physics (Ukraine), ⁶A. V. Rzhanov Institute of Semiconductor Physics (Russia)

We report on resonant plasmon enhancement of Raman scattering (RS) and photoluminescence (PL) from colloidal CdSe-based nanocrystals (NCs) deposited on Au nanodisk arrays. The enhancement factors of RS and PL depend on the nanodisk size and reach maximal values at 75 and 7, respectively. These maxima are observed for the nanodisks having plasmon energy close to the interband transition energies in semiconductor NCs. Tip-Enhanced RS allowed us to perform local spectral analysis of CdSe NCs with nanometer spatial resolution.

17:00 : Controlling surface plasmons using all-metallic structures

Joseph Riley, Noel Healy, Victor Pacheco Peña

Newcastle University (United Kingdom)

In this work, we propose a method to manipulate and control the propagation of surface plasmons using all-metallic plasmonic structures. In this technique, a metallic structure is inserted within a semi-infinite block made of a different metal. The effective medium approach at each air-metal interface is exploited to control the propagation direction of surface plasmons. As an example, a plano-convex lens is designed and evaluated demonstrating the ability to focus the incident surface plasmons to the desired focal distance.

17:15 : Plasmonic metasurfaces for magnetic field enhancement at THz frequencies

Katarana Rovenska¹, Martin Hrton¹, Filip Ligmajer¹, Peter Kepic¹, Vlastimil Krapek¹, Rainer Hillenbrand², Tomas Sikola¹

¹Brno University of Technology (Czech Republic), ²CIC Nanogune (Spain)

Enhanced magnetic fields at terahertz frequencies are indispensable in many situations where analysis or modification of magnetic properties of matter is needed. Here we study a plasmonic metasurface formed by diabolical antennas, which exhibits such magnetic field enhancement, and using numerical simulations and subsequent terahertz time-domain spectroscopy we confirm a clear relationship between the antenna geometry and the spectral shape and position of the plasmonic resonance.

16:30 - 17:30 — Ibn Al-Haytham Room**Session 4A35****Plasmonic Nanomaterials for Bio-diagnostics, Environmental Monitoring and Food Safety**

Organized by: Lucia Petti

Chaired by: Lucia Petti and Massimo Rippha

16:30 : Invited talk**Light-induced Aggregation of Gold Nanorods on Graphene controlled by Radiation Pressure for SERS Detection of Biomolecules****Antonino Foti, Maria Grazia Donato, Onofrio Maria Marago, Pietro Giuseppe Gucciardi***CNR-IPCF (Italy)*

Radiation pressure is used to push gold nanorods on multilayered graphene and create hybrid active surfaces for Surface-Enhanced Raman Spectroscopy (SERS) in liquid. As a proof of concept, ultrasensitive detection of bovine serum albumin is shown, and the aggregation kinetics is studied as a function of the irradiation time. Our results enlarge the spectrum of materials that can be used for optical aggregation and SERS detection of biomolecules, highlighting the importance of controlling the physical properties of the surfaces

16:50 : Invited talk**Plasmonic tools to study interactions relevant for food science and eco/nanotoxicology****Duncan Sutherland***Aarhus University (Denmark)*

Nanosized sensors based on the local refractive index sensitivity of plasmonic nanostructures can be applied to study interactions occurring at biointerfaces relevant in understanding protein effects in eco and nanotoxicology and in food science. Plasma protein interactions with silver nanoparticles modulate the biotransformation of silver to silver sulfide while saliva proteins coupled to plasmonic sensors can be used to quantify astringency through interactions with polyphenols.

17:10 : Invited talk**Enhanced Refractive Index Imaging Based on Quasi-Bound States in the Continuum****Silvia Romano¹, Maria Mangini², Stefano Cabrini³, Erika Penzo³, Anna Chiara De Luca², Ivo Rendina¹, Vito Mocella¹, Gianluigi Zito¹***¹National Research Council ISASI (Italy), ²National Research Council IBBC (Italy), ³Lawrence National Laboratory of Berkeley (USA)*

We achieve a condition of enhanced hyperspectral refractometric, imaging on all-dielectric photonic crystal slabs, (PhCS) using a scheme of surface-enhanced fluorescence, (SEF) combined with refractometric sensing. The scheme is based on two high-Q resonances in proximity of bound, states in the continuum (BICs). The mechanism of quasi-, BIC spatially-variant gain, tracked by quasi-BIC refractometric, sensing, can find application in many fields for monitoring, physical and biochemical processes. In this study it is applied for surface cell analysis.

16:30 - 17:00 — Lawrence Bragg Room**Session 4A36****Light-matter interactions in new materials and meta-architectures**

Organized by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

Chaired by: Nicolò Maccaferri, Vincenzo Caligiuri and Mario Miscuglio

16:30 : Ultrastrong interaction between plasmons and photons in a terahertz photonic crystal cavity
Fanqi Meng, Mark D. Thomson, Bernhard Klug, Dovile Cibiraite, Qamar UI-Islam, Hartmut G. Roskos
Goethe-University Frankfurt (Germany)

Realization of strong interaction between photons and the plasmons of metamaterials in a cavity is of practical interest. Here, by employing a photonic crystal cavity and a metallic metamaterial, we achieve strong light-matter interaction between plasmons and photons in the terahertz frequency range. The measured Rabi splitting indicates that the ultrastrong coupling regime is reached. We demonstrate the nonlocal collective character of the interaction. Moreover, ultrastrong coupling between Babinet-complementary metamaterials and photons is also observed and explained.

16:45 : Ultrafast reflectance switching based on artificial epsilon-near-zero modes in a metal-insulator-metal nanocavity

Joel Kuttruff¹, Denis Garoli², Jonas Allerbeck¹, Roman Krahn², Antonio De Luca³, Daniele Brida⁴, Vincenzo Caligiuri³, Nicolo Maccaferri⁴

¹University of Konstanz (Germany), ²Istituto Italiano di Tecnologia (Italy), ³Universita della Calabria (Italy), ⁴University of Luxembourg (Luxembourg)

Ultrafast control of light-matter interactions is crucial in view of new technological frontiers of information processing. Here, we exploit high-quality-factor artificial epsilon-near-zero (ENZ) modes of a metal-insulator-metal nanocavity to establish all-optical, ultrafast control of light reflectance. Pumping the system at its high-energy ENZ mode, the low-energy mode strongly redshifts because of the transient increase of the local dielectric function, which leads to a sub-3-ps control of the reflectance with a modulation depth of about 120%.

Index

Abdeddaim Redha : 2A21
 Abdollahi Mohammad A. : 1P2
 Abdulhalim Ibrahim : 1P2
 Abebe Muluneh G. : 3A14
 Abraham Ekeroth Ricardo M. : 2A19, 2A28
 Abujetas Diego R. : 1A15
 Achouri Karim : 4A5
 Acuautla Monica : 1A17
 Adam Pierre Michel : 2A21
 Adibi Ali : 3A32, 3A41
 Aers Geoffrey C : 1A30
 Agircan Hasret : 4A17
 Agouram Said : 1A18
 Agrawal Amit : 1A22, 3A12, 3A34
 Agreda Adrian : 4A18
 Aharonovich Igor : 4A12
 Ahmed Gazi A. : 3P1
 Ahmed Shakeel : 2P2
 Ahmed Waqas : 2A8
 Ahmed Waqas W. : 2A14, 2A37
 Aizpurua Javier : 3A23, 4A18
 Akbari Mahmood : 1P2
 Akbıyık Alp : 3P2
 Akiki Rock : 4P1
 Akpan Usenobong : 2A34
 Aktas Djeylan : 4A23
 Al Haddadin Zaid : 3A36
 Al Maruf Rubayet : 1A26
 Al Rsheed Anwar : 3P1
 Al Saffar Norah A. : 3P2
 Al-Hamry Ammar : 3A28
 Alabastri Alessandro : 4A31
 Alarcon Llado Esther : 2A20
 Albert Matthias : 3A28
 Albisetti Edoardo : 4A31
 Aldossary O. M. : 3A37
 Aldossary Omar M. : 3P1
 Alecci Marcello : 1A19
 Algarni Hamed : 3P2, 3P2
 Ali Abubakar Sani : 1P2
 Ali Munazza Zulfiqar : 3A27
 Alibart Olivier : 3A21
 Aliev Ali : 2A24
 Allasia Giorgio : 4A29
 Allerbeck Jonas : 4A36
 Alonso-Gonzalez Pablo : 1A5, 2A3
 Alonso-Ramos Carlos : 1A22, 3A21
 Alpat Behcet : 3A23
 Alqahtani Mohammed : 3P2, 3P2
 Altland Alex : 3A4
 Altuzarra Charles : 3A35
 Alvarez Fernandez Alberto : 1A14
 Alvarez Ortiz Maria Monserrat : 1P2
 Alvarez-Fernandez Alberto : 4A8
 Alzaidy Ghadah : 2P2
 Amanaganti Sushanth R. : 4A12, 4P1
 Ambrosio Antonio : 3A24, 3A30
 Amelchenko Maria : 2P2
 Amirsalari Abdolvahab : 2P2
 Amo Alberto : 1A20
 An Sensong : 2A27
 Anand Srinivasan : 2A2
 Anders Krzysztof : 3P1
 Andrianov Evgeny : 4A16
 Andrés Daniel : 2A17, 3P1
 Angelskar Hallvard : 3A28
 Angelucci Francesco : 2A19
 Angeris Guillermo : 2A32
 Anikin Kirill : 4A34
 Anopchenko Aleksei : 3A32
 Antonietti Markus : 2A13
 Antosiewicz Tomasz J. : 1A28
 Anuszkiewicz Alicja : 1A5
 Apostoluk Alexandra : 3A22
 Aprano Salvatore : 4A29
 Aradian Ashod : 1A22
 Arai Yukinaga : 1A3
 Arbouet Arnaud : 1A12, 3A25
 Ardini Matteo : 2A19
 Ardizzone Vincenzo : 3A1
 Argiolas S. : 4A3
 Arias-González Felipe : 4A26
 Arik Kamalodin : 1P2
 Arlou Yauheni : 4A26
 Armelles Gaspar : 4A19
 Arreola Arturo : 4A27
 Arriaga Jesus : 3P2
 Arruda Tiago J. : 1A22
 Aryshev Alexander : 2P2
 Asano Takashi : 3A3
 Ashida Yuto : 2A5
 Askarpour Amir Nader : 1P2
 Assadillayev Artyom : 1P1
 Atlasbaf Zahra : 1P1
 Atmakuri Madhunika : 4A12, 4P1
 Atwater Harry : 1A6
 Atwater Harry A. : 2A22, 2A40
 Atzrodt Heiko : 1A9
 Auad Yves : 3A25
 Auernhammer Gunter K. : 4A24
 Auslander Mark : 1A28, 4P1
 Auyeung Raymond : 1A30
 Avalos-Ovando Oscar Rodrigo : 1A23
 Avishan Nardin : 3P2
 Avrahamy Roy : 1A28, 4P1
 Ayuso David : 2A24
 Azad Abul K. : 2A38
 Baba Akira : 3A3
 Babani Suleiman : 1P2
 Babicheva Viktoriia : 2A22, 3A36
 Babiker M. : 3A40
 Babiker Mohamed : 3A30

Bachelard Romain : 1A22
 Bachelot Renaud : 2A13
 Badloe Trevon : 4A1
 Bae Kideok : 2A7
 Baez Gabriela : 4A27
 Baffou Guillaume : 3P1
 Bagheri Amirmasood : 3P2
 Bahl Mayank : 2P2
 Bajcsy Michal : 1A26
 Baldi Andrea : 3A25
 Balena Antonio : 4A29
 Balestri Dario : 3A5
 Bali Rantej : 3A16
 Ballantine Kyle : 3A37
 Ballarini Dario : 3A1
 Bancerek Maria : 1A28
 Banerjee Debasish : 3A26
 Bang Hyunsung : 3A11
 Bardyszewski Witold : 3A40
 Barelli Matteo : 4A31
 Barnard Edward S. : 4A23
 Barnes Bill : 3A39
 Barois Philippe : 1A22, 1A24, 2A21, 4A8
 Baron Alexandre : 1A22, 1A24, 2A21, 4A8
 Barton David R. : 3A36
 Bartosewicz Bartosz : 2P1, 2P2
 Basani Jasvith Raj : 1P1, 4P1, 4P1
 Bason Mark G : 3A35
 Battle Xavier : 4A28
 Battie Yann : 2A12
 Baturina Olga : 3A33
 Baudrion Anne Laure : 2A21
 Bauman Stephen J. : 2P2
 Baumberg Jeremy J. : 4A21
 Baxamusa S. : 1A25
 Beal Jeremie : 2A21
 Becker Sven : 3A21
 Bedu Frederic : 2A21, 4A4
 Begou Thomas : 2A21
 Bek Alpan : 3P2
 Belardini Alessandro : 1A5
 Belkin Mikhail A. : 3A12
 Bellessa Joel : 3A1, 4A2, 4P1
 Bello Frank : 1A15
 Belosludtsev Alexandr : 1A18
 Belotelov Vladimir : 2A1
 Belov Pavel : 1P1
 Ben-Abdallah Philippe : 4A1
 Benedetti Stefania : 2A9
 Benedikovic Daniel : 1A22, 3A21
 Benisty Henri : 2A14
 Bennett Anthony J. : 1A20
 Benoit Jean-Michel : 3A1, 4A2
 Berakdar Jamal : 1A19, 3A20
 Bercioux Dario : 1A11
 Bercu Nicolas : 1P2
 Berezovska Nataliya : 1P2
 Bergamini Andrea E. : 1A17
 Bergamini Luca : 4A18
 Bergman David J : 1A5
 Berini Pierre : 2A26
 Bertacco Riccardo : 4A31
 Berte Rodrigo : 4A10
 Berthelot Johann : 2A21
 Bertoncini Andrea : 3A33
 Bertuch Thomas : 1P1
 Berté Rodrigo : 3A34, 4A23
 Bescond Marc : 1A8
 Bessonov Vladimir : 2A37
 Besteiro Lucas V. : 1A23, 1A26
 Betancourt Diego : 1P1
 Betz Fridtjof : 2A17
 Bhardwaj Amit : 4P1, 4A19
 Bi Chenghao : 4A7
 Bi Lei : 3A7, 3P1, 4A19
 Bi Yangang : 1A20, 1A20
 Biagioni Paolo : 4A18
 Bidault Sebastien : 2A21
 Biehs Svend-Age : 4A23
 Biffi Giulia : 4A12, 4A24
 Binkowski Felix : 2A17
 Bir Anastasia : 2P2
 Bisio Francesco : 1A23, 2A9, 2A9, 4A8, 4A17
 Bittrich Eva : 1A23
 Blake Jolie : 3A17
 Bleu Olivier : 3A1
 Bliokh Konstantin : 1A7
 Bloch Jacqueline : 1A20
 Blugel Stefan : 2A15
 Bo Fang : 3A9
 Bodunov Evgeny N. : 4A2
 Boehm Gerhard : 3A12
 Boer-Duchemin Elizabeth : 1P1
 Boeuf Frédéric : 1A22
 Bogdanov Andrei : 4A32
 Bogdanov Andrey : 2A4
 Bogdanowicz Janusz : 3A5
 Boggiano Hilario D. : 4A23
 Bohite Omkar : 2P1
 Bohm Julian : 2A33
 Bohn Justus : 3A39
 Boisron Olivier : 2A21
 Bok Eun : 1A8
 Bollani Monica : 4A18
 Boltasseva Alexandra : 1A14, 3A39
 Boltersdorf Jonathan A : 3A33
 Bonello Bernard : 1A17, 4P1, 4A30
 Bonn Mischa : 4A17
 Bonod Nicolas : 2A21
 Booth Tim : 1P1
 Bories Serge : 3P2
 Borisov Vladislav : 4A26
 Borrisé Xavier : 4A28
 Bosman Michel : 1A10
 Botey Muriel : 2A8, 2A14, 2A37
 Both Steffen : 3A22

Bouchon Patrick : 3P2, 3A35, 4A28
 Bouhelier Alexandre : 1A4, 4A18
 Bouras Khedidja : 2P2
 Bouras Mounir : 2P2, 3P1
 Boutinguiza Mohamed : 4A26
 Bouzouad Mouloud : 2P2
 Bowen Patrick : 2A27
 Boyd Robert W. : 2A29
 Boyd Stephen : 2A32
 Bozhevolnyi Sergey L. : 4A32
 Bozhko Dmytro A. : 1A16
 Bradlyn Barry : 1A11
 Bragas Andrea : 3A34
 Bragas Andrea V. : 4A23
 Brar Victor : 2A4
 Brasse Yannic : 1A23
 Braun Kai : 1A13
 Braun Paul V. : 3A32
 Bravo-Abad Jorge : 2P1
 Brener Igal : 3A39
 Brennan Grace : 1A22, 4A22
 Bretel Rémi : 1P1
 Bricchi Beatrice Roberta : 2A9
 Brida Daniele : 4A36
 Broch Laurent : 2A12
 Brongersma Mark L. : 1P1, 3A5
 Brulon Cyprien : 3A35
 Brunelli Matteo : 3A15
 Bruno Vincenzo : 3P2, 3A39
 Buatier de Mongeot Francesco : 4A31
 Buchnev Oleksandr : 1A18, 3A1
 Buchwald Walter R. : 3A39
 Buczynska Dorota : 4A28
 Buczynski Ryszard : 1A5
 Budai Judit : 1A18
 Budich Jan C. : 3A4
 Budner Bogusław : 2P2
 Burger Johannes : 4A20
 Burger Sven : 2A17
 Burguete-Lopez Arturo : 1P2, 1P2
 Burokur Shah Nawaz : 2A10, 2A10, 2A15, 4A5
 Busch Mark : 3A5
 But Dmytro : 4A32
 But Dmytro B. : 1A29
 Bychkov Igor : 1P2
 Bösch Cyrill : 3A10
 Cabrini Stefano : 3A34, 4A35
 Caglayan Humeyra : 3A5
 Cai Jinze : 4A33
 Caironi Mario : 3A16
 Calajò Giuseppe : 3A15
 Caligiuri Vincenzo : 4A12, 4A24, 4A36
 Caloz Christophe : 2A35, 4A23
 Calzolari A. : 1A12
 Camayd-Munoz Phil : 2A22
 Camayd-Munoz Philip : 2P2
 Cambiasso Javier : 1P1
 Campbell Sawyer D. : 2A32
 Campion James : 1A28
 Canales Claudio : 1A28
 Canepa Maurizio : 1A23, 2A9, 4A8, 4A17
 Canepa Paolo : 4A17
 Canos Valero Adria : 4A28
 Cantoni Matteo : 4A31
 Cao Shuiyan : 1P1
 Cao Yaoyu : 3A18
 Capasso Federico : 1A21, 1A31
 Capers James : 2P1
 Cardin Andrew E. : 2A38
 Cardozo de Oliveira Edson : 4A30
 Carletti Luca : 1A19, 4A24
 Carlotti Giovanni : 4A31
 Carminati Remi : 1A14
 Carollo Angelo : 3A15
 Casals Blai : 2A34
 Cassan Eric : 1A22, 3A21
 Castilla Marion : 3A13
 Castillo Miguel : 3A22
 Catchpole Kylie : 3P1
 Catellani A. : 1A12
 Catone Daniele : 2A9
 Cavalleri Ornella : 4A17
 Celano Umberto : 3A5
 Celebrano Michele : 3A5, 4A18
 Cerutti Laurent : 4A28
 Cervera Francisco : 3A37
 Cesca Tiziana : 1A5
 Chaabani Wajdi : 2A21
 Chacon Reinaldo : 1A4
 Chalimah Siti : 2A1
 Chan C. T. : 2A36, 4A1, 4A34
 Chan Che Ting : 1A11, 3A18, 3A20
 Chang Chieh-Chun : 3P1
 Chang Huan-Cheng : 2P1
 Chang Sih-Wei : 2P1
 Chao Pengning : 2A32
 Chapman Craig T. : 2A31
 Charipar Kristin : 1A30, 2P2
 Charipar Nicholas : 1A30, 2P2
 Charley Anne L. : 3A5
 Charra Fabrice : 2A13
 Chauhan Ankit : 4A25
 Chavel Pierre : 2A17
 Cheben Pavel : 1A22, 3A21
 Chehaidar Abdallah : 2A21
 Chekhov Alexander : 3P2
 Chen Bigeng : 4A18
 Chen Chen : 4A1
 Chen Chung-Chia : 3A11
 Chen Hong : 3A7
 Chen Hou-Tong : 2A38
 Chen Hsuen-Li : 2P1
 Chen Hui : 1A27, 1A27
 Chen Jian : 2A11
 Chen Jie : 1A8
 Chen Jin-Ju : 3A28

Chen Kuo-Ping : 3A3, 3A3
 Chen Lei : 1A6
 Chen Liang : 3A34
 Chen Lifeng : 2P2
 Chen Lin X. : 2A31
 Chen Lu : 3A12
 Chen Oliver Y. : 2P1
 Chen Parry : 1A5
 Chen Qin : 3A7
 Chen Rui : 2A11, 3A6
 Chen Ruipin : 3A6
 Chen Shangzhi : 3A17
 Chen Shuqi : 1A20, 1A20
 Chen Ting-Wei : 1P1
 Chen Wenjie : 2A11, 3A6
 Chen Xianfeng : 3A9, 3A10
 Chen Xianzhong : 4A20
 Chen Xiaoshuang : 3A19, 4A9
 Chen Xingqiao : 4A17
 Chen Xu : 1A6
 Chen Yangyang : 1A27
 Chen Yen-Wei : 2P1
 Chen Yixin : 4A33
 Chen Yujie : 1P1
 Chen Yuntian : 1A11, 3A18
 Chen Yushao : 2P2
 Chen Ze-An : 3A6
 Chen Zhigang : 1A4
 Cheng Hengbin : 3A7
 Cheng Hua : 1A20, 1A20
 Cheng Jian-chun : 2A14
 Cheng Jianchun : 4A13
 Cheng Ming : 4A11
 Cheng Szu-Cheng : 1P1
 Cheng Wen-Hui : 1A6
 Cheng Ying : 1A9
 Chenot Stephane : 2A21
 Cheref Yannis : 3A29
 Cherkashin Nikolay : 3A25
 Chernov Alexander : 2A1
 Chernyadiev Alexander Vladimirovich : 4A32
 Chevrier Kevin : 3A1
 Chew Soo Hoon : 2P1
 Chiang Yan Kei : 2A18
 Chioar Ioan-Augustin : 4A12
 Chirvony Vladimir S. : 3P2
 Chistensen Johan : 1A9
 Chittur K. Subramaniam : 4A19
 Cho Kyung-Sang : 2A7
 Cho Sung-Sil : 2P1
 Choa Pengning : 3A38
 Chochol Jan : 4P1
 Choi Duk-Yong : 4A15
 Choi Jun Hee : 2A3
 Choi Sungho : 2P1
 Choo Hyuck : 2A7, 2A11, 4A1, 4A4
 Chou Yu Cheng : 4A15
 Chowdhury Brinta : 1P2
 Chowdhury Sarah N. : 1A14
 Choy Jennifer T. : 2A38
 Choy Wallace : 4A7
 Christensen Johan : 1A9, 2A14, 2A14, 3A37
 Christensen T. : 3A29
 Christiansen Rasmus E. : 3A38
 Christodoulides Demetrios : 2A33
 Chu Anshi : 2A2
 Chu Zeshi : 3A19
 Chua Soo Jin : 3A19
 Chubich Dmytro : 4A2
 Chung Chin-Chien : 3A11
 Chung Hyeongju : 3A12
 Ciardi Matteo : 3A5
 Ciattoni Alessandro : 2A4
 Cibraite Dovile : 4A36
 Ciccarello Francesco : 2A23, 3A15, 3A15
 Ciraci Cristian : 2A1, 2A25
 Ciraci Cristian : 4A28
 Ciraci Cristian : 4P1
 Clarke Daniel : 2A13
 Clementi Marco : 3A39
 Cleret de Langavant Capucine : 3A29
 Clerici M. : 4A30
 Clerici Matteo : 3P2, 3A39
 Cluzel Benoit : 4A18
 Cohen Oren : 2A24
 Colas des Francs Gérard : 1A4, 1A12
 Coletti Camilla : 4A17
 Colleu-Banse Tanguy : 1P2
 Colom Remi : 2A17
 Comoretto Davide : 4A26
 Conti Claudio : 2A4
 Cookson Tamsin : 3A1
 Cornier Thibaut : 3A22
 Correa-Duarte Miguel A. : 3A13
 Cortes Cristian L. : 3A33
 Cortes Emiliano : 1A13
 Cortés Emiliano : 1P1, 3A34, 3A34, 4A23
 Costa da Silva Eduardo : 2P2
 Cours Robin : 3A25
 Courteille Philippe W. : 1A22
 Cox Joel Douglas : 1A15
 Crescitelli Alessio : 4A29
 Csaki Andrea : 4P1, 4A29
 Cuche Aurelien : 1A12
 Cuerva Miguel : 3P1
 Cui Tie Jun : 4A9
 Cumming David : 2P2, 3A39
 Cummins Cian : 4A8
 Curto Alberto G. : 1P2, 2P1, 3A13
 Cusano Angela M : 4A29
 Cwierzona Maciej : 4A28
 Cwik Michal : 4A28
 Cywinski Grzegorz : 1A29
 Czajkowski Krzysztof M. : 1A28
 D’Orazio Antonella : 4A29
 d’Orsonnens William : 2P1

Dahlin Andreas : 3A17
 Dai Daoxin : 2A7
 Dai Xinhai : 3A19
 Dai Xu : 3A19
 Dai Yanan : 1A7
 Daiguji Hirofumi : 1A3
 Dalacu Dan : 1A30
 Dalir Hamed : 1A12
 Dalvit Diego A. R. : 2A38
 Dammak Kamal : 3P2, 3P2
 Damodaran Anoop R. : 4P1
 Danièle Stéphane : 3A22
 Dao Thang Duy : 3A3
 Daquin Priscillia : 3P2
 Dardano Principia : 4A29
 Darvill Daniel : 2A19, 2A28
 Daskalaki Christina : 4A14
 Dauliat Romain : 4A23
 David Christin : 2A35
 Davis Timothy : 1A3
 Dawson M. D. : 4A30
 de Abajo Javier Garcia : 2A3
 de Almeida Jorge V. : 2P2
 De Angelis Costantino : 1A19, 3A5, 4A24
 De Angelis Francesco : 2A19, 2A19, 2A28, 2A28
 De filippo Giovanni : 4A29
 De Giorgi Milena : 3A1
 De Girolamo Del Mauro Anna : 4A29
 de Groot C. H. : 4A18
 de Groot C.H. (Kees) : 3A23
 De Liberato Simone : 4A10
 De Luca Anna Chiara : 4A35
 De Luca Antonio : 4A12, 4A24, 4A36
 De Luca Federico : 2A1
 de Lustrac André : 2A10
 De Marco Maria Letizia : 1A22
 de Paz Maria Blanco : 1A11
 De Proft Anabel : 3A17
 De Stefano Luca : 4A29
 De Vittorio Massimo : 4A28, 4A29
 De Wolf Ingrid : 3A5
 Decher Gero : 1A5
 Deck-Leger Zoe-Lise : 2A35
 Decleva Piero : 2A24
 Deinert Jan-Christoph : 4A17
 Del Valle Nuria : 3A35
 Delaunay Jean-Jacques : 1A3
 Delerue Christophe : 2A21
 Deletang Thibault : 4P1
 Deligeorgis George : 4A14
 Della Valle Giuseppe : 2A9, 4A31
 Demichel Olivier : 4A18
 Demir Hilmi Volkan : 3A2
 Demmerle Frederic : 3A12
 Deng Chih-Zong : 1A3
 Deng Fu : 2A18
 Deng Longjiang : 3A7, 3P1, 4A19
 Denkova Denitza : 4A9
 Derkowska Beata : 3A22
 Dermeche Nassim : 2P2, 3P1
 Deschauer Niels : 1A9
 Desprez Sylvain : 3A14
 DeVault Clayton : 2A22, 3A39
 Devaux Thibaut : 1A8
 Devescovi Chiara : 1A11
 Dezert Romain : 1A22, 1A24, 2A21
 Dhama Rakech : 4A25
 Di Falco Andrea : 2A34
 Di Meo Valentina : 4A29
 Diana Rosita : 4A29
 Diehl Sebastian : 3A4
 Dietrich Kay : 1A24
 Dimaki Maria : 1P2
 Ding Desheng : 1A9
 Ding Fei : 4A32
 Ding Jie : 4A9
 Ding Kun : 1P1, 2A36, 3A7, 3A11, 3A18, 4A34
 Ding Wen Jun : 1A10
 Ding Wendu : 2A31
 Ding Xumin : 4A5, 4A5
 Dinh Thi-Thuy-Duong : 3A21
 Dionne Jennifer A. : 3A36
 Dirdal Christopher A. : 3A28, 4A32
 Dixon Katelyn : 3A34
 Djafari Rouhani Bahram : 4A30
 Djafari-Rouhani Bahram : 1A17, 1A27, 4P1
 Dmitruk Igor : 1P2
 Do Hue T. B. : 1A10
 Dolati Noushin : 3P1
 Domagalska Martyna : 4A28
 Dombi Peter : 2A20
 Dombi Péter : 1A18
 Dominguez Owen : 3A36
 Dominici Lorenzo : 3A1
 Donato Maria Grazia : 4A35
 Donegan John F. : 1A15
 Dong Tianyu : 3A11
 Dongare Pratiksha D. : 4A31
 Donnelly Fearghal C. : 1A4
 Dontabhaktuni Jayasri : 4A12, 4P1, 4P1
 Doporto José : 3P2
 Doppler Jorg : 2A33
 Dorche Ali E. : 3A37
 Doronin Iliya : 4A16
 Dorozhko Elena : 3A29
 Dose Wesley M. : 4A21
 Douillard Ludovic : 2A13
 Doyle Dennis : 2P2
 Drachev Vladimir P. : 2A24
 Drechsler Martin : 3A30
 Drisko Glenna : 1A22
 Droste Marvin : 1A9
 Droulias Sotiris : 4A22
 Drozdovskii Andrei V. : 1A16
 Drozd Piotr A. : 4P1
 Du Fan : 2A22

Du Kang : 3A11, 3A19
 Du Shengwang : 3A4, 4A11
 Duan Xueke : 4A18
 Dub Maksym : 1A29
 Dubcek Tena : 3A10
 Duda Tatyana A. : 4A34
 Dudek Michal : 1P1, 4P1
 Dudek Michał : 1A24
 Duguet Etienne : 1A22, 2A21
 Dujardin Gérald : 1P1
 Duo Lamberto : 4A18
 Duong Ngoc My Hanh : 4A12
 Dupré Cécilia : 1A22
 Duran Valdeiglesias Elena : 3A21
 Dutt Avik : 3A10
 Duvigneau Régis : 2A32
 Duò Lamberto : 3A5
 Dvorak Petr : 2P2, 4A26
 Dzhagan Volodymyr : 4A28, 4A34
 D’Orazio Antonella : 4P1
 Eckmann Felix : 1P1
 Economou Eleftherios : 3A27, 4A14, 4A22
 Economou Eleftherios N. : 4A10
 Edes Zoltan : 4A26
 Ehrler Jonathan : 3A16
 El Shamy Eslam : 3P2
 Elancheliyan Rajam : 1A22, 1A24, 4A8
 Eleftheriades George V. : 3A34
 Elezzabi Abdulhakem Y. : 2A39
 Elhadj S. : 1A25
 Elsayy Mahmoud M. R. : 2A32
 Emani Naresh Kumar : 2P1, 2P1, 3A8
 En Naciri Aotmane : 2A12
 Engheta Nader : 1A25, 1A28, 3A31, 3A39
 Epshteyn Albert : 3A33
 Epstein Eliav D. : 1A7
 Ergoktas M. Said : 4A14
 Erni Daniel : 1A19
 Eroglu Abdullah : 1P2
 Escorcía Carranza Ivonne : 2P2
 Esmann Martin : 1A17, 4A30
 Espinoza Shirly : 1A18
 Esposito Emanuela : 4A29
 Estakhri Nasim Mohammadi : 2A34
 Estakhri Nooshin M. : 2A34
 Euser Tijmen G. : 4A21
 Eustache Clement : 1P1
 Evans Dean R. : 1A18
 Eyi R. : 3P2
 Faccio Daniele : 3P2, 3A35, 3A39
 Fages Frederic : 4A4
 Fan Alvin : 2A16
 Fan Qingbin : 3A12
 Fan Ren-Hao : 1A10
 Fan Shanhui : 3A10
 Fan Xu-dong : 2A14
 Fan Zheng : 2A11
 Fang Chen : 3A7
 Fang Xiaoguo : 4A11, 4A11
 Fang Xinyuan : 4A20
 Farhi Asaf : 1A5
 Farle Michael : 3A16
 Fasolino Tommaso : 4A29
 Fathnan Ashif Aminulloh : 3A6
 Fedotov Vassili A. : 1A18, 3A1
 Fedyanin Andrey : 1A18, 2A37
 Fedyanin Andrey A. : 3A27, 4A9
 Fehrembach Anne Laure : 2A21
 Feigenbaum Eyal : 1A25
 Feilhauer Juraj : 2A33
 Feng Jing : 1A20
 Feng Lantian : 2A7
 Feng Rui : 2A10
 Fergestad Halvor : 3A9
 Fernandez-Corbaton Ivan : 2A16
 Fernandez-Dominguez Antonio I. : 1A29
 Fernández-Arias Mónica : 4A26
 Ferrera Marcello : 3A39
 Ferrera Marzia : 2A9, 4A8, 4A17
 Ferreyra Pablo : 4A33
 Fery Andreas : 1A23, 3A28, 4A24
 Fesenko Volodymyr : 2A15, 4A5
 fichtner Andreas : 3A10
 Figat Aleksandra Maria : 2P1
 filipiak Maciej : 1A29, 4P1
 filipkowski Adam : 1A5
 finazzi Marco : 3A5, 4A18
 finizio Simone : 4A31
 fink Mathias : 2A16
 finot Eric : 2P1
 fiore Andrea : 3A5, 4A3
 Fiorini-Debuisschert Céline : 2A13
 fip Tassilo : 3A21
 fix Baptiste : 3A35
 Flach Sergej : 1A3
 Flamant Quentin : 1A24
 Flannery Jeremy : 1A26
 Fleischer Monika : 1A13, 3A1
 Fleischmann Claudia : 3A5
 Fleury Guillaume : 1A14, 4A8
 Fleury Romain : 2A23, 4A30
 Flores Enrique : 4A27
 Flores Fernando Gonzalez-Posada : 4A28
 Florescu Marian : 2A20, 4A3
 Floris F. : 4A3
 Fontana Jake : 2P2, 4A33
 Forcherio Gregory T : 3A33
 Forestiere Carlo : 4A29
 Forti Stiven : 4A17
 Foti Antonino : 4A35
 Fournel Franck : 1A12
 Fowler Clayton : 2A27
 Fowler Daivid : 1A22
 Fox Maayan : 1A7
 Fraile Rodríguez Arantxa : 4A28
 Francone A. : 3A14

Franczyk Marcin : 1A5
 Frank Thomas : 3A1
 Fratalocchi Andrea : 1P2, 1P2, 3A23
 Freimuth Frank : 2A15
 Frej Antoni : 4A31, 4A31
 Friman Alexander : 4A26
 Fritzsche Wolfgang : 4A29
 Frolov Aleksandr Yu. : 4A9
 Frosz Michael H. : 4A21
 Fröch Johannes E. : 4A12
 Fuhrmann Bodo : 4P1
 Fujii Kensuke : 3P1
 Fujii Minoru : 1P1
 Fujita Kentaro : 1A3
 Fujiwara Hideki : 3A2
 Fumeaux Christophe : 3A8
 Fung Kin Hung : 4A1
 Gacoin Thierry : 1A4, 3A29
 Gajc Marcin : 1A18
 Galante Angelo : 1A19
 Galdi Vincenzo : 4A29
 Galiffi Emanuele : 1A24, 3A11, 4A21
 Galimov Aidar : 1A18
 Galinski Henning : 2P2
 Gallas Bruno : 2A21
 Galli Matteo : 3A39
 Gallo Katia : 3A9
 Ganapathy Boopalan : 4A19
 Gandolfi Marco : 4A24
 Gao Feng : 3A9
 Gao Jie : 2A12, 3P1
 Gao Kun : 3A11
 Gao Penglin : 3A37
 Gao Song : 4A15
 Gao Weilu : 3P2
 Gao Wenlong : 1A11, 1A20, 1A20
 Gao Xiaomei : 3A9
 Gao Yuan : 3A2
 Garcia de Abajo F. Javier : 2A7
 Garcia Esteban : 1A14
 Garcia P. D. : 3A14
 Garcia Parajo Maria F. : 2A21
 Garcia-Elcano Inaki : 2P1
 Garcia-Etxarri Aitzol : 1A11
 Garcia-Raffi Lluís M. : 1A8
 Garoli Denis : 2A19, 2A28, 4A36
 Gaspari Matteo : 3A23
 Gassenq Alban : 4A2
 Gawinkowski Sylwester : 2A28, 2P2
 Gawlik Andrzej : 3A5
 Ge Dandan : 2A13
 Geldmeier Jeff : 4A33
 Geldmeier Jeffrey A. : 2P2
 Genevet Patrice : 2A15, 2A32, 3P1
 Gensch Michael : 4A17
 Gerace Dario : 3A39
 Gerard Davy : 2A2
 Getman Fedor : 1P2, 1P2
 Ghaemidzicheh Hamed : 2P1, 3A15
 Gianfrate Antonio : 3A1
 Giannino G. : 4A3
 Giesz Valérian : 1A17
 Gigli Carlo : 1A19, 2A17, 3A5
 Gil Javier : 4A26
 Giloan Mircea : 4A25
 Giner Lambert : 1A30
 Ginzburg Pavel : 1P1, 2A19
 Giordano Maria Caterina : 4A31
 Giovannini Giorgia : 2A19, 2A28
 Girard Christian : 1A12
 Giraudet Louis : 1P2
 Gjessing Jo : 3A28, 4A32
 Gkantzounis G. : 2A20
 Gleeson Matthew : 1A29
 Gliserin Alexander : 2P1
 Gnoatto Lorenzo : 4A28
 Goddard Lynford : 4A27
 Godiksen Rasmus H. : 1P2
 Goessling Johannes. W. : 3A22
 Goffi Fatima Zohra : 2A16
 Gogoi Ankur : 3P1
 Golovynskyi Sergii : 1P2
 Gomez Rivas Jaime : 1P2
 Gomez-Carbonell Carmen : 1A17
 Gomonay O. : 2A15
 Goncalves P. A. D. : 3A29
 Gong Qihuang : 2A9, 4A18
 Gong Yongkang : 1A20
 Gonzalez-Tudela Alejandro : 2P1
 Gorbatshevich Alexander : 4A26
 Gorji Setatira : 3P2
 Gorodestki Yuri : 1A16
 Gorodetski Yuri : 1A7
 Gould Stephen : 2A18
 Govorov Alexander : 1A4, 1A26, 3A33
 Govorov Alexander O. : 1A13, 1A23
 Goya Shinya : 3A3, 3A23
 Grafe Markus : 2A33
 Graham Thomas : 1A9
 Grajower Meir : 1A6, 2A22
 Granchi Nicoletta : 3A5, 4A3
 Grande M. : 4A3
 Grande Marco : 4A29
 Grange Rachel : 2A30
 Grant James : 2P2
 Grant James P. : 3A35
 Gratus Johnathan : 2A35
 Gratus Jonathan : 2A25
 Gray Stephen K. : 3A33
 Grey Clare P. : 4A21
 Greybush Nicholas J. : 2P2, 4A33
 Gric Tatjana : 3P2
 Grinblat Gustavo : 3A34, 3A34
 Grishin Sergey : 2P2, 2P2
 Groby Jean-Philippe : 1A8, 1A8
 Grosjean Thierry : 1P1, 1A13

Gruszecki Pawel : 2A24
 Gu Min : 3A40
 Gu Ying : 2A9, 4A18
 Gualdrón-Reyes Andrés F. : 3P2
 Guan Chunsheng : 4A5
 Gubarev Fedor : 3A28
 Gubbin Christopher R. : 4A10
 Gucciardi Pietro Giuseppe : 4A35
 Guenneau Sébastien : 2A1
 Guerber Sylvain : 1A22
 Guerin Sarah : 1A29
 Guerrero Albert : 4A28
 Gugole Marika : 3A17
 Guilhabert B. : 4A30
 Guldin Stefan : 1A14
 Gun'ko Yurii K : 1A4
 Gunther Matthias : 1P1
 Guo Jinying : 4A20
 Guo L. Jay : 3A26
 Guo Qinghua : 1A20, 3A20
 Guo Shangkun : 4A9
 Guo Yangyu : 1A8
 Guo Zhiwei : 3A7
 Gupta Shilpi : 3P1
 Gupta Vaibhav : 4A24
 Gurdal Emre : 3A1
 Gurioli Massimo : 3A5, 4A3
 Gurung Sudip : 3A32
 Gurvitz Egor A. : 4A28
 Gutowski Łukasz : 2P2
 Gutt Robert : 4A25
 Gérard Davy : 3A13
 Günzler Antonio : 4A26
 Ha Céline : 3P2
 Ha Kyoungho : 4A4
 Hafermann Martin : 4A32
 Hafez Hassan A. : 4A17
 Haffner Christian : 3A34
 Haffouz Sofiane : 1A30
 Haidar Riad : 4A28
 Hainey Jr. Mel F. : 1A3
 Haji Ebrahim Mehdi : 3P2
 Hajian Hodjat : 3A29
 Halas Naomi J. : 4A31
 Hamarneh Ramez : 4A5
 Hamazaki Ryusuke : 2A5
 Hammerschmidt Martin : 2A17
 Han Joo Hun : 2A3
 Han Narae : 2A7
 Han Sangjun : 2A4
 Han Seunghoon : 2A7
 Hananel Uri : 3A13
 Hannema Gwenaël : 1A17
 Hao Zhenzhong : 3A9
 Haraguchi Masanobu : 1A3
 Haran Gilad : 1A29
 Harouri Abdelmounaim : 1A17, 1A20, 4A30
 Harutyunyan Hayk : 2A31
 Hasanpour Tadi Saeed : 3P2
 Hashimoto Rei : 2A1
 Hawak Dan : 3A23
 Hayati Raad Shiva : 1P1
 Haïdar Riad : 3P2, 4A28
 He Huilin : 2A2, 4A11
 He Runqiu : 3A19
 Healy Noel : 4A34
 Heinrich Matthias : 2A33
 Heller Evan : 2P2
 Hendry Euan : 3A39
 Hennard Luc : 1A18, 1P2
 Hernandez Martinez Pedro Ludwig : 3A2
 Hernandez Romain : 4A18
 Hernandez Sergi : 1P2
 Hernández-García Carlos : 3A40
 Herranz Gervasi : 2A34
 Herrero Ramon : 2A8, 2A14, 2A37
 Hess Ortwin : 1A15, 2A13
 Hewak Daniel W. : 2P2
 Hibbins Alastair : 1A9, 2A10, 2P1, 2P2
 Hidaka Mio : 3P1
 Hierro Adrian : 1A18
 Higashino Makoto : 3A3, 3A23
 Hillebrands Burkard : 1A16
 Hillenbrand Rainer : 4A34
 Hilmi Volkan Demir : 1A4
 Hinamoto Tatsuki : 1P1
 Hisaeda Yoshio : 2P2
 Hjorvarsson Bjorgvin : 4A12
 Hmima Abdelhamid : 1P2, 2P1
 Ho Ya-Lun : 1A3
 Ho Ying -Lung Daniel : 2P2
 Hocini Abdesselam : 2P2, 3P1
 Hoffman Anthony : 3A36
 Holland Glenn : 3A34
 Holleitner Alexander : 2A12
 Holman Hoi-Ying : 3A34
 Hong Ic-Pyo : 2P1
 Hong Min Ji : 4A23
 Hong Qilin : 4A17
 Hopmann Eric : 2A39
 Horak Michal : 3A28, 3A29, 4A9, 4A26
 Horn Alexander : 4A17
 Horneber Anke : 3A1
 Horsley Simon : 2P1
 Hotta Ikuto : 1A10
 Hou Bo : 4A1
 Houdellier Florent : 3A25
 Hrton Martin : 3P2, 4A9, 4A26, 4A34
 Hrtoň Martin : 4A15
 Hsu Jin-Chen : 3P1
 Hu Changyu : 3A39
 Hu Hui : 3A9
 Hu Juejun : 3A38
 Hu Mengying : 3A7
 Hu Xiao : 4A13
 Hu Yabin : 1A8

Huang Chen-Bin : 1A7
 Huang Chung-Che : 2P2
 Huang Guoliang : 1A27, 1A27, 1A27
 Huang Jian-An : 2A28
 Huang Jianan : 2A28
 Huang Lingling : 4A11
 Huang Lujun : 2A18
 Huang Yu : 1A27
 Huang Zhen-Ting : 4A13
 Huang Zhifeng : 3A1
 Hubarevich Aliaksandr : 2A19, 2A28
 Huber Patrick : 3A5
 Huebner Uwe : 2A4
 Huffaker Diana L. : 1A20
 Huh Ji-Hyeok : 3A11
 Hui Yuen-Yung : 2P1
 Huidobro Paloma : 1A24
 Huidobro Paloma A. : 1A20, 4A21
 Huidobro Paloma Arroyo : 1A11
 Humbert Mélodie : 1A12
 Huq Tahiyat : 3A39
 Hussain Abdulrahman : 3P2
 Hussainova Irina : 1A28
 Huttenhofer Ludwig : 1P1
 Hwang Inoh : 4A4
 Hwang Inyong : 3A12
 Hyaran Zeki : 2A14
 Höfling Sven : 2A3
 Hübner Uwe : 1A24
 Iadanza S. : 4A3
 Iadanza Simone : 3A39, 4A3
 Iarossi Marzia : 2A19, 2A28
 Ibarias Martin : 3P2
 Ibrir Miloud : 3P2
 Ichiji Naoki : 4P1
 Iff Oliver : 2A3
 Ignatyeva Daria : 2A1
 Ikamas Kestutis : 4A32
 Ikeda Naoki : 2A1
 Ikeda Tatsuhiko N. : 2A6
 Ikezawa Satoshi : 3A14
 Ilyas Saddia : 2P2
 Im Eunji : 4A16
 Imamura Riku : 3A1
 Indiveri Ivano : 3A23
 Inoue Takashi : 1A7
 Intaravanne Yuttana : 4A20
 Intonti Francesca : 3A5, 4A3
 Ioannidis Thanos : 3P2
 Ippoliti Rodolfo : 2A19
 Iqbal Naeem : 3A6
 Iranzo David Alcaraz : 4A17
 Isella Giovanni : 4A18
 Ishaaya Amiel Avraham : 1A28
 Ishihara Hajime : 1A7, 4P1
 Ishii Satoshi : 3A3, 3A3, 3A23
 Issa Ali : 3A29
 Ivanov Misha : 2A24
 Ivanov Roman : 1A28
 Ivanov Viktor : 4A26
 Iwami Kentaro : 3A14
 J. Beck Fiona : 3P1
 Jacassi Andrea : 3A39
 Jacob Jesly : 3P1
 Jaeck Julien : 3P2
 Jakob Lukas : 1A13
 Jamadi Omar : 1A20
 Jamier Raphael : 4A23
 Jandieri Vakhtang : 1A19
 Jang Jaehyuck : 3A17, 4A20
 Jang Min Seok : 2A4
 Jankiewicz Bartłomiej J. : 2P1, 2P2
 Jaramillo Juliana : 3A14
 Jariwala Deep : 2A34
 Jarrah Mona : 2A30
 Jatteau Tom : 1A22
 Jauho A.-P. : 3A29
 Jeffrey Stefanie S. : 3A36
 Jenkins Ronald P. : 2A32
 Jenne Jurgen : 1P1
 Jensen Geir Uri : 3A28
 Jeon Seokwoo : 4A4
 Jeong Byung Gil : 4A4
 Jeong Myung Yung : 2P1
 Jeong Young-Gyun : 3A33
 Jheng Shih-Da : 1P1
 Ji Chengang : 3A26
 Ji Wenye : 4A32
 Jia Danchen : 2A22
 Jiang Haitao : 3A7
 Jiang Shun Liu : 3A6
 Jiang Shunliu : 3A6
 Jiang Xi : 3A7
 Jiang Xue : 2A36, 4A13
 Jiang Yunlu : 2A3
 Jiao Yuqing : 4A12
 Jimba Yoji : 1A3
 Jin Jeongwan : 1A30
 Jin Jicheng : 3A10
 Jin Weiliang : 2A32, 3A38
 Jin Yabin : 1A17, 4A30
 Jin Yuqi : 1A27
 Jing Yun : 4A13
 Joglekar Yogesh : 2A23
 Johns Ben : 2A9
 Johns Paul : 2P2, 4A33
 Johnson M. A. : 1A25
 Johnson Noble : 3A37
 Johnson Steven G. : 3A38, 3A38
 Jonsson Magnus : 3A16
 Jonsson Magnus P. : 3A17
 Joseph Joby : 1A10
 Jradi Safi : 3A29, 3A33
 Juliano-Martins Renato : 4A18
 Jung Pawel : 2A33
 Jurow Matthew : 4A23

Jusza Anna : 3P1
 Kachorovskii Valentin : 4P1
 Kaczmarek Malgosia : 3A1
 Kafesaki Maria : 3A27, 4A10, 4A14, 4A22
 Kahlouche Ahmed : 2P2, 3P1
 Kaiser Florian : 4A23
 Kaivola Matti : 1P2
 Kajikawa Kotaro : 3A3, 3P1
 Kakuno Tsutomu : 2A1
 Kalfagiannis Nikolaos : 3A25
 Kalinikos Boris A. : 1A16
 Kall Mikael : 2A17
 Kamali Khosro Zangeneh : 2A18
 Kameche Farid : 2A13
 Kamenetskii Eugene : 2A24, 3A30
 Kanamori Yoshiaki : 3A3
 Kanega Minoru : 2A6
 Kaneko Kei : 2A1
 Kang Ming : 4A11
 Kang Tongtong : 3A7, 4A19
 Kanoun Olfa : 3A28
 Kapaev Vladimir : 4A26
 Kapaklis Vassilios : 4A12
 Kapoor Rohit : 3P2
 Kapush Olga : 4A28
 Karakhanyan Vage : 1P1
 Karimi Habil Mojtaba : 3P2
 Karki Dolendra : 2A1
 Karmakar Subhajit : 3P1
 Karvounis Artemios : 2A30
 Kasaya Takeshi : 1A3
 Kashif Muhammad Fayyaz : 4A29
 Kaspar Corinna : 2A13
 Kastl Christoph : 3A16
 Kasztelanik Rafal : 1A5
 Kato Keizo : 3A3
 Kats Mikhail : 4A32
 Katsantonis Ioannis : 4A22
 Kaur Manpreet : 3A3
 Kaur Sarabjeet : 3A33
 Kawabata Kohei : 2A5
 Kawano Akihiro : 3A14
 Kawazu Takuya : 1A3
 Kazlou Artsiom : 3P2
 Kedze Kam Eucharist : 4P1
 Kehagias N. : 3A14
 Kejik Lukas : 2P2, 3A28, 4A26
 Kenanakis George : 4A10, 4A14
 Kenis Karine : 3A5
 Kenney Mitchell G. : 2P2
 Kopic Peter : 4A34
 Kerič Peter : 3P2, 4A15
 Kern Dieter P. : 3A1
 Keshavarz Hedayati Mehdi : 4A21
 Khabarov Kirill : 4A26
 Khadir Samira : 3P1
 Khajavikhan Mercedeh : 2A33
 Khalfi Roumaissa : 1P2
 Khalid Muhammad : 2A25
 Khan Irfan : 3A36
 Khan Pritam : 1A22, 1A29, 4A22
 Kharitonova Olga : 1P2
 Khatib Omar : 2A27
 Khavasi Amin : 1P2, 2P1, 2P2, 3P2
 Kherani Nazir : 3A34
 Kherani Nazir P. : 3A34
 Khodzitsky Mikhail : 2P1
 Kholpin Dmitry : 2A2
 Khomeriki Ramaz : 1A19
 Khort Pavlo : 4A28
 Khousakoun Eric : 3A14
 Khrabustovskyi Andrii : 2A16
 Kik Pieter G. : 3A5
 Kildebro Lars : 3A23
 Kildishev Alexander V. : 1A14
 Kim Daeik : 3A12
 Kim Dong Eon : 2P1
 Kim Doo-In : 2P1
 Kim Heungsoo : 1A30
 Kim Hyeon Ho : 4A16
 Kim Hyochul : 2A7
 Kim Inki : 4A1, 4A11, 4A24
 Kim Jaekwan : 2A7
 Kim Jeongmo : 1A4
 Kim Jeongyong : 3A12
 Kim Ji Su : 4A4
 Kim Jisoo : 2A4
 Kim Jong-Wook : 1A4
 Kim Jongwook : 3A29
 Kim Joosung : 2A3
 Kim Jun Hyun : 2P1
 Kim Jung-Woo : 3P1
 Kim Kwangjin : 4A10
 Kim Min Soo : 4A4
 Kim Minkyung : 1A11
 Kim Nakhyun : 2A3
 Kim Sang-Hoon : 3P1
 Kim Sejeong : 4A12
 Kim Seungchul : 2P1
 Kim Seyoon : 2A4
 Kim Shin Young : 4A15
 Kim Shinho : 2A4
 Kim Soo Jin : 4A4
 Kim Sun Il : 4A4
 Kim Sung-Soo : 1P1
 Kim Yongsung : 2A7
 Kim Yoon Young : 4A15
 Kim Zee Hwan : 2A2
 Kinsler Paul : 2A25, 2A25, 2A35
 Kipp Jonathan : 2A15
 Kirchner Robert : 3A28
 Kirschner Matthew S. : 2A31
 Kishen Saurabh : 2P1, 2P1, 2P1, 3A8
 Kisielewski Jan : 2A24
 Kislov Denis : 4A28
 Kitajim Hiroki : 2A6

Kityk Andriy V. : 3A5
Kivshar Yuri : 2A4
Klar Thomas A : 2A9
Klauck Friederike U. J. : 2A33
Kley Ernst-Bernhard : 1A24
Klimko Grigorii : 1A18
Klimov Vasily V. : 2A8, 2A26
Klitis C. : 4A30
Klos Andrzej : 3P1
Klug Bernhard : 4A36
Klymov Oleksii : 1A18
Knap Wojciech : 1A29, 4P1
Ko Woong : 2A7
Kocabas Coskun : 4A14
Koch Alexander : 4A32
Kochanowska Olga : 1A28
Kociak Mathieu : 2A2, 3A25
Kodera Toshiro : 4A23
Koenderink Femius : 2A20
Koenig Tobias A. F. : 1A23, 4A24
Koenig Tobias A.F. : 1A10
Koksal Koray : 3A30
Kolaciński Cezary : 4A32
Kolkowski Radoslaw : 4A29
Kolymagin Danila : 4A2
Konecna Andrea : 2A7, 4A9
Konecny Martin : 4A26
Kong Kiho : 2A3
Kong Ling Qi : 3A6
Kong Lingqi : 3A6
Kong Xiang Kun : 3A6
Kong Xiangkun : 3A6
Kongsuwan Nuttawut : 1A15
Kono Junichiro : 3P2
Konotop Vladimir : 2A8
Konstandin Simon : 1P1
Kontoleta Evgenia : 2A20
Kooksal Koray : 3A40
Koppens Frank : 4A17
Kort-Kamp Wilton J. M. : 2A38
Koschny Thomas : 3A27
Kosik Miriam : 3A37
Kostina Natalia : 2A19
Kostina Nataliia : 1P2
Koulouklidis Anastasios : 4A14
Koutserimpas Theodoros T. : 2A23
Kovalev Sergey : 4A17
Kowerdziej Rafal : 1P1, 4P1
Kowerdziej Rafał : 1A24
Kozhaev Mikhail : 2A1
Krahne Roman : 4A12, 4A24, 4A36
Krajczewski Jan : 4A8
Krajewska Aleksandra : 2P1
Krapek Vlastimil : 3A28, 4A9, 4A26, 4A34
Krawczyk Maciej : 1P2, 2A24, 3P2
Krishnamoorthy Harish N. S. : 4A12
Krishnan Sivarama : 4P1
Krivitsky Leonid : 3A34
Krokhin Arkadii : 1A27, 3P2
Krol Mateusz : 3A40
Kroychuk Maria : 1A18
Krushynska Anastasiia O. : 1A17
Kuai Yan : 3A11
Kubo Atsushi : 4P1
Kubo Wakana : 3A2
Kudrya Vladislav : 1P2
Kudyshev Zhaxylyk : 1A14, 3A39
Kues M. : 4A30
Kuhl Ulrich : 2A33
Kula Przemyslaw : 3A40
Kulkarni Manas : 2A23
Kumar Anshuman : 1P1
Kumar Govind : 3P1
Kuo Shan-Jen : 2P1
Kura Naoto : 2A5
Kuroda Takashi : 2A1
Kurt Hamza : 2A14
Kutsevol Nataliya : 4A28
Kuttruff Joel : 4A36
Kuwana Yasuhiro : 3P1
Kuzmin Dmitry : 1P2
Kuznetsov Arseniy : 3A34
Kuznetsov Arseniy I. : 4A12
Kuznetsova Svetlana : 1A8
Kuznetsova Vera A. : 1A4
Kvapil Michal : 2P2, 4A26
Kwak B.Leo : 3A11
Kwon Namseop : 2A7
Kwon O-Pil : 3A33
Kyaw Aung Ko Ko : 4A7
Kyriakou Eydokia : 4A14
Käll Mickael : 3P1
L. Muskens Otto : 4A18
Labartaa Amílcar : 4A28
Labiad Abdelhadi : 2P2
Labonté Laurent : 3A21, 4A23
Laborieux Axel : 3A29
Laferrière Patrick : 1A30
Lafforgue Christian : 3A21
Lafitte Maeva : 1A24
Lagoudakis Pavlos : 3A1
Lagoudakis Pavlos G. : 3A40
Lagutchev Alexei S. : 1A14
Lahlil Khalid : 1A4
Lai Yun : 4A11
Laible Florian : 1A13
Lakhtakia Akhlesh : 2A25
Lalanne Philippe : 2A17, 3A4, 3A5
Lamaitre Aristide : 1A20
Lamaze Florian : 1P2, 2P1, 3A13
Lan Shoufeng : 4A33
Lanco Loic : 4A30
Landi Gabriel : 3A15
Langevin Denis : 3P2
Lanoy Maxime : 1A27
Lanteri Stéphane : 2A32

Lanzillotti-Kimura Daniel : 3A21
 Lanzillotti-Kimura Norberto Daniel : 1A17, 4A30
 Lapointe Jean : 1A30
 Larquet Eric : 3A29
 Larrey Vincent : 1A12
 Larrieu Guilhem : 1A12, 2A12
 Latyshev Alexander V. : 4A34
 Lau Kai Ming : 3A4, 4A11
 Lauri Alberto : 1P1
 Lavrinenko Andrei V. : 3A23, 4A22
 Lawrence Mark : 3A36
 Le Gratiet Luc : 1A20, 4A30
 Le Moal Eric : 1P1
 Le Roux Xavier : 1A22, 3A21
 Leach Gary : 3A34
 Leahu Grigore : 1A5
 Lecestre Aurélie : 1A12, 2A12
 Lee Byounggho : 1A6
 Lee Duhyun : 4A4
 Lee Eunsung : 2A3
 Lee Gun-Yeal : 1A6
 Lee Howard : 3A32
 Lee In Ho : 4P1
 lee jaesoong : 3A20
 Lee Jaewon : 3A11
 Lee Jeong Yub : 2A7, 2A7
 Lee Jisan : 4A4
 Lee Jongwon : 3A12
 Lee Joong Seok : 4A15
 Lee S. : 3P2
 Lee Sam Hyeon : 1A8
 Lee Sang-Shin : 4A15
 Lee Sangyun : 4A1
 Lee Seung-Heon : 3A33
 Lee Seungwoo : 3A11, 4A4, 4A10, 4A16
 Lee Sung Won : 4A4
 Lee Woo-Bin : 4A15
 Lee Woorim : 4A15
 Lee Yang-Chun : 1A3
 Lee Yeon Ui : 4A4
 Lefebvre Isabelle : 2A21
 Leff Asher C : 3A33
 Lefier Yannick : 1P1
 Lei Dangyuan : 1A6, 2A1, 2A8
 Leite Marina S. : 3A36
 Leitner Nikolaus S. : 2P2
 Lemaire Vincent : 1A5
 Lemaitre Aristide : 4A30
 Lemaître Aristide : 1A17
 Lembessis V. E. : 3A37, 3A40
 Lembessis Vasileios E. : 3P1
 Lembessis Vassilis E. : 3A30
 Lemoult Fabrice : 1A27, 2A16
 lempicka Karolina : 3A40
 Lenz Kilian : 3A16
 Leo Giuseppe : 1A19, 3A5
 Lepeshov Sergei : 4A32
 Lepore Maria : 4A29
 Lera-Leri Roger : 1P2
 Lerario Giovanni : 3A1
 Leray Aymeric : 1A4, 2P1
 Lermusiaux Laurent : 1A22, 2A21
 Lerner Gavriel : 2A24
 Lerosey Geoffroy : 1A27
 Leroy Valentin : 1A27
 Lertvachirapaiboon Chutiparn : 3A3
 Levy Miguel : 2A1
 Lewandowski Wiktor : 4P1, 4P1, 4A24
 Lezaic Marjana : 2A15
 Lezec Henri J. : 3A12, 3A34
 Lheurette Eric : 4P1
 Li Bassi Andrea : 2A9
 Li Bing : 1A8
 Li Hailiang : 3A19
 Li Hang : 2A27
 Li Hao-Xiang : 2A14
 Li Heng : 4A13
 Li Jensen : 1A8, 2A8, 2A16, 3A4, 4A11, 4A11
 Li Jiaqi : 4A9
 Li Jinxiang : 4A30
 Li Juntao : 4A6
 Li Ke : 2A2
 Li Liangsheng : 3P1
 Li Pei : 3A11
 Li Qiang : 1A3, 4A19
 Li Qitong : 3A5
 Li Tao : 4A1
 Li Xiangping : 3A18
 Li Xiaopeng : 1A27
 Li Xiaowei : 4A11
 Li Xiaoyan : 2A2
 Li Xiaoying : 4A18
 Li Yang : 2A22, 2P2
 Li Yao : 4A16
 Li Yi : 1P1, 3A34, 3A34, 3A40, 4A10
 Li Yong : 4A13
 Li Yongqian : 4A18
 Li Yue : 3P1
 Li Yunhui : 3A7
 Li Yuyu : 4A33
 Li Zhancheng : 1A20
 Li Zhe : 1A29
 Li Zheng : 1A8, 1A27
 Li Zhichao : 2A9
 Li Zhipeng : 2A9
 Liang Bin : 2A14, 4A13
 Liang G. H. : 3A19
 Liang Haowen : 4A6
 Liang Hong : 3A4
 Liberal Inigo : 1A25
 Liberale Carlo : 3A33
 Lieu Simon : 2A33
 Ligmajer filip : 2P2, 4A9, 4A34
 Ligmajer Filip : 3P2, 4P1, 4A15
 Lim C. W. : 1A9
 Lim Minwoo : 4A1

Limonov Mikhail : 1P1
 Lin Ching-Che : 2P1
 Lin Qian : 3A10
 Lin Tzy-Rong : 3P1
 Lin Wan-Yu : 3A11
 Lin Xiao-Min : 2A31
 Lin Zin : 3A38
 Lindner Jürgen : 3A16
 Lioubtchenko Dmitri : 2P1
 Lioubtchenko Dmitry : 1A28
 Lipovka Anna : 3A28
 Lissauskas Alvydas : 4A32
 Liska Jiri : 2P2
 Liska Jiri : 3P2
 Lisiansky Alexander : 4A16
 Liszewska Malwina : 2P1, 2P2
 Littlefield Alexander : 4A27
 Liu Changxu : 1P1
 Liu H. : 3A19
 Liu Hui : 3A7
 Liu Jianxun : 2A2, 4A11, 4A11
 Liu Jing-jing : 4A13
 Liu Liu : 2A11
 Liu Ning : 1A22, 1A29, 4A22
 Liu Rongjuan : 3A7
 Liu Shijie : 3A9
 Liu Tian : 1P1
 Liu Tongjun : 4A30
 Liu Wenwei : 1A20
 Liu Xiao-Jun : 3A6
 Liu Xiaojun : 1A9
 Liu Yan Jun : 4A11
 Liu Yanjun : 2A2, 4A11
 Liu Yi : 4A23
 Liu Yongmin : 2A27
 Liu Yongquan : 1A8
 Liu Yueyang : 2A22
 Liu Yun-Kai : 4P1
 Liu Yupeng : 2A6
 Liu Yuzhou : 2A33
 Lizunova Anna : 4A26
 Liégeois Vincent : 1P2
 Liška Jiří : 4P1
 Lodari Mario : 4A18
 Lodde Matteo : 4A3
 Lodewijks Kristof : 3A17
 Loh Joel Y. Y. : 3A34
 Lohmüller Theobal : 3A22
 Loncar Marki : 2P2
 Lopez Garcia Martin : 3A22
 Lopez-Aymerich Elena : 1P2
 Lopez-Garcia Martin : 3A33
 Lorenzo Salvatore : 3A15, 3A15
 Lorenzon Monica : 4A23
 Lotti Francesco : 2A3
 Lova Paola : 4A26
 Low Tony : 1A30, 2A4
 Lozovik Yurii : 4A16
 Lu Junchi : 3A36
 Lu Li : 4A12
 Lu Ling : 3A7
 Lu Lizhen : 3A11
 Lu Tien-Chang : 4A13
 Lu Wei : 3A19, 4A9
 Lu Xuyang : 3A35
 Lu Yi-Hsien : 4A23
 Lu Yu-Jung : 3A6
 Lucklum Frieder : 1A8
 Luk Ting Shan : 3A39
 Lumeau Julien : 2A21
 Luo Dan : 2A2, 4A11
 Luo Jie : 4A11
 Luo Qiang : 3A9
 Lux Fabian : 2A15
 Lv Ruizhen : 4A16
 Lyons Ashley : 3A35
 Lyras A. : 3A37
 Lyras Andreas : 3P1
 Lyubin Evgeny V. : 3A27
 Lähderanta Erkki : 1A16
 M. Gaskell Jeffrey : 4A18
 Ma Guancong : 2A36
 Ma Renmin : 2A8, 3A18
 Ma Xikui : 3A11
 Ma Xuezhi : 4A33
 Ma Yixuan : 2A18
 Ma Yun : 4A18
 Ma Yungui : 2A11, 3A6
 Maccaferri Nicolo : 2A19, 2A28, 4A36
 MacDonald Kevin F. : 4A30
 Mach-Battle Rosa : 3A35, 4A28
 Macias Demetrio : 1A28
 Mackay Tom : 2A25
 Mackowski Sebastian : 4A2, 4A28, 4A28
 MacNab finlay : 3A34
 Maes Bjorn : 3A14
 Maes Clément : 4A28
 Maglione Maria Grazia : 4A29
 Magnozzi Michele : 1A23, 2A9, 4A8, 4A17
 Mahfoud Zackaria : 1A10
 Mahmoodi Maryam : 4A22
 Mahmoud Ahmed : 1A15
 Maier Stefan : 3A34, 3A34, 3A39
 Maier Stefan A. : 1P1, 1P1, 2A4, 4A10, 4A20, 4A23
 Maier Stefan Alexander : 4P1
 Mailybaev Alexei A. : 2A33
 Maiti Rishi : 1A12
 Majorel Clément : 1A12
 Majumdar Arka : 1A25
 Majérus Bruno : 1A18
 Makarenko Maksim : 1P2
 Makarov Sergey : 3P1
 Makhsiyani Mathilde : 4A28
 Maksim Makarenko : 1P2
 Mallet Nicolas : 1A12
 Malléjac Matthieu : 1A8

Malo Dana : 4A26
 Malof Jordan : 2A27
 Malpuech Guillaume : 3A1
 Malureanu Radu : 1P1, 4A22
 Maly Sergei : 4A26
 Malzard Simon : 2A23
 Manasreh O. : 3P2
 Mancarella Cristina : 2A9
 Mancini Andrea : 4A10
 Mangini Maria : 4A35
 Manhaes Mosso Marbey : 2P2
 Manjarrez Bryan : 4A27
 Manjavacas Alejandro : 2A26
 Mano Takaaki : 1A3
 Manushyna Daria : 1A9
 Many Véronique : 2A21
 Manyakin Ilya : 4A21
 Mao Peng : 1P1
 Mao Wenbo : 3A9
 Marago Onofrio Maria : 4A35
 Marguet Sylvie : 2A13
 Marigo Jean-Jacques : 2A1
 Marini Andrea : 2A4, 3P2
 Marino Giuseppe : 1A19, 3A5
 Marko Gilad : 4A16
 Markovich Gil : 3A13
 Marks Daniel : 2A27
 Marris-Morini Delphine : 1A22, 3A21
 Martin Jérôme : 2A2, 3A13
 Martin Nicolas : 1A13
 Martin Olivier J. F. : 4A5
 Martin-Sanchez Javier : 2A3, 3A24
 Martinez Castellano Eduardo : 1A18
 Martinez-Llinas Jade : 2A17
 Martinez-Pastor Juan P. : 3P2
 Martinez-Tomas Carmen : 1A18
 Martínez Castells Marc : 4A28
 Marzban Mahmood R. : 2P2
 Masell Jan : 1A7
 Masenelli Bruno : 1A12, 2A21, 3A22
 Masenelli-Varlot K. : 2A21
 Maserati Lorenzo : 3A16
 Mathew Sanro : 1A4
 Mathur Avi : 4P1, 4P1
 Matsuda Osamu : 1A3, 1A8
 Matsukata Taeko : 2A7
 Matsuo Mamoru : 2A6
 Mattei Giovanni : 1A5
 Mattera Lorenzo : 2A9
 Matuszewski Michal : 3A40
 Matveev Oleg : 3P1
 Maurel Agnes : 2A1
 Maurel Agnès : 2A16
 Maurer Thomas : 1P2, 2P1, 3A13
 May S. : 4A30
 Mayer Martin : 4A24
 Mazeas Florent : 4A23
 Maziewski Andrzej : 2A24
 Mazur Eric : 2A22, 2P2
 Mazur Rafal : 3A40
 Mazzanti Andrea : 4A31
 McCall Martin : 2A16
 McCall Martin W : 2A25
 McPhillimy J. : 4A30
 Meckenstock Ralf : 3A16
 Medina Judith : 2A14
 Megahd Heba : 4A26
 Mehmood Muhammad Q. : 4A24
 Mei Hongyan : 4A32
 Mei Ting : 3A11, 3A19
 Meixner Alfred J. : 1A13, 3A1
 Mello Olivia : 2A22
 Mello Olivia L. : 2P2
 Melloni filippo : 3A16
 Mendez-Sanchez Rafael A. : 4A27
 Meng Chao : 4A32
 Meng Fanqi : 4A36
 Mengali Sandro : 3A23
 Merkel Aurélien : 1A8
 Merte Maximilian : 2A15
 Merzouk Hamid : 1P2, 2P1
 Meuret Sophie : 3A25
 Mezil Sylvain : 1A8
 Michon Jérôme : 3A38
 Miele Ermanno : 4A21
 Mignuzzi Sandro : 3A39
 Mikelson Jelena : 4P1
 Mikkelsen Maiken H. : 3A31
 Milekhin Alexander : 4A34
 Milekhin Ilya A. : 4A34
 Milgrom Benjamin : 4P1
 Milicevic Marijana : 1A20
 Miller Owen : 3A38
 Min Changjun : 4A11
 Minamimoto Hiro : 2P1
 Minarini Carla : 4A29
 Miniaci Marco : 1A17
 Minin Igor : 4A27
 Minin Igor V : 3P2
 Minin Oleg : 4A27
 Minin Oleg V : 3P2
 Minkov Momchil : 3A10
 Minn Khant : 3A32
 Miranda Bruno : 4A29
 Mirek Rafal : 3A40
 Miroshnichenko Andrey E : 2A18
 Miscioscia Riccardo : 4A29
 Mitra Joy : 2A9
 Mivelle Mathieu : 2A21
 Miyazaki Hideki T. : 1A3
 Miyazaki Hiroshi : 1A3
 Mnasri Karim : 2A16
 Mnaymneh Khaled : 1A30
 Moccia Massimo : 4A29
 Mocella Vito : 4A35
 Mochizuki Masahito : 1A7

Moczala-Dusanowska Magdalena : 2A3
 Mohammed Miami Mohammed : 3P2
 Mohammedi Abdelkader : 3P2
 Moiseyev Nimrod : 2A33
 Moitra Parikshit : 4A12
 Mokrousov Yuriy : 2A15
 Molesky Sean : 2A32, 3A38
 Molinaro Céline : 2A13
 Mondain-Monval Olivier : 1A22, 4A8
 Monneret Serge : 3P1
 Montambaux Gilles : 1A20
 Montazeri Arthur : 3A34
 Montes Bajo Miguel : 1A18
 Montesinos Miguel : 3A21
 Monval Olivier Mondain : 1A24
 Mora-Sero Ivan : 3P2
 Moradi Golamreza : 1P2
 Morandotti Roberto : 3A33
 Morassi Martina : 4A30
 Morawiak Przemyslaw : 3A40
 Moreno Mauricio : 1P2
 Morgan Katrina A. : 2P2
 Morin Henri : 1A30
 Moritake Yuto : 4A13
 Moroshkin Petr : 3P2
 Morozova Maria : 3P1
 Mortensen N. A. : 3A29
 Morvillo Pasquale : 4A29
 Mosali Raviteja R. : 4A12
 Moses Jebin J. : 3A36
 Moshchalkov Victor V. : 4A9
 Mousavi Mansoureh Z. : 2A28
 Movsesyan Artur : 2A21
 Muhammad F. : 1A9
 Mukherjee Abhishek : 4P1, 4P1
 Muljarov Egor A. : 3A22
 Muller Moos : 1P2
 Muller Philipp : 4A29
 Munday Jeremy : 1A25
 Munoz Elias : 1A18
 Munoz-Sanjosé Vicente : 1A18
 Mura G. : 4A3
 Murai Shunsuke : 3A2, 3A3, 3A23
 Muraki Yusuke : 3A5
 Murakoshi Kei : 2A7, 2P1
 Murastov Gennadiy : 3A28
 Musa Umar : 1P2
 Muskens Otto L. : 2A12, 2A18, 3A23
 Musslimani Ziad H. : 2A33
 Mélinon Patrice : 2A21
 N. V. Jayanth : 4A19
 Na Byunghoon : 2A7
 Naciri Jawad : 2P2, 4A33
 Nagao Tadaaki : 3A3, 3A3, 3A23
 Nagashima Keigo : 3A1
 Nahvi Ehsan : 1A25
 Nakanishi Toshihiro : 4A13
 Nakata Yosuke : 4A13
 Naldoni Alberto : 1A4
 Naoi Yoshiki : 1A3
 Narang Prineha : 2A20
 Narushima Tetsuya : 2A2
 Naryanan Remya : 2P1
 Nasir M. Farooq : 2P2
 Nasir Mazhar : 2A21
 Nasir Mazhar E. : 2A3
 Nassiri Alireza : 1A30
 Nath Partha P. : 3P1
 Navaratna Nikhil : 1P1
 Navarro-Urrios Daniel : 1P2
 Nenna Giuseppe : 4A29
 Neogi Arup : 1A27
 Neshev Dragomir N. : 2A18
 Neufeld Ofer : 2A24
 Neumann Oara : 4A31
 Nguyen H. T. : 1A25
 Nguyen Huy : 1A27
 Nicholls Luke H. : 2A3, 2A21
 Niedziolka-Jönsson Joanna : 4A2, 4A28
 Nielsen Michael : 3A34
 Niemi Tapio : 1A14
 Nikitin Andrey A. : 1A16
 Nikitov Sergey : 2P2
 Nishijima Yoshiaki : 4A13
 Niu Jiebin : 4A6
 Niv Avi : 3A39, 4A17
 Noda Susumu : 3A3
 Noguchi Yuki : 1A9
 Nomura Masahiro : 1A8
 Noor Ahsan : 4P1
 Nordlander Peter : 4A31
 Northeast David B : 1A30
 Notomi Masaya : 1A2, 1A11, 4A13
 Nowack Thomas S. : 2P2
 Nowaczynski Rafal : 1A18, 4P1
 Nunez-Lasus Zoilo : 1P2
 Nuytten Thomas : 3A5
 Nyga Piotr : 1A14
 O'Brian P. : 4A3
 O'Carroll Deirdre M. : 1A21
 O'Faolain L. : 4A3, 4A3
 O'Faolain Liam : 3A39
 O'Gorman Nicholass Alan : 1P2
 O'Keeffe Patrick : 2A9
 Oberhammer Joachim : 1A28, 2P1
 Ocana-Pujol Jose L. : 2P2
 Ochiai Tetsuyuki : 1A3
 Oda Reiko : 2A12
 Odent Jeremy : 3A14
 Odintsov Sergey : 2P2
 Ogrin Feodor : 1A16, 2P2
 Oh Joo Hwan : 4A4
 Oh JooHwan : 1A9
 Oh Sang Hyun : 4P1
 Oh Sang Soon : 1A11, 1A20, 3A4
 Ohashi Michio : 3A6

Okamoto Hiromi : 3A2
 Okamoto Takayuki : 2A7, 3P1
 Okimura Kunio : 4A13
 Olbrich Markus : 4A17
 Olifierczuk Marek : 1A24, 4P1
 Olk Andreas E. : 3A6
 Olsson Oliver : 3A17
 Olthaus Jan : 3A37
 Omatsu Takashige : 2A6
 Omeis Fatima : 2A28
 Oniki Yusuke : 3A5
 Ordonez Andres Felipe : 2A24
 Ortiz Omar : 1A17, 3A21
 Ortolani Michele : 2A1
 Orzechowski Kamil : 4P1
 Osato Hirotaka : 1A3
 Oser Dorian : 3A21
 Otaki Katsura : 3A6
 Otome Yuta : 3A14
 Otsuji Taiichi : 4A14
 Otsuka Paul H. : 1A8
 Otten Matthew : 3A33
 Ou Jun-Yu : 4A30
 Oudjertli Salah : 3A28, 3P2
 Oue Daigo : 1P1, 4A34
 Oulton Rupert : 3A34
 Ozawa Tomoki : 1A20
 Ozbay Ekmel : 3A29
 Ozerov Igor : 2A21, 4A4
 Pace Simona : 4A17
 Pacheco Peña Victor : 1A28, 4A34
 Pacheco Peña Victor Manuel : 3A39
 Paddubrouskaya Hanna : 4A9
 Padilla Willie : 2A27
 Page John : 1A27
 Pagliano Francesco : 3A5
 Paiella Roberto : 4A33
 Paillard Vincent : 1A12
 Pala Ragip A. : 1A6
 Paladini Alessandra : 2A9
 Palange Elia : 1A19
 Palei Milan : 3A36
 Palekar Shubhangi Mahendra : 3P1
 Palhinha Castillo Miguel A. : 3A33
 Palma Gioacchino Massimo : 3A15, 3A15
 Pan Lei : 4A16
 Pan Meiyang : 4A19
 Pan Yi-Ming : 4P1
 Pan Ying : 3A38
 Pande Divya : 2A27
 Pandolfi Giuseppe : 4A29
 Panella Gloria : 2A19
 Paniagua-Dominguez Ramon : 4A12
 Pankin Pavel S. : 3A3
 Panov Vladimir I. : 4A9
 Papa Zsuzsanna : 1A18
 Papadakis Georgia Theano : 4A12
 Papaioannou Evangelos : 4A12
 Papas Dimitrios : 4A30
 Park Chul-Soon : 4A15
 Park Haedong : 1A11
 Park Hongkyu : 4A1
 Park Hyeonsoo : 2A7
 Park Ikmo : 4P1
 Park Jinjoo : 2A3
 Park Jung Hun : 2A3
 Park Junghyun : 4A4
 Park Namkyoo : 2A5, 4A14
 Park Sang-Min : 2A2
 Park Seongjin : 3A12
 Park Yeonsang : 2A7
 Park Younghwan : 2A3
 Parka Janusz : 1P1, 1A24, 4P1
 Parnell William J. : 4A26
 Parto Midya : 2A33
 Pascarella Francesco : 4A29
 Pashaei Adl Hamid : 3P2
 Passarella Bianca : 3A16
 Pastier Florian : 1A17
 Paszke Piotr : 1A18, 4P1
 Patra Aniket : 4A12
 Pauly Matthias : 1A5, 2A12
 Pavlov Alexander A. : 4A28
 Pawlak Dorota : 3P1
 Pawlak Dorota A. : 3P1, 3P1
 Pawlak Dorota Anna : 1A18, 4P1
 Payeur Stéphane : 3A33
 Pecorario Stefano : 3A16
 Pellegrino Daniele : 3A5
 Pendry J. B. : 4A34
 Pendry John : 1P1, 3A11, 4A21
 Pendry John B. : 1A24
 Pendry John Brian : 2A8
 Peng Bo : 3P1
 Peng Chao : 3A10
 Peng Ruwen : 1A10
 Peng Zheng : 3A7, 3P1
 Pennec Yan : 1A17, 4P1, 4A30
 Pensa Evangelina : 1P1
 Penzo Erika : 4A23, 4A35
 Perez-Galacho Diego : 3A21
 Perez-Sanchez Juan B. : 1A23
 Perinot Andrea : 3A16
 Pernet Nicolas : 1A20
 Pernice Wolfram : 2A13
 Perrakis George : 4A10
 Pertsch Thomas : 1A24
 Petek Hrvoje : 1A7
 Petit Marlène : 4A18
 Petronijevic Emilija : 1A5
 Petrov Ilia : 3A28, 3A29
 Petruzzella Maurangelo : 4A3
 Petti Daniela : 4A31
 Petti Lucia : 4A29, 4A29
 Pfeiffer Loren N. : 3A1
 Pflug Theo : 4A17

Pham Kim : 2A1, 2A16
 Philipp Dennis : 1P1
 Pianelli Alessandro : 1P1, 1A24, 4P1
 Piao Xianji : 2A5, 4A14
 Piatkowski Dawid : 4A28
 Picardi Michela Florinda : 4A31
 Piccardo Marco : 3A24
 Piccoli Riccardo : 3A33
 Piecek Wiktor : 3A40
 Piechulla Peter M. : 4P1
 Pierrat Romain : 1A14
 Pike Trinity C. : 3A36
 Pin Christophe : 3A2
 Pinho Pedro V. : 2P2
 Pinton Nadia : 4A24
 Pintos Jean-François : 3P2
 Piotrowski Piotr : 1A18, 4P1
 Piqué Alberto : 1A30
 Piramidowicz Ryszard : 3P1
 Pisanello Ferruccio : 4A28, 4A29
 Pisanello Marco : 4A29
 Pisano filippo : 4A28, 4A29
 Pisarenko Anastasia : 4A2
 Pittner Angelina : 4A29
 Piętka Barbara : 3A40
 Plain Jerome : 2A21
 Plain Jérôme : 3A13
 Plidschun Malte : 2A4
 Plum Eric : 4A30
 Plum Michael : 2A16
 Poblet Martin : 3A34
 Pohlit Merlin : 4A12
 Politi Alberto : 4A10
 Pommier Delphine : 1P1
 Ponsinet Virginie : 1A14, 1A22, 1A24, 2A21, 4A8
 Poole Philip J : 1A30
 Portaccio Marianna : 4A29
 Postnikov Pavel : 3A29
 Pothuraju Renuka Devi : 4A12
 Potzger Kay : 3A16
 Pou Juan : 4A26
 Pou Pablo : 4A26
 Pouget Emilie : 2A12
 Poulikakos Lisa V. : 3A36
 Powell Alex W. : 2A10
 Powell David A. : 3A6
 Pradhan Akash Kumar : 2P1
 Prajapati Ashish : 4A31
 Prajapati Ashsih : 4A16
 Prat-Camps Jordi : 3A35
 Prato Mirko : 3A16
 Prencipe Alessandro : 3A9
 Primot Jérôme : 4A28
 Priya Priya : 1A17, 4A30
 Probst Patrick T. : 4A24
 Proctor Matt : 1A11
 Prodan Emil : 1A27
 Proust Julien : 1P2, 2P1, 2A21, 3A13, 3A33
 Prystawko Pawel : 1A29, 4P1
 Przewłoka Aleksandra : 2P1
 Pucci Andrea : 4A26
 Pukhov Alexander : 4A16
 Purcell-Milton finn : 1A4
 Purdy Andrew : 3A33
 Puri Aniket S. : 3A36
 Purkayastha Archak : 2A23
 Pylypovskyi Oleksandr V. : 1A16
 Pysz Dariusz : 1A5
 Qi Jingbo : 2A6
 Qian Zhiyuan : 2A9
 Qiao Tong : 3A7
 Qin Jun : 3A7, 4A19
 Qin Mingpu : 3A10
 Qiu Chengwei : 4A13
 Qiu Min : 1A3, 1A6, 4A19
 Qu Junle : 1P2
 Qualtieri Antonio : 4A29
 Quan Cong : 2A10
 Quan Jia-Qi : 4P1
 Quevedo-Teruel Oscar : 4A5
 Raabe Jorg : 4A31
 Raagavi B. : 3P1
 Raczkiewicz Marcin : 3P1
 Raczkiewicz2 Marcin : 3P1
 Rafailov Edik : 3P2
 Rahaman Mahfujur : 4A34
 Rahm Marco : 3A21
 Rahmani Mohsen : 2A18
 Rahmani Motahareh : 3P2
 Rakhlin Maxim : 1A18
 Rakovich Yury P. : 2A39
 Ramaswamy Kannan : 1P1, 4P1, 4P1
 Ramò Lorenzo : 4A17
 Rana Ahsan S. : 4A24
 Rao Neeraj : 3P1
 Raquez Jean-Marie : 3A14
 Rarity John G. : 2P2
 Rastelli Armando : 2A3
 Ratni Badreddine : 2A10, 2A10, 2A15, 4A5
 Ravaine Serge : 1A22, 2A21
 Ravets Sylvain : 1A20
 Ravishankar Ajith Padyana : 2A2
 Ravnik Miha : 4A12
 Ray N. J. : 1A25
 Raza Soren : 1P1, 2A39
 Razdolski Ilya : 3P2
 Raziman T. V. : 1P2, 2P1, 4A30
 Razzari Luca : 3A33
 Rea Ilaria : 4A29
 Real Bastian : 1A20
 Rebarz Mateusz : 1A18
 Reben Manuela : 3P2, 3P2
 Rechcinska Katarzyna : 3A40
 Redka Dmitrii : 4A28
 Refaely-Abramson Sivan : 3A16
 Regev Oren : 1A24

Regmi Raju : 2A21
Reid Barry : 1A14
Reif Johanna : 3A28
Reimer Michael : 3A34
Reimhult Erik : 2P2
Reineke Bernhard : 4A11
Reinhard Bjoern : 1A26
Reisner Mattis : 4A23
Reiter Doris E. : 3A37
Rejaei Behzad : 2P1
Remesh Vikas : 4A9
Ren Haoran : 2A4, 3A40, 4A20
Ren Juanjuan : 4A18
Ren Simiao : 2A27
Ren Xifeng : 2A7
Rendina Ivo : 4A35
Reniers Sander F. G. : 4A12
Rensberg Jura : 4A32
Reshetnyak Victor : 1A18
Reuter Cornelia : 4A29
Rho Junsuk : 1A11, 2A2, 2A18, 3A17, 4A1, 4A11, 4A20, 4A24
Ricciardi Rosa : 4A29
Rice Daragh : 4A22
Richetti Philippe : 1A22, 2A21
Richter Felix : 2A30
Ricouvier Joshua : 1A14
Riedo Elisa : 4A31
Rigneault Herve : 2A21
Riley Joseph : 4A34
Rinaldi Christian : 4A31
Rippa Massimo : 4A29, 4A29
Riveiro Antonio : 4A26
Rivera N. : 3A29
Rizza Carlo : 1A19
Roach Lucien : 2A21
Roberto Li Voti Roberto : 1A5
Roberts Ann : 1A3
Roberts Nathan : 4A21
Roccati Federico : 3A15, 3A15
Rocco Davide : 1A19, 3A5, 4A24
Rockstuhl Carsten : 1A24, 2A16, 4P1
Rodriguez A. : 3A21
Rodriguez Alejandro : 2A32
Rodriguez Alejandro W. : 3A38
Rodriguez Alvarez Javier : 4A28
Rodriguez Anne : 1A17, 4A30
Rodriguez Raul : 3A29
Rodriguez Raul D. : 3A28
Rodríguez Fortuño Francisco : 4A31
Rodríguez Daniel : 4A26
Rodyakina Ekaterina E. : 4A34
Roeder Ryan : 3A36
Roh Sookyoung : 4A1
Romanenko Dmitrii : 3P1
Romanenko Dmitriy : 2P2
Romano Silvia : 4A35
Romano-Rodriguez Albert : 1P2
Romero-Garcia Vicent : 1A8, 1A8
Ronning Carsten : 4A32
Rontgen Malte : 1A19
Rosendo Lopez Maria : 1A9
Rosendo-López María : 2A14
Roskos Hartmut G. : 3P2, 4A36
Rottenberg Xavier : 3A17
Rotter Stefan : 2A33
Rovenska Katarana : 4A34
Rovenska Katarina : 3P2
Rovenská Katarína : 4P1
Rovere Andrea : 3A33
Rowan-Robinson Richard : 4A12
Roy Chowdhury Dibakar : 3P1
Roy Philippe : 4A23
Rozniecka Ewa : 4A28
Rumpf R. : 3P2
Rumyantsev Sergey : 1A29
Ruostekosk Janne : 3A37
Russell Philip St.J. : 4A21
Rybin Mikhail : 1P1
Röhrich Ruslan : 2A20
S. Brindhu Malani : 1P1
S. Simitha : 3P1
Saba Matthias : 4A26
Sadasivan Jayesh : 4P1
Sadecka Katarzyna : 3P1
Saeidi Hooman : 3A35
Saenz Juan J. : 1A15
Safari Mahdi : 3A34
Safou Abdelkader : 3A28
Safronov Kirill : 2A37
Sagne Isabelle : 1A19
Sagnes Isabel : 1A20
Sagnes Isabelle : 1A17, 4A30
Saha Soham : 3A39
Sai Pavlo : 1A29
Sain Basudeb : 4A11
Saito Shinji : 2A1
Sakai Kyosuke : 2A6
Sakhare Pratiksha A. : 4P1
Sakoda Kazuaki : 2A1
Sakowicz Maciej : 1A29
Sakthivel M. S. : 3P1
Salas-Montiel Rafael : 4A18
Salim Flora : 3A40
Sallet Vincent : 2A21
Salman Jad : 4A32
Salmeron Miquel : 4A23
Salmon Arthur : 3A35
Salut Roland : 1A13
Samanta Kartik : 2A15
Sambles John Roy : 2A10
Sambles Roy : 1A9, 2P2
Sammon Michael : 1A30
Samusev Kirill : 1P1
San-Jose Pablo : 3A37
Sanchez-Dehesa Jose : 3A37

Sanchez-Gil Jose A. : 1A15
 Sandomenico Annamaria : 4A29
 Sannomiya Takumi : 2A7
 Santiago Eva Y : 3A33
 Sanvitto Daniele : 3A1
 Sapienza Riccardo : 2A20, 3A39, 4A3
 Saplacan Gavril : 4A25
 Sarkar Swagato : 1A10
 Sartori Carlos A. F. : 2P2
 Sasaki Keiji : 2A6, 3A2
 Sato Daiki : 2P1
 Sato Masahiro : 2A6
 Sauvan Christophe : 2A36
 Savaresi Matteo : 2A3
 Savona Vincenzo : 4A30
 Scaramuzzi Giuseppe : 4A31
 Scarpettini Alberto F. : 4A23
 Schadenko Sergey : 3A28
 Schaller Richard D. : 2A31, 3A21
 Schatz George C. : 2A31
 Scheibner Colin : 1A27
 Schindler Frank : 3A10
 Schletz Daniel : 3A28
 Schmidt Markus A. : 2A4
 Schmidt-Kaler Ferdinand : 3A30
 Schmiegelow Christian Tomas : 3A30
 Schneider Christian : 2A3
 Schneider Philipp-Immanuel : 2A17
 Schneidewind Henrik : 2A4
 Schomerus Henning : 2A23, 2A36, 3A4
 Schrunner Philip P. J. : 3A37
 Schuck Carsten : 3A37
 Schuermans Silvère : 3A13
 Schulz Sebastian A. : 2A34, 3A39
 Schumacher Cedric : 4A26
 Schumer Alexander : 2A33
 Schwartz David E. : 3A37
 Schwartzberg Adam : 4A23
 Sekar Sribharani : 1A5
 Semenov Alexander A. : 1A16
 Semisalova Anna : 3A16
 Semnani Behrooz : 1A26
 Sen Mrinal : 2P1
 Senellart Pascale : 1A17, 4A30
 Sengupta Kaushik : 3A35
 Sentker Kathrin : 3A5
 Serebryannikov Andriy : 1P2, 3P2
 Serebryannikov Andriy E. : 3A29
 Serebryannikov Andriy E. : 1P2
 Serga Alexander A. : 1A16
 Sergeev Aleksandr : 4A27
 Sergeeva Daria : 2P2
 Sergeeva Ksenia : 4A27
 Serna Rosalia : 3A21
 Serra-Garcia Marc : 3A10
 Serras Florenci : 1P2
 Seviour Rebecca : 2A35
 Sha Wei E. I. : 2P1
 Sha Yixin : 3A7
 Shah Wiqar Hussain : 3P1, 4A32
 Shah Yash D. : 2P2
 Shah Yash Diptesh : 3A35, 3A39
 Shahbazyan Tigran V : 4A23
 Shalaev Vladimir : 2A29, 3A39
 Shalaev Vladimir M. : 1A14
 Shalev Gil : 4A16, 4A25, 4A31
 Shalin Alexander : 2A19
 Shalin Alexander S. : 1P2, 4A28
 Shamkhi Hadi K. : 4A28
 Shan Lingxiao : 4A18
 Shang Xiao : 4A6
 Shao Linda : 2A10
 Shao Wei : 3A40
 Shaqqura Nadim : 3A1
 Sharma Naresh : 3P1
 Sharp Ian D. : 1P1
 Shavrov Vladimir : 1P2
 Shayegannia Moein : 3A34
 Shcherbakov Maxim R. : 4A9
 Shcherbinin Vitalii : 4A5
 Shearer Tom : 4A26
 Sheremet Evgenia : 3A29
 Sheremet Evgeniya : 3A28
 Shi Lei : 3A7
 Shi Lina : 4A6
 Shi Yue : 4A11
 Shigetou Akitsu : 1A3
 Shilkin Daniil A. : 3A27
 Shin Dong Chul : 2A3
 Shin Jaikwang : 2A3
 Shin YeJeong : 1A9
 Shinbo Kazunari : 3A3
 Shiraki Hirofumi : 4P1
 Shirmanesh Ghazaleh : 2A22
 Shirmanesh Ghazaleh Kafaie : 1A6
 Shishkin Ivan : 1P1
 Shkondin Evgeniy : 3A23
 Shokri Babak : 3P2
 Shorokhov Alexander : 1A18
 Shubin Nikolay : 4A26
 Shubina Tatiana : 1A18
 Sibilia Concita : 1A5
 Siebadji Tchuimeni Daniel : 4A25
 Siefke Thomas : 1A24
 Sielezin Karol : 1A24, 4P1
 Siena Julya : 2A13
 Sigmund Ole : 3A38
 Sikola Tomas : 2P2, 3A29, 3P2, 4A9, 4A26, 4A34
 Sikola Tomas J. : 3A28
 Sileo Leonardo : 4A28, 4A29
 Silien Christophe : 1A22, 4A22
 Silien christophe : 1A29
 Silov Andrei : 3A5
 Silva Sinhara R. : 2A38
 Silvani Raffaele : 4A31
 Silveirinha Mario : 4A21

Simeoni Mirko : 3A23
Simon Thomas : 2A2
Simpkins Blake : 3A33
Simpson Robert E. : 4A12
Simões Gamboa Ana Luisa : 4A2
Sinelnik Artem : 1P1
Singh Leeju : 1A7
Singh leeu : 1A16
Sinkevich Eugene : 4A26
Sisler Jared : 2A22
Sivun Dmitry : 2A9
Slama Sebastian : 1A22
Slivina Evgeniia : 4P1
Slowik Karolina : 3A37
Slowikowski Mateusz : 4P1
Smirnov Serguei : 2P1
Smirnova Daria A. : 2A18
Smirnova Olga : 2A24
Smith David R. : 2A27
Smith Terry : 1A30
Soci Cesare : 4A12
Sofiel Mathew : 1A27
Sokhoyan Ruzan : 1A6, 2A22
Soliman Amr Mohamed : 4P1
Soljadic M. : 3A29
Soljadic Marin : 3A10
Solnyshkov Dmitry : 3A1
Song Jung-Hwan : 3A5
Song Junyeob : 3A12, 3A34
Song Qinghua : 3P1
Soppera Olivier : 2A13
Sorel M. : 4A30
Sorger Volker J. : 1A12
Sotomayor-Torres C. M. : 3A14
Soukoulis Costas : 3A27, 4A14, 4A22
Spalding Richard : 2A20, 4A3
Spolenak Ralph : 2P2
Sprafke Alexander N. : 4P1
Staliunas Kestutis : 2A8, 2A14, 2A37
Stanfield Leanne D. : 2P2
Staples Bethany : 1A9
Star Dmitry : 3A34
Steglich Martin : 1A24
Steiner Anja Maria : 4A24
Stelmaszczyk Kamil : 1A29
Sternklar Shmuel : 1A7
Sternklar Shumuel : 1A16
Stoger-Pollach Michael : 4A9
Stognij Alexander : 3P2
Stoja Endri : 1P1
Stomeo Tiziana : 4A29
Strain M. J. : 4A30
Strait Jared H. : 3A34
Stranik Ondrej : 4A29
Strusch Tanja : 3A16
Strzezysz Olga : 4P1
Stupakiewicz Andrzej : 1A12, 3P2, 4A31, 4A31
Suarez Isaac : 3P2
Suarez Miguel : 1A13
Suemitsu Masahiro : 3A3
Suffit Stephan : 1A19
Sugimoto Yoshimasa : 1A3, 2A1
Sukham Johneph : 4A22
Sukhorukov Andrey A. : 2A8
Sulowska Karolina : 4A2, 4A28, 4A28
Sun Hongbo : 1A20, 1A20
Sun Kai : 3A23
Sun Mao-Guo : 3A3
Sun Mingyu : 1P1
Sun Yong : 3A7
Sung Jangwoon : 1A6
Surma Barbara : 3P1
Surma Hancza : 1A18
Surma Hancza Barbara : 4P1
Sutherland Duncan : 4A35
Suzuki Tatsuro : 3A2
Svendsen Winnie E. : 1P2
Sviridova Elizaveta : 3A29
Swan Anna : 4A33
Swathi Sugumar : 3P1
Sygletou Maria : 2A9, 2A9
Symonds Clementine : 3A1, 4A2, 4P1
Sypniewska Malgorzata : 3A22
Szameit Alexander : 2A33
Szczytko Jacek : 3A40
Szelag Bertrand : 1A22
Szlachetko Kamil : 1A18, 3P1, 4P1
Szola Maria : 4P1
Słowikowski Mateusz : 1A29
Tabeling Patrick : 1A14
Taboada-Gutiérrez Javier : 2A3
Tacchi Silvia : 4A31
Tagliabue Giulia : 3A25
Taha Mohamed K. H. : 2A21
Tahara Azusa : 3A14
Tai Yi-Hsin : 1A3
Takagi Hidetsugu : 3A6
Takahara Junichi : 1A10, 3A14
Takashima Yuusuke : 1A3
Takayama Osamu : 3A23, 4A22
Talentikit Djahida : 2P1
Talentikite-Touati Djahida : 1P2
Taliercio Thierry : 4A28
Tallarico Domenico : 1A17
Tam Wing Yim : 2A16, 3A4, 4A11
Tamagnone Michele : 3A24
Tamayo-Arriola Julen : 1A18
Tanabe Takasumi : 3A1
Tanaka Katsuhisa : 3A2, 3A3, 3A23
Tanaka Takuo : 1A6
Tanaka Yasuhiro : 1A7
Tang Haoning : 2A22
Tang Hong : 2A27
Tang Weiyuan : 2A36
Tanzilli Sebastien : 3A21, 4A23
Tapar Jinal : 2P1, 3A8

Tapar Jinal Kiran : 2P1
 Tapaszto Levente : 1A18
 Tarnowska Katarzyna : 2P2
 Tasolamprou Anna : 3A27, 4A14
 Tasolamprou Anna C. : 4A10
 Tassin Philippe : 2A17
 Tassini Paolo : 4A29
 Tavakol Mohammad Reza : 3P2
 Tavakoli Nasim : 2A20
 Tavassoli Seyed Hassan : 4A22
 Taverne Mike P. C. : 2P2
 Tedeschi Davide : 2A3
 Tedesco A. : 4A3
 Teselko Petro : 1P2
 Theron Christophe : 3A22
 Thiele Matthias : 4A29
 ThiThanhTam Hoang : 3P1
 Thollar Zac : 2A7
 Thomas Anna Anupa : 3A16
 Thomas Vibin Ipe : 3P1
 Thompson Robert : 2A16
 Thomson Mark D. : 4A36
 Thrane Paul : 4A32
 Thureja Prachi : 2A22
 Thyagarajan Krishnan : 3A37
 Tian Jianguo : 1A20, 1A20
 Tian Jianjun : 4A7
 Tielrooij Klaas-Jan : 4A17, 4A17
 Tilmann Benjamin : 3A34
 Timofeev Ivan V. : 3A3
 Timpu Flavia : 2A30
 Timucin Dogan : 3A37
 Tishchenko Alexey : 2P2, 2A35
 Tisso Jayur : 3P1
 Titov Aleksey : 2P2
 Tizei Luiz : 3A25
 To Naoki : 4A13
 Toanen Vincent : 4A2, 4P1
 Toba Hidemitsu : 3A6
 Tofail Syed A. M. : 1A22, 1A29, 4A22
 Tognazzi Andrea : 4A24
 Tollerton Craig : 3A39
 Toma Mana : 3P1
 Tomchuk Anastasiia : 1P2
 Tomczyk Monika : 3P1
 Tomoda Motonobu : 1A3, 1A8
 Tongay Sefaattin : 2A3
 Tonielli Federico : 3A4
 Tonkaev Pavel : 3P1
 Torelli Piero : 2A9
 Toropov Alexey : 1A18
 Torrent Daniel : 1A9, 2A14, 3A37, 4A30
 Toschi Francesco : 2A9
 Toudert Johann : 3A21
 Tounsi Amel : 1P2
 Tourin Arnaud : 1A27
 Tournat Vincent : 1A8
 Tozawa Hiroya : 1A8
 Traviss Daniel : 4A18
 Treguer-Delapierre Mona : 1A22
 Trotta Rinaldo : 2A3
 Tréguer-Delapierre Mona : 2A21
 Tupikowska Martyna : 4P1
 Turchinovich Dmitry : 4A17
 Tuz Vladimir : 2A15, 4A5
 Tzortzakis Stelios : 4A10, 4A14
 Tünnermann Andreas : 1A24
 Ueda Masahito : 2A5, 2A5
 UI-Islam Qamar : 4A36
 Umatham Reiner : 1P1
 Umed Singh Dipti : 2P1
 Urade Yoshiro : 4A13
 Urbach Paul : 4A32
 Urbani Alessandro : 3A23
 Urbas Augustine : 2A22
 Ustinov Alexey B. : 1A16
 V. Grayli Sasan : 3A34
 Vaagen Thrane Paul C. : 3A28
 Vakarin Vladyslav : 3A21
 Valev Ventsislav K. : 3A2
 Valuckas Vytautas : 4A12
 Van Damme Bart : 1A17
 van de Groep Jorik : 3A5
 van der Heide Paul : 3A5
 Van Dorpe Pol : 3A17, 4A9
 van Hulst Niek F. : 1P2, 4A9
 van Otten Frank : 3A5
 Vancso Peter : 1A18
 Vandenbosch Guy A. E. : 4A9
 Vantaraki Christina : 4A12
 Vardeny Shai. R. : 2A38
 Varshney Ravendra Kumar : 3P1
 Vasco Juan Pablo : 4A30
 Vashistha Vishal : 1P2, 3P2
 Vasiliev Roman B. : 4A34
 Vazquez Besteiro Lucas : 1A13
 Vembris Aivars : 4P1
 Venkataram Prashanth S. : 2A32, 3A38
 Venkatesh Suresh : 3A35
 Verellen Niels : 4A9
 Vergniory Maia Garcia : 1A11
 Verlhac Clément : 4A28
 Verre Ruggero : 2A17, 3P1
 Vestri Ambra : 4A29
 Vezzoli Stefano : 3A39
 Vial Alexandre : 1A28
 Vidal François : 3A33
 Vijaya R. : 3P1
 Vilas-Iglesias Ana M. : 4A26
 Vincent Grégory : 4A28
 Vinogradov Alexey : 4A16
 Visser C. Peter : 2P1
 Visser Rober J. : 3A11
 Viswanath P. : 1P1
 Vitelli Vincenzo : 1A27
 Vivien Laurent : 1A22, 3A21

Vogler-Neuling Viola Valentina : 2A30
Volz Sebastian : 1A8
Voronov Andrey : 2A1
Vuckovic Jelena : 2A32
Vynck Kevin : 3A5
W. Sheel David : 4A18
Wackenhut Frank : 1A13
Waldschmidt Geoff : 1A30
Walker Ezekiel : 1A27
Wambold Ray : 4A32
Wan Chenghao : 4A32
Wan Chenglong : 2A20
Wan Dehui : 2P1
Wan Otten Frank : 4A3
Wang Baiming : 1P1
Wang Binbin : 4A18
Wang Feng : 2A31
Wang Heng : 3A11, 3A19
Wang Hongfei : 2A8
Wang Jianfang : 1A10
Wang Jiawei : 2A2, 4A11
Wang Mengjia : 1A13
Wang Mu : 1A10
Wang Qi : 3A6
Wang Qiang : 3A7
Wang Shaojun : 1P2
Wang Shixun : 4A7
Wang Shubo : 4A1
Wang Teng : 4A20
Wang Wan : 4A30
Wang Wei : 1A17, 4A30
Wang Wenhao : 1A4
Wang Wenhui : 1A11
Wang Xinke : 4A20
Wang Yansong : 3P1
Wang Yongtian : 4A11
Wang Yudong : 4A18
Wang Yuxiang : 2A10
Wang Zengbo : 4A25, 4A27
Wang Zhiming : 1A4, 1A23
Wang Zhiming M. : 1A13
Wang Zhiyu : 1A3
Wang Zhuochao : 4A5
Wang Zishuai : 2P1
Wani Waseem A. : 1P1
Wardley William P. : 3A22, 3A33
Watad Ibrahim : 1P2
Watanabe Kazuhiro : 1A3
Weber Sebastien : 3A25
Weber-Bargioni Alexander : 4A23
Weeber Jean-Claude : 4A18
Wegener Martin : 2A3
Wehrspohn Ralf B. : 4P1
Wei Qi : 3A6
Wei Qunshuo : 4A11
Wei Zhengxing : 2A6
Weigand Helena : 2A30
Weiner John : 1A22
Weiss Thomas : 3A22
Wen Jing : 1A6
Wen Long : 3A7
Wen Xinhua : 1A8, 2A16
Wen Yuanhui : 1P1
Wenger Jerome : 2A21
Werner Douglas : 1A19
Werner Douglas H. : 2A32
Werner Pingjuan L. : 2A32
Wesemann Lukas : 1A3
West Kenneth : 3A1
Weyher Jan L. : 2P2
White Thomas : 3P1
Whiting Eric B. : 2A32
Whitworth G. L. : 3A14
Wiecha Matthias M. : 3P2
Wiecha Peter : 1A12
Wiecha Peter R. : 2A12, 2A18
Wiederrecht Gary : 1A23
Wiersig Jan : 2A26, 3A4
Wilke Robin Niklas : 1P1
Wilkinson Timothy : 4P1
Williams Calum : 4P1
Williams Robin L : 1A30
Winkler Felix : 3A28
Winkler Pamina M. : 2A21
Wintz Sebastian : 4A31
Withayachumnankul Withawat : 3A8
Wiwatowski Kamil : 4A2
Wolf Sebastian : 3A30
Wolinski Tomasz R. : 4P1
Won Rachel : 1A5
Wong Stephan : 1A11, 1A20, 3A4
Wong Thomas T. Y. : 1A30
Wong Wai Chun : 3A4
Wong Zi Jing : 4A33
Wozniak Pawel : 1P2, 4A9
Wright Oliver B. : 1A3, 1A8
Wu Bei : 3A18
Wu Da-Jian : 3A6
Wu Dajian : 1A9
Wu Fan : 2A33
Wu Hong Wei : 1A8
Wu Hong-Wei : 4P1
Wu Jeong Weon : 4A4
Wu Jingyi : 2A21
Wu Lin : 1A10
Wu Nianqiang : 2A31
Wu Pin Chieh : 1A6
Wu Qian : 1A27
Wu Qun : 2A10, 2A15, 4A5, 4A5
Wu Tianhao : 4A5
Wu Tong : 3A5
Wu Wenbing : 1A5
Wu Ying : 2A8, 2A37
Wunderer Thomas : 3A37
Wätzel Jonas : 3A20
Xenidis Nikolaos : 1A28

Xia Rongyu : 1A27
 Xiang Yuanjiang : 1A11
 Xiao Fajun : 2A10
 Xiao Jialing : 2A34
 Xiao Meng : 3A10
 Xiao Min : 3A7
 Xiao Wei : 3A23
 Xiao Yi-Xin : 2A36
 Xie Changqing : 3A19, 4A6
 Xie Jianliang : 3P1
 Xing Lei : 3A6
 Xiong Daxi : 4A25
 Xiong Kunli : 3A17
 Xiong Xiang : 1A10
 Xiong Zhongfei : 1A11
 Xu Beibei : 4A1
 Xu Chenglin : 2P2
 Xu Hongbo : 4A16
 Xu Hongxing : 1A29
 Xu Jimmy : 1A29, 3P2
 Xu Jingjun : 3A9
 Xu Lei : 2A18
 Xu Ouyang : 3A18
 Xu Ting : 2P1, 3A12
 Xu Yi : 3A18
 Xu Ziquan : 1A3, 4A19
 Xue Bin : 1P2
 Xue Jiancai : 1A10
 Ya'akobovitz Assaf : 4A17
 Yablonovitch Eli : 2A30
 Yagudin Damir : 1A18
 Yamada Takayuki : 1A9
 Yamamoto Naoki : 2A7
 Yamane Hidemasa : 1A7
 Yan Bing : 4A25, 4A27
 Yan Xiang Xi : 3A6
 Yang Biao : 3A20
 Yang Jinghui : 3A12
 Yang Jingyi : 3A32
 Yang Xibin : 4A25
 Yang Xingyu : 4A18
 Yang Yang : 1A20
 Yang Zhengmei : 3A26
 Yang Zih-Ying : 3A3, 3A3
 Yankin Sergey : 4A28
 Yao Jie : 3A6
 Yao Yuanzhao : 2A1
 Yavorski Dmitri : 4P1
 Yazhgur Pavel : 1A14
 Yen Ta-Jen : 3A6, 3A11, 4A15
 Yermakov Oleh : 2A4
 Yeschhenko Oleg : 4A28
 Yeshchenko Oleg : 1P2
 Yeung Cheuk-Him : 4A26
 Yeung Edith : 1A30
 Yi Jianjia : 2A10
 Yin Hongcheng : 3P1
 Yin Xuefan : 3A10
 Yin Zhen : 2A2, 4A11
 Yokoshi Nobuhiko : 1A7, 4P1
 Yokoyama Takahiro : 3A3
 Yokoyama Takehito : 2A6
 Yoo J. H. : 1A25
 You Jang-Woo : 2A7
 You Oubo : 1A20
 You Xiaolong : 3A8
 Yousef Elsayed : 3P2, 3P2
 Yu Binbin : 1A6
 Yu Jaeyeon : 3A12
 Yu Meng-Ju : 1A29
 Yu Peng : 1A4
 Yu Siyuan : 1P1
 Yu Sunkyu : 2A5, 4A14
 Yu Xiaochang : 1P1
 Yu Ye : 3A28
 Yu Yidyu : 4A18
 Yu Yu : 4A9
 Yu Zongfu : 2A32, 2A37, 3A23
 Yuan J. : 3A40
 Yuan Jifeng : 4A7
 Yuan Jun : 3A30
 Yuan Luqi : 3A10
 Yuan Xiaodong : 2A10, 4A17
 Yuan Yueyi : 2A10, 2A15, 4A5
 Yue Liyang : 4A25, 4A27
 Yuen-Zhou Joel : 1A23
 Yumigeta Kentaro : 2A3
 Yun Seokho : 4A1
 Yung Tsz Kit : 4A11
 Yunusa Zainab : 1P2
 Yüce Emre : 3P2
 Zabala Nerea : 4A18
 Zagaglia L. : 4A3
 Zahn Dietrich R.T. : 4A34
 Zahradnik Martin : 1A18
 Zaleska Anastasiia : 2A21
 Zalewski Tomasz : 4A31, 4A31
 Zambrana-Puyalto Xavier : 2A19
 Zapata-Rodriguez Carlos J. : 3P2
 Zarei Sanaz : 2P2
 Zayats Anatoly V. : 2A3, 2A21
 Zebrowski Michal : 4A28
 Zecca Roberto : 2A27
 Zeimpekis Ioannis : 3A23
 Zeisberger Matthias : 2A4
 Zemanek Pavel : 4A28
 Zemp Armin : 1A17
 Zeng Hao : 4A11
 Zentgraf Thomas : 4A11
 Zetterstrom Oskar : 4A5
 Zezyulin Dmitry A. : 2A8
 Zhang Cheng : 3A12
 Zhang D. G. : 3A11
 Zhang Dai : 3A1
 Zhang Dawei : 1A6
 Zhang Fan : 2A9

Zhang Guoquan : 3A9
Zhang Hailin : 2A10
Zhang Haizhong : 3A34
Zhang Haoyang : 2A18
Zhang Hongkuan : 1A27
Zhang Hualiang : 2A27
Zhang Jianfa : 2A10, 4A17
Zhang Jianhao : 3A21
Zhang Kuang : 2A10, 2A15, 4A5, 4A5
Zhang Li : 3A9
Zhang Likun : 4A13
Zhang Ming : 2A7
Zhang Qi : 2A9, 4A18
Zhang Ruo-Yang : 1A11, 3A20
Zhang Shuang : 1P1, 1A11, 1A20, 1A20
Zhang Wei : 1A10
Zhang Wending : 3A19
Zhang Xiao : 1A11
Zhang Xiaomeng : 3A19
Zhang Xin : 3A34
Zhang Xuanru : 4A9
Zhang Yan : 4A20
Zhang Yu : 2A31, 3A22
Zhang Zhao-Qing : 2A36
Zhang Zhiwang : 1A9, 1A9
Zhang Zhongwei : 1A8
Zhao Jianlin : 2A10
Zhao Meiyang : 1A8
Zhao Yingqi : 2A28
Zheludev Nikolay I. : 4A30
Zhen Bo : 3A10
Zheng Bowen : 2A27
Zheng Changxiong : 4A11
Zheng Di : 4A28
Zheng Liyang : 1A9
Zheng Xu : 2P2
Zheng Xuezhi : 4A9
Zheng Yuanlin : 3A9
Zhou Changyi : 4A15
Zhou Chenzhang : 2A25
Zhou Jing : 3A19, 4A9
Zhou Jun : 4A29
Zhou Ming : 3A23
Zhou Ning : 1A30
Zhou Wei : 3A34
Zhou Xiaoming : 1A27
Zhou Yi : 2A11, 3A6
Zhou Zhang-Kai : 1A10
Zhu Bo O. : 2A1
Zhu Huanzheng : 1A3, 4A19
Zhu Jiangbo : 1P1
Zhu Jie : 2A16
Zhu Jinlong : 4A27
Zhu S. N. : 3A19
Zhu Shining : 3A7, 4A1
Zhu Weiren : 2A10
Zhu Wenqi : 3A12
Zhu Xing-Feng : 3A6
Zhu Xinghong : 1A8, 2A8, 2A16
Zhu Xuefeng : 4A13
Zhu Yi-fan : 2A14
Zhu Yifan : 4A13
Zhu Yong : 3P1
Zhu Zhihong : 2A10, 4A17
Zhukova Elena : 4A2
Zhuo Xiaolu : 1A10
Zilk Matthias : 1A24
Zilli Attilio : 3A5
Zinoviev Alexey : 3A28
Zito Gianluigi : 4A35
Zograf George : 3P1
Zohar Moshe : 4P1
Zolfaghary Saeed : 2P1
Zopf David : 4A29
Zou Jinglan : 2A10
Zschiedrich Lin : 2A17
Zubair Muhammad : 4A24
Zubov Yurii : 1A27
Zvagelsky Roman : 4A2
Zyablovsky Alexander : 4A16
Zyss Joseph : 4A29
Šikola Tomáš : 4P1, 4A15